

Global Ocean Monitoring: Recent Evolution, Current Status, and Predictions

Prepared by
Climate Prediction Center, NCEP/NOAA

November 9, 2023

<http://www.cpc.ncep.noaa.gov/products/GODAS/>

This project, to deliver real-time ocean monitoring products, is implemented
by CPC in cooperation with NOAA's Global Ocean Monitoring and Observing Program (GOMO)



- Overview
- Recent highlights
 - Pacific Ocean
 - Arctic & Antarctic Oceans
 - Indian Ocean
 - Atlantic Ocean
- Global SSTA Predictions

• Pacific Ocean

- El Niño conditions continued to grow in Oct 2023.
- The strong warming in the far eastern Pacific weakened in Oct 2023.
- NOAA “ENSO Diagnostic Discussion” on 9 Nov 2023 *stated “El Niño is anticipated to continue through the Northern Hemisphere spring (with a 62% chance during April-June 2024)”*.
- The PDO has been in a negative phase since Jan 2020 with PDOI = -1.7 in Oct 2023.
- Strong subsurface warming has persisted in the central north Pacific Ocean since 2020.

• Arctic and Antarctic Oceans

- Average Arctic sea ice extent during Oct 2023 ranked the seventh lowest Oct since 1979.
- Antarctic sea ice extent remained at historical low level.

• Indian Ocean

- A positive Indian dipole event continued to develop in Oct 2023.


• Atlantic Ocean

- SST warming in the north Atlantic during Aug-Oct 2023 hit the historical high for the same season since 1982.
- Hurricane activity was active in Oct 2023.
- Marine heatwaves continued in west coast of North Africa, Caribbean and Labrador basin.


Global Oceans

Global SST Anomaly (°C) and Anomaly Tendency

OCT 2023 SST Anomaly (°C)
(1991–2020 Climatology)



OCT 2023 – SEP 2023 SST Anomaly (°C)





- SSTs were above average across most of the equatorial Pacific Ocean.
- Strong coastal El Niño condition weakened in Oct 2023.
- Positive SSTAs persisted in the North Pacific.
- Positive SSTAs dominated the Atlantic Ocean.
- Negative (positive) SSTA were present in the eastern (western) Indian Ocean, reflecting a grow positive Indian dipole event.

- Both negative and positive SSTA tendencies were observed in the North Pacific Ocean.
- Negative SSTA tendencies dominated the North Atlantic Ocean.
- Negative (positive) SSTA tendencies were present in the eastern (far western) equatorial Indian Ocean.

SSTAs (top) and SSTA tendency (bottom). Data are derived from the Olv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.

Global Monthly SST anomaly and Marine Heat Waves



OISSTv2.1 OCT2023 SST Anom. ($^{\circ}\text{C}$)
Hatch area: MHW on OCT-2023-31



- MHWs were observed in the Northeast subtropical Pacific, Northwest Pacific, central-eastern equatorial Pacific, Northeast Atlantic, Caribbean Sea, Labrador basin and the western Indian Ocean.

((Left panel) Monthly SST anomaly (shaded) and locations experiencing marine heat waves (hatched) by the end date labelled in the plot. (right panel) SST evolution at a specific location. Green line and blue line are the 90th percentile and daily climatology, respectively. Shaded area denotes the periods experiencing MHW. MHW is defined as a prolonged warming exceeding 90th percentile of daily SST for at least 14 consecutive days. Data is derived from NCEI OISSTv2.1 and the reference period is 1991-2020

Longitude-Depth Temperature Anomaly and Anomaly Tendency in 2°S-2°N




- Positive temperature anomalies were present along the central-eastern thermocline in the Pacific.
- Negative(positive) temperature anomalies were observed along the eastern (western) thermocline in the Indian Ocean, favoring a positive IOD development.
- Negative temperature anomaly dominated the thermocline in the Atlantic Ocean.

- Negative (positive) temperature anomaly tendency were observed in the western and eastern (central) thermocline in the Pacific.
- Dipole pattern of anomaly tendency continued in the Indian Ocean.









Equatorial depth-longitude section of ocean temperature anomalies (top) and anomaly tendency (bottom). Data is from the NCEP's GODAS. Anomalies are departures from the 1991-2020 base period means.

Tropical Pacific Ocean and ENSO Conditions




SSTAs (top-left), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right; positive means heat into the ocean), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the OIv2.1 SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.

Westward Expansion & Evolution of Coastal El Niño



Equatorial Pacific SST ($^{\circ}\text{C}$), HC300 ($^{\circ}\text{C}$), u850 (m/s) Anomalies


2 $^{\circ}\text{S}$ –2 $^{\circ}\text{N}$ Average, 3 Pentad Running Mean



- Since Feb 2023, a set of westerly wind surges triggered downwelling Kelvin waves, helping to reinforce the subsurface warming in the central-eastern Pacific.
- Westerly wind anomalies prevailed over most of equatorial Pacific Ocean in Oct 2023.
- Positive SST anomalies strengthened in the central and eastern Pacific in Oct 2023.

Evolution of Equatorial Pacific Surface Zonal Current Anomaly (cm/s)


U (15m), cm/s, 2°S–2°N (Shading=Anomaly; Contour=1993–2020 Clim)




- Anomalous eastward currents were present in the equatorial Pacific in both OSCAR and GODAS since Feb 2023, which were consistent with the growth of the positive SSTA.
- Anomalous eastward currents enhanced west of 150W in Oct 2023.

Multiple Ocean Reanalysis Intercomparison: Temperature Anomaly and Tendency at Equator

Anomalous Temperature (C) Averaged in 1S-1N: OCT 2023




OCT 2023 - SEP 2023 1S-1N Temp Anomaly (C)





The monthly subsurface temperature data is obtained from the Real-time Ocean Reanalysis Intercomparison Project (https://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html).

Oceanic ENSO Precursors: WWV & CTP



Warm water volume (WWV) is defined as an average of D20 anomaly across the equatorial Pacific (120° E – 80° W, 5° S-5° N) (Meinen and McPhaden 2000). Central tropical Pacific (CTP) index is calculated as the averaged D20 anomaly in the central tropical Pacific (160° W-110° W, 10° S-10° N) (Wen et al. 2014). The monthly D20 data is obtained from the Real-time Ocean Reanalysis Intercomparison Project (https://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html).

Evolution of Pacific Niño SST Indices





- Niño 4 and Niño3.4 indices strengthened in Oct 2023, with Niño3.4 = 1.6°C.
- Positive Niño1+2 weakened in Oct 2023, with Niño1+2 = 2.5°C.
- Compared with Oct 2022, the tropical Pacific was much warmer in Oct 2023.
- The indices may have differences if based on different SST products.

Niño region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the specified region. Data are derived from the Olv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.

Evolution of Pacific Niño SST Indices

Monthly Tropical Pacific SST Anomaly




- Relative Niño3.4 index is now included in ENSO monitoring, which is defined as the conventional Niño3.4 index minus the SSTA averaged in the whole tropics (0°-360°, 20°S-20°N), in order to remove the global warming signal. Also, to have the same variability as the conventional Niño3.4 index, the relative Niño3.4 index is renormalized (van Oldenborgh et al. 2021: ERL, 10.1088/1748-9326/abe9ed).

[Relative Niño3.4 data updated monthly at: https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt](https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt)

2023 Pacific Hurricane Season Activities



NOAA's 2023 Hurricane Season Outlooks



For the Eastern Pacific hurricane region, the outlooks indicate a 55% chance of an above-normal season, a 35% chance of a near-normal season, and a 10% chance of an above-normal season. The odds for the Central Pacific are 50% for an above-normal season, 35% for a near-normal season, and 15% for a below-normal season.

These outlooks are for the overall seasonal activity. They are not a hurricane landfall forecast.









https://en.wikipedia.org/wiki/2023_Pacific_hurricane_season

E. Pacific	Observations (By Nov 6)	Outlook (May 25) 55% above-normal	(1991-2020)
Total storms	16	14-20	15
Hurricanes	10	7-11	8
Major hurricanes	8	4-8	4



North Pacific & Arctic Oceans

Last 3- month North Pacific SST, SLP, and uv925 anomalies





Subsurface Temperature Anomaly in the Northcentral Pacific

Anomalous Temperature (C) in [160E-150W, 30N-45N]




OCT 2023 SST Anom. (°C)




- Positive temperature anomaly ($>0.9^{\circ}\text{C}$) penetrated to 150m deep and persisted since 2020.
- Subsurface warming in the last three years is the strongest since 1979.

Two Oceanic PDO indices

SST-based PDO (Wen et al. 2014: GRL)



H300-based PDO (Arun and Wen 2016: Mon. Wea. Rev.)



- The negative phase of PDO has persisted since Jan 2020 with PDOI = -1.7 in Oct 2023.

- Negative H300-based PDO index has persisted since Nov 2016, with HPDO = -1.8 in Oct 2023.


- SST-based PDO index has considerable variability both on seasonal and decadal time scales.

- H300-based PDO index highlights the slower variability and encapsulates an integrated view of temperature variability in the upper ocean.


SST-based PDO is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly ERSSTv5 SST anomalies onto the 1st EOF pattern. H300-based Pacific Decadal Oscillation is defined as the projection of monthly mean H300 anomalies from NCEP GODAS onto their first EOF vector in the North Pacific. PDO indices are downloadable from https://www.cpc.ncep.noaa.gov/products/GODAS/ocean_briefing.shtml.

North America Western Coastal Upwelling

Pentad Coastal Upwelling for West Coast North America
($\text{m}^3/\text{s}/100\text{m}$ coastline)



Standard Positions of Upwelling Index Calculations



- Anomalous upwelling dominated along the coastline in the second-half of Oct 2023.


(top) Total and (bottom) anomalous upwelling indices at the 15 standard locations for the western coast of North America. Derived from the vertical velocity of the NCEP's GODAS and are calculated as integrated vertical volume transport at 50-meter depth from each location to its nearest coast point ($\text{m}^3/\text{s}/100\text{m}$ coastline). Anomalies are departures from the 1991-2020 base period pentad means.

- Area below (above) black line indicates climatological upwelling (downwelling) season.

- Climatologically upwelling season progresses from March to July along the west coast of North America from 36°N to 57°N.




Weekly SST anomaly and MHWs in the North Pacific

Weekly OISSTv2.1 Anom. (°C)
Hatch area: MHW location





- MHW persisted in the northwest Pacific Ocean in Oct 2023.
- MHWs were observed in the northeastern tropical Pacific in Oct 2023.
- Considerable amount of anomalously warm waters persisted near the coast of California.

(Left panel) Weekly SST anomaly (shaded) and locations experience Marine heat waves (hatched) by the date labelled in the plot. (right panel) SST evolution at a specific location. Green line and blue line denote the seasonal 90th percentile and daily climatology, respectively. Shaded area denotes the periods experiencing MHW. MHW is defined as a discrete prolonged warmer than 90th percentile of daily SST for at least 14 days. Data is derived from NCEI OISSTv2.1 and the climatology reference period is 1991-2020




- Average Arctic sea ice extent during Oct 2023 was 6.37 million square kilometers, the seventh lowest Oct in the satellite record.



-Antarctic sea ice extent was 15.79 million square kilometers on Oct 31, remaining at record low levels.

UFS

Arctic sea ice extent (SIE) forecast
Experimental UFS initialized October 21–25, 2023





- UFS forecasts suggest SIE maximum in Mar 2023 will be close to 1991-2020 climatology.

https://www.cpc.ncep.noaa.gov/products/people/jszhu/seaiice_seasonal/index.html


Indian Ocean

Evolution of Indian Ocean SST Indices



- Indian dipole mode index (DMI) has been greater than 0.4°C since Aug 2023. It indicated a positive IOD event is underway.
- Positive Indian dipole mode index (DMI) strengthened in Oct 2023, with DMI= 1.43 °C.

Indian Ocean region indices, calculated as the area-averaged monthly mean SSTA (°C) for the SETIO [90°E–110°E, 10°S–0] and WTIO [50°E–70°E, 10°S–10°N] regions, and Dipole Mode Index, defined as differences between WTIO and SETIO. Data are derived from the OIv2.1 SST analysis, and anomalies are departures from the 1991–2020 base period means.




SSTAs (top-left), SSTA tendency (top-right), OLR anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the Olv2.1 SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.


Tropical and North Atlantic Ocean

Evolution of Tropical Atlantic SST Indices


Monthly Tropical Atlantic SST Anomaly
(OISSTv2.1, 1991–2020 Climatology)



OCT2023 SST Anom. (°C)




OCT2023 – OCT2022 SST Anom. (°C)



- Strong positive SSTA in the tropical north Atlantic weakened slightly in Oct 2023.
- Positive ATL3 index weakened in Oct 2023.

Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the TNA [60°W-30°W, 5°N-20°N], TSA [30°W-10°E, 20°S-0] and ATL3 [20°W-0, 2.5°S-2.5°N] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the OIv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.





Tropical Atlantic: SST, SST tend., TCHP, OLR, 200 hPa wind, wind share, heat flex, & RH anom.



Top Row: SSTA (left; OI SST), SSTA tendency (central), Tropical Cyclone Heat Potential anomaly (right; GODAS).
 Middle row: OLR (left; NOAA 18 AVHRR IR), UV200 (central; NCEP CDAS), UV200-UV850 (right; NCEP CDAS) anomalies.
 Bottom row: SW+LW (left), LH+SH (central), Relative humidity at 700 hPa (right; NCEP CDAS) anomalies.
 Anomalies are departures from the 1991-2020 base period means.


Evolution of SST anomaly in the North Atlantic

OISSTv2.1 Anomaly (°C) : AUG-OCT 2023






Evolution of 200mb-850mb Wind Shear Anomaly

200mb - 850mb Wind Shear Anom. (m/s): AUG-OCT 2023



- Statistically, Wind shear tends to enhance (weaken) over the Hurricane Main Development region (MDR, green box) during El Niño (La Niña) events.
- Negative wind shear anomalies dominated in the MDR during Aug-Oct 2023, favouring tropical storm development.





- NOAA forecasters have increased the likelihood of an above-normal Atlantic hurricane season from **30%** in outlook issued in May to **60%** in August outlook update.
- Main climate factors expected to influence the 2023 Atlantic hurricane activity are : the ongoing El Niño, the warm phase of the Atlantic Multi-Decadal Oscillation and record-warm Atlantic SSTs.
- **Likelihood of greater activity rises due to record-warm Atlantic SST .**
(<https://www.noaa.gov/news-release/noaa-forecasters-increase-atlantic-hurricane-season-prediction-to-above-normal>)

2023 Atlantic Hurricane Season Activities





https://en.wikipedia.org/wiki/2023_Atlantic_hurricane_season

- By 6 Nov 2023, twenty tropical storms formed, with seven developing into hurricane and three major hurricanes.

Atlantic	Observations (By Nov 6)	Updated Outlook (Aug) 60% above-normal	Outlook (May 25) 40% near-normal	(1991-2020)
Total storms	20	14-21	12-17	14
Hurricanes	7	6-11	5-9	7
Major hurricanes	3	2-5	1-4	3

Weekly SST anomaly and MHWs in the North Atlantic




Weekly OISSTv2.1 Anom. (°C)
Hatch area: MHW location



- Strong MHWs has been persist near the west coast of North Africa since Apr 2023.
- MHWs continued near the Caribbean and the Labrador basin.

(Left panel) Weekly SST anomaly (shaded) and locations experience Marine heat waves (hatched) by the date labelled in the plot. (right panel) SST evolution at a specific location. Green line and blue line denote the seasonal 90th percentile and daily climatology, respectively. Shaded area denotes the periods experiencing MHW. MHW is defined as a discrete prolonged warmer than 90th percentile of daily SST for at least 14 days. Data is derived from NCEI OISSTv2.1 and the climatology reference period is 1991-2020

NAO and SST Anomaly in North Atlantic




- Negative NAO strengthened substantially in Oct 2023, with NAO = -1.7.
- Strong warming continued in the eastern North Atlantic Ocean.


Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N. Time-latitude section of SSTAs averaged between 80°W and 20°W (bottom). SST are derived from the OlV2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.

ENSO and Global SST Predictions

CFSv2 IC:Nov for DJF 2023




NMME IC:Nov for DJF 2023




NWS/NCEP/CPC

Last update: Tue Nov 7 2023
Initial conditions: 28Oct2023-6Nov2023

CFSv2 forecast Nino3 SST anomalies (K)



NMME Nino3.4 Fcst, IC=202311





<https://www.cpc.ncep.noaa.gov/products/CFSv2/CFSv2seasonal.shtml>


<https://www.cpc.ncep.noaa.gov/products/NMME/>

Uncertainty in ocean initial conditions and NMME forecasts


Anomalous Temperature (C) Averaged in 1S-1N: OCT 2023




CFSv2 and CCSM4 forecasts at different initial months



Both CFSv2 and CCSM4 use CFSR as ocean initial conditions.



Impact of NSST cold bias on CFSv2




Bias pathway : NSST \Rightarrow CFSR \Rightarrow CFSv2

Wen,C., A. Kumar, W. Wang, M.L Heureux, P.Xie, Z. Hu and B.Katz (2022): Communicating uncertainty in SST analysis. Extended Summary, *Climate Prediction S&T Digest*, 46th NOAA Climate Diagnostics and Prediction Workshop, Virtual Online, DOC/NOAA, page range. DOI: 10.25923/rj6c-rk11


Individual Model Forecasts: A strong to very strong El Niño in 2023

EC: Niño3.4, IC= 1 Nov 2023




ECMWF

UKMO: Niño3.4, Updated 11 Oct 2023




BOM: Niño3.4, Updated 21 Oct 2023



www.bom.gov.au/climate
Commonwealth of Australia 2023, Australian Bureau of Meteorology




Model: ACCESS-S2
Base period 1981-2018
Model run: 21 Oct 2023

JMA: Niño3.4, Updated 11 Oct 2023





All rights reserved. Copyright(c)Japan Meteorological Agency

IRI/CPC Niño3.4 Forecast




- Most of models forecasted that El Niño conditions will continue through the Northern Hemisphere spring 2024.
- Dynamical model ensemble mean favors of a strong El Niño through Jan-Mar 2023.
- **ENSO Alert System Status issued on Nov 9 2023: El Niño Advisory**
- Synopsis: "El Niño is anticipated to continue through the Northern Hemisphere spring (with a 62% chance during April-June 2024) "

Evolution of NINO34 & NINO1+2 in El Niño Years



Comparisons among 1982, 1997 and 2023 El Niño events




Nino 3.4 and NINO1+2 indices are derived from the NCEP OI SST analysis, and anomalies are departures from the 1991-2020 base period means

NMME Forecasts in the Indian Ocean


NMME Sea Surface Temperature Anomalies (DecC)

Dec2023–Feb2024 November2023 initial conditions



https://www.cpc.ncep.noaa.gov/products/international/ocean_monitoring/indian/IO_monitoring_fcsts/io_index.shtml




NMME IOD fcst, IC=202311




- All NMME members predict the positive IOD event will last through Feb 2024.

NMME Forecasts in the Atlantic Ocean

NMME Sea Surface Temperature Anomalies (DecC)




standardized PDO index



- CFSv2 predicts the negative phase of PDO will continue through spring 2024.

CFS Pacific Decadal Oscillation (PDO) index predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means. PDO is the first EOF of monthly ERSSTv3b anomaly in the region of [110°E-100°W, 20°N-60°N]. CFS PDO index is the standardized projection of CFS SST forecast anomalies onto the PDO EOF pattern.



- NMME forecasts suggest that MHW coverage will remain elevated near 30% of the global oceans through the end of 2023.
- MHW conditions are expected to persist in the eastern tropical Pacific, central North Pacific, Northwest Pacific near Japan and tropical North Atlantic through the end of year.
- MHW condition will persist in the Caribbean Sea through the end of 2023 with 70-100% probability.

Acknowledgement

- ❖ Drs. Arun Kumar, Zeng-Zhen Hu, Jieshun Zhu reviewed PPT, and provide insightful suggestions and comments
- ❖ Dr. Pingping Xie provided the BASS/CMORPH/CFSR EVAP package
- ❖ Drs. Jieshun Zhu and Wanqiu Wang provided the upgraded sea ice forecasts

Please send your comments and suggestions to:

Arun.Kumar@noaa.gov

Caihong.Wen@noaa.gov

Jieshun.Zhu@noaa.gov

Zeng-Zhen.Hu@noaa.gov

- **NCEP/CPC Ocean Monitoring & Briefing Operation (Hu et al., 2022, BAMS)**
- **Weekly Optimal Interpolation SST (OIv2.1 SST; Huang et al. 2021)**
- **Extended Reconstructed SST (ERSST) v5 (Huang et al. 2017)**
- **Blended Analysis of Surface Salinity (BASS) (Xie et al. 2014)**
- **CMORPH precipitation (Xie et al. 2017)**
- **CFSR evaporation adjusted to OAFlux (Xie and Ren 2018)**
- **NCEP CDAS winds, surface radiation and heat fluxes (Kalnay et al. 1996)**
- **NESDIS Outgoing Long-wave Radiation (Liebmann and Smith 1996)**
- **NCEP's GODAS temperature, heat content, currents (Behringer and Xue 2004)**
- **Aviso altimetry sea surface height from CMEMS**
- **Ocean Surface Current Analyses – Realtime (OSCAR)**
- **In situ data objective analyses (IPRC, Scripps, EN4.2.1, PMEL TAO)**
- **Operational Ocean Reanalysis Intercomparison Project**
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora_body.html
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html

Backup Slides

Global Sea Surface Salinity (SSS): Anomaly for October 2023

New Update: The NCEI SST data used in the quality control procedure has been updated to version 2.1 since May 2020;


Enhanced fresh water flux (driven by strong precipitation) continued during October 2023, across the equatorial Pacific and Atlantic, creating freshened SSS anomalies there. Strong convection shifted northward over the tropical Indian Ocean, resulting wet / dry precipitation anomalies there. SSS is freshened over regions of wet precipitation, but maintains weak negative anomaly over regions of dry precipitation anomalies, likely attributable to other oceanic processes (e.g. advection). Saltier SSS anomalies are also observed over the Bay of Bengal and the Gulf of Mexico.

**SSS : Blended Analysis of Surface Salinity (BASS) V0.Z
(a CPC-NESDIS/NODC-NESDIS/STAR joint effort)**


<ftp.cpc.ncep.noaa.gov/precip/BASS>

Precipitation: CMORPH adjusted satellite precipitation estimates

Evaporation: Adjusted CFS Reanalysis




Precipitation and ocean-going fresh water flux anomalies enhanced substantially over the tropical eastern Pacific off the coast of central America, resulting in freshening SSS tendency there. Saltier SSS tendencies, meanwhile, are observed over the Bay of Bengal, consistent with the tendencies of Precipitation and E-P.



Monthly SSS Anomaly Evolution over Equatorial Pacific

NOTE: Since June 2015, the BASS SSS is from in situ, SMOS and SMAP; before June 2015, The BASS SSS is from in situ, SMOS and Aquarius.


- Hovermoller diagram for equatorial SSS anomaly (5°S - 5°N);
- Freshened SSS anomalies over the western and central equatorial Pacific (130°E - 170°W) continued and are further enhanced as a result of the strong ITCZ activities during October 2023. Saltier SSS anomalies over the eastern equatorial Pacific continues weakening.



Pentad SSS Anomaly Evolution over Equatorial Pacific


Figure caption:

Hovermoller diagram for equatorial (5°S - 5°N) 5-day mean SSS, SST and precipitation anomalies. The climatology for SSS is Levitus 1994 climatology. The SST data used here is the OISST V2 AVHRR only daily dataset with its climatology being calculated from 1985 to 2010. The precipitation data used here is the adjusted CMORPH dataset with its climatology being calculated from 1999 to 2013.






AVISO & GODAS SSH Anomaly (cm) and Anomaly Tendency

OCT 2023 SSH Anomaly (cm)
(climo. 1993–2020)




OCT 2023 – SEP 2023 SSH Anomaly (cm)
(limo. 1993–2020)




- SSHs were above normal in the equatorial Pacific in GODAS & AVISO.
- The tendencies indicated an increase (decrease) of SSH in the eastern (western) tropical Pacific.

Evolution of Monthly Mean Zonal Wind Stress Anomaly across [5S-5N]





Monthly Mean Taux(shaded) & SST(contour) Anomaly(5S-5N: 1991-2020 Climatology)



Evolution of Monthly Mean D20 Anomaly across [2S-2N]



Pacific Decadal Oscillation (PDO) Index









- The PDO has been in a negative phase since Jan 2020 with PDOI = -1.9 in Jul 2023.

- Statistically, ENSO leads PDO by 3-4 months, through teleconnection via atmospheric bridge, with El Niño (La Niña) associated with positive (negative) PDO Index.

• PDO is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly SST anomalies onto the 1st EOF pattern.


North Pacific & Arctic Ocean: SSTA, SSTA Tend., OLR, SLP, Sfc Rad, Sfc Flx Anomalies



SSTA (top-left; Olv2.1 SST Analysis), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) (middle-left; NOAA 18 AVHRR IR), sea surface pressure (middle-right; NCEP CDAS), sum of net surface short- and long-wave radiation (bottom-left; positive means heat into the ocean; NCEP CDAS), sum of latent and sensible heat flux (bottom-right; positive means heat into the ocean; NCEP CDAS). Anomalies are departures from the 1991-2020 base period means.

Hurricane-friendly Climate Conditions

Hurricane-friendly climate conditions during “active” eras: warm phase of AMO









(Gerry Bell, 2014)


<https://www.climate.gov/news-features/blogs/enso/impacts-el-ni%C3%B1o-and-la-ni%C3%B1a-hurricane-season>

Established theories:

- Warm phase of Atlantic Multi-decadal Oscillation (AMO)
- Warmer SSTs across the Atlantic hurricane main development region
- Reduced wind shear (i.e ENSO impact)
- Stronger West African monsoon




Niño3.4 SST anomalies (K)



- The latest CFSv2 forecasts call for an El Niño in the second half of 2023.

CFS Niño3.4 SST prediction from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means.

Indian Ocean Dipole SST anomalies (K)



- CFSv2 predicts a positive phase of IOD in the 2nd half of 2023.

CFS Dipole Model Index (DMI) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1991-2020 base period means.

NMME forecasts from different initial conditions

