

# **Global Ocean Monitoring: Recent Evolution, Current Status, and Predictions**

Prepared by  
Climate Prediction Center, NCEP  
**November 5, 2010**

**<http://www.cpc.ncep.noaa.gov/products/GODAS/>**

**This project to deliver real-time ocean monitoring products is implemented  
by CPC in cooperation with NOAA's Office of Climate Observation (OCO)**

# Outline

- **Overview**
- **Recent highlights**
  - **Pacific/Arctic Ocean**
  - **Indian Ocean**
  - **Atlantic Ocean**
- **CFS SST Predictions**

# Overview

- **Pacific/Arctic Ocean**

- ENSO cycle: La Niña conditions persisted with  $\text{NINO3.4} = -1.6^\circ\text{C}$  in Oct 2010.
- NOAA/NCEP Climate Forecast System (CFS) predicted a strong La Niña, to last through the Northern Hemisphere spring 2011.
- PDO has been below-normal since Jul 2010, with  $\text{PDOI} = -1.2$  in Oct 2010.
- Arctic sea ice extent increased seasonally after mid-Sep 2010, and was still well below-normal.


- **Indian Ocean**

- The tropical Indian Ocean Basin warming weakened in the north and slightly strengthened in the south in Oct 2010.
- Dipole Mode index has been below-normal since May 2010, and strengthened to be about  $-0.9^\circ\text{C}$  in Oct 2010.


- **Atlantic Ocean**

- NAO index has been persistently below-normal since Oct 2009, and it was  $-0.9$  in Oct 2010.
- Strong positive SSTA ( $>2.5^\circ\text{C}$ ) presented in the high latitudes since Sep 2010.
- SST in the tropical North Atlantic (TNA) has increased steadily from Dec 2009 to May 2010, gradually weakened from Jun to Aug 2010, and persisted in Sep-Oct 2010.
- The active hurricane season in 2010 was well predicted by NOAA. Multiple factors, including the current La Nina, the delayed impacts of the 2009/10 El Nino, persistent negative NAO, and decadal trend, seem all have contributed to the above normal hurricane activity.

# Global SST Anomaly ( $^{\circ}\text{C}$ ) and Anomaly Tendency




- Negative SSTA in the tropical eastern and central Pacific, indicating the La Niña conditions.
- Negative phase PDO SSTA pattern presented in N. Pacific.
- Positive SSTA existed in the tropical Indian Ocean and tropical W. Pacific.
- Strong positive SSTA in the high latitude North Atlantic, Arctic, and tropical Atlantic.



- Small SSTA tendency in the west-central tropical Pacific and large negative SSTA tendency in the eastern tropical Pacific suggested a persistency in the central and an enhancement in the eastern tropical Pacific for the La Niña associated negative SSTA.
- SST tendency was large in N. Pacific and Arctic.
- Positive (negative) SSTA tendency presented in the middle (high) latitudes of N. Atlantic.

**Fig. G1.** Sea surface temperature anomalies (top) and anomaly tendency (bottom). Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.

# Global SSH/HC Anomaly (cm/°C ) and Anomaly Tendency




- Negative (positive) SSHA and HCA persisted along the Equator in the central-eastern (western) tropical Pacific, consistent with status of La Nina conditions.
- Positive (negative) HCA and SSHA strengthened in the eastern (central) tropical Indian Ocean, consistent with strengthening of negative Indian Ocean Dipole (IOD).
- The tripole SSHA and HCA pattern in North Atlantic persisted.
- SSHA and HCA anomalies as well as their tendencies were largely consistent, except in the Southern Ocean where biases in GODAS climatology are large (not shown).

**Fig. G2. Sea surface height anomalies (SSHA, top left), SSHA tendency (bottom left), top 300m heat content anomalies (HCA, top right), and HCA tendency (bottom right). SSHA are derived from <http://www.aviso.oceanobs.com>, and HCA from GODAS.**

# Regional SSH Anomaly (cm)

Monthly SSH Time Series (Aviso Altimetry, Clima. 1993–2005)




- Positive SSHA persisted or slightly weakened for global oceans, tropical oceans, and Southern Ocean.

- Positive SSHA slightly strengthened in North Pacific and North Atlantic in October.

- Negative SSHA intensified in the equatorial Pacific since Jun 2010, consistent with occurrence of La Nina conditions.

# Longitude-Depth Temperature Anomaly and Anomaly Tendency in 2°S-2°N

OCT 2010 Eq. Temp Anomaly (°C)  
(GODAS, Climo. 82-04)




- Negative (positive) subsurface ocean temperature anomalies dominated in the equatorial east-central (western) Pacific, consistent with the La Niña conditions.

- Negative (positive) subsurface ocean temperature anomalies dominated in the equatorial western (eastern) Indian Ocean, consistent with the negative IOD.

- Positive temperature anomalies dominated near the thermocline of the Atlantic Ocean.

OCT 2010 - SEP 2010 Eq. Temp Anomaly (°C)



- Positive (negative) temperature anomaly tendency presented near the thermocline of the western (eastern) Pacific.


- Subsurface temperature anomalies associated with the negative IOD intensified.

- Strong positive ocean temperature anomaly tendency existed in the tropical Atlantic Ocean.

**Fig. G3. Equatorial depth-longitude section of ocean temperature anomalies (top) and anomaly tendency (bottom). Data are derived from the NCEP's global ocean data assimilation system which assimilates oceanic observations into an oceanic GCM. Anomalies are departures from the 1982-2004 base period means.**

Tropical Storm Tracks

Year 2010





**Tropical Cyclone Activity in Atlantic Jun-Oct 2010: Active year and well predicted by NOAA**

Category	Observed Number (Jun-Oct)	NOAA Prediction (Aug 5, 2010)	Mean (1951-2000)
Major HR (5+4+3)	5 (0+4+1)	4-6	2
Total HR	12	8-12	6
Total Named TS	21	14-20	10
Accumulated Cyclone Energy	155	170-260	93.2



# Tropical Pacific Ocean

# Evolution of Pacific NINO SST Indices




- Strengthened for the Indices in the east (Nino3 and Nino1+2) and slightly weakened in the west (Nino4) in Oct 2010, Consist with SSTA tendency in Slide 4.

- The indices were calculated based on OISST. They may have some differences compared with those based on ERSST.v3b.

**Fig. P1a. Nino region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the specified region. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.**

# NINO3.4 Heat Budget



- Tendency ( $dT/dt$ ) in NINO 3.4 became positive in Oct 2010, probably implying weakening tendency of the cooling.

- All dynamical terms ( $Q_u$ ,  $Q_v$ ,  $Q_w+Q_{zz}$ ) were still negative.

- The thermodynamic processes ( $Q_q$ ) were small positive since Jun 2010.

- There was a clear difference between the tendency ( $dT/dt$ ) and the total budget term (RHS) since Jun 2010. This may be due to the fact that (1) the cooling from  $Q_u$  and  $Q_v$  might be overestimated in the GODAS due to too strong zonal and meridional current, and (2) surface heat flux ( $Q_q$ ) damping was too weak.

$Q_u$ : Zonal advection;  $Q_v$ : Meridional advection;

$Q_w$ : Vertical entrainment;  $Q_{zz}$ : Vertical diffusion

$Q_q$ :  $(Q_{net} - Q_{open} + Q_{corr})/pcph$ ;  $Q_{net} = SW + LW + LH + SH$ ;


$Q_{open}$ : SW penetration;  $Q_{corr}$ : Flux correction due to relaxation to OI SST

# Warm Water Volume (WWV) and NINO3.4 Anomalies

-WWV is defined as average of depth of 20°C in [120°E-80°W, 5°S-5°N] (Meinen and McPhaden, 2000).

-Since WWV is intimately linked to ENSO variability (Wyrтки 1985; Jin 1997), it is useful to monitor ENSO in a phase space of WWV and NINO3.4 (Kessler 2002).

- Increase (decrease) of WWV indicates recharge (discharge) of the equatorial oceanic heat content.




- Negative Nino3.4 persisted in Oct 2010 and became less than -1C since July 2010, indicating moderate-strong La Nina conditions.

- Negative WWV persisted since Jul 2010.

Fig. P3. Phase diagram of Warm Water Volume (WWV) and NINO 3.4 SST anomalies. WWV is the average of depth of 20°C in [120°E-80°W, 5°S-5°N] calculated with the NCEP's global ocean data assimilation system. Anomalies for WWV (NINO 3.4) are departures from the 1982-2004 (1971-2000) base period means.

# Evolution of Equatorial Pacific SST ( $^{\circ}\text{C}$ ), 0-300m Heat Content ( $^{\circ}\text{C}$ ),

## 850-mb Zonal Wind (m/s), and OLR ( $\text{W}/\text{m}^2$ ) Anomaly




- Negative SSTA and HC300A persisted, consistent with the occurrence of La Niña conditions.
- Convection was suppressed in the equatorial eastern Indian Ocean since mid of Oct and in the equatorial central Pacific since May 2010, consistent with near surface wind convergence and divergence.

**Fig. P4.** Time-longitude section of anomalous pentad sea surface temperature (left), upper 300m temperature average (heat content, middle-left), 850-mb zonal wind (U850, middle-right) averaged in 2°S-2°N and Outgoing Long-wave Radiation (OLR, right) averaged in 5°S-5°N. SST is derived from the NCEP OI SST, heat content from the NCEP's global ocean data assimilation system, U850 from the NCEP CDAS. Anomalies for SST, heat content and U850/OLR are departures from the 1971-2000, 1982-2004, 1979-1995 base period pentad means respectively.


## Evolution of Equatorial Pacific Surface Zonal Current Anomaly (cm/s)

U (15m), cm/s, 2°S–2°N




- Anomalous zonal current was in transition from negative (westward) to positive (eastward) in OSCAR since Sep 2010, implying reduction of the zonal advection contribution to the cooling associated with the La Niña conditions.
- However, anomalous zonal current remained strong in GODAS in recent months, which contributed to the imbalance of mixed layer heat budget.

## ENSO cycle as indicated by 1st EOF of surface current and SST anomalies



- Westward surface current anomaly weakened since Jul 2010, similar to the case in 1998 in both its amplitude and evolution.



- On average, ocean surface zonal current anomaly leads the SSTA by a few months.

First EOF mode of ocean surface current (SC) and SST anomalies for the past decade extending through the latest 10-day period. The amplitude time series (top panel) are computed by fitting the data sets to 10-year base period eigenvectors (1993-2002). The amplitudes are then normalized by their respective standard deviations. The bottom panel shows the corresponding EOF maps, scaled accordingly. The El Niño signal can be seen as periods of positive excursions ( $> 1$  Std. Dev.) of the amplitude time series. The near real-time SC are the output from a diagnostic model.


(supplied by Dr. Kathleen Dohan and see ["http://www.esr.org/enso\\_index.html"](http://www.esr.org/enso_index.html) for details)



# Equatorial Pacific Temperature Anomaly

TAO

GODAS




## TAO climatology used


- Negative anomaly of ocean temperature in the eastern-central equatorial Pacific persisted in Oct 2010, consistent with the La Niña conditions.
- Negative temperature anomaly in the central and eastern equatorial Pacific had little propagation.
- Temperature anomaly in TAO intensified from Sep 30 to Oct 15, then weakened slightly, but the intensity change was small in GODAS.



# Oceanic Kelvin Wave Indices




## Standardized Projection on EEOF 1



- Upwelling Kelvin wave occurred in late Feb 2010 in the W. Pacific and propagated eastward, which may have contributed to the transition of ENSO cycle from the warm phase to the cold phase.
- There were no Kelvin wave propagations since Jun 2010.

- Extended EOF (EEOF) analysis is applied to 20-120 day filtered equatorial temperature anomaly in the top 300m using 14 lagged pentads (similar to that in Seo and Xue, GRL, 2005).
- EEOF 1 describes eastward propagation of oceanic Kelvin wave cross the equatorial Pacific in about 70 days.
- Oceanic Kelvin wave indices are defined as standardized projections of total anomalies onto the 14 patterns of EEOF 1.

# Tropical Pacific: SST Anom., SST Anom. Tend., OLR, Sfc Rad, Sfc Flx, 925-mb & 200-mb Winds



- Negative SSTA persisted in the central-eastern equatorial Pacific in Oct 2010.

- Convection was enhanced (suppressed) over the Maritime Continent (in the western-central tropical Pacific).

- Negative SSTA tendency presented mainly in the eastern tropical Pacific.


- Net surface heat flux anomalies damped SSTA between 150W-80W.

- Easterly (westerly) wind anomaly in low level (high level) dominated in the western and central tropical Pacific, consistent with the La Nina conditions.

**Fig. P2. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1979-1995 base period means except SST anomalies are computed with respect to the 1971-2000 base period means.**

# **North Pacific & Arctic Ocean**

# North Pacific & Arctic Ocean: SST Anom., SST Anom. Tend., OLR, SLP, Sfc Rad, Sfc Flx






- Strong positive SSTA in western-central N. Pacific and Arctic Ocean, and negative SSTA in the SE and NE N. Pacific in Oct 2010, consistent with the negative PDO index in Oct 2010 (next slide).

- SSTA tendency showed positive (negative) in the southern (northern) part of N. Pacific, which is generally consistent with latent and sensible heat flux (LH+SH).

- Negative (positive) SLP anomaly presented in the northern (southern and central) part of N. Pacific.

**Fig. NP1.** Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sea surface pressure anomalies (middle-right), sum of net surface short- and long-wave radiation anomalies (bottom-left), sum of latent and sensible heat flux anomalies (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, sea surface pressure and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1979-1995 base period means except SST anomalies are computed with respect to the 1971-2000 base period means.

# PDO index



- The PDO index was near -1.2 in Oct 2010.

- The PDO index has been below normal since Jun 2010.


- Negative PDO index was coincident with the La Nina conditions.

- Pacific Decadal Oscillation is defined as the 1<sup>st</sup> EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly SST anomalies onto the 1st EOF pattern.


- The PDO index differs slightly from that of JISAO, which uses a blend of UKMET and OIv1 and OIv2 SST.

# Arctic Sea Ice

National Snow and Ice Data Center  
<http://nsidc.org/arcticseaicenews/index.html>





- The Arctic sea ice extent in 2010 was well below normal.
- The Arctic sea ice extent was slightly larger during mid Jun-mid Oct 2010 than that in 2007, and became comparable since mid Oct between 2010 and 2007.
- Sea ice extent increased seasonally after mid-Sep 2010.




# North America Western Coastal Upwelling

Pentad Coastal Upwelling for West Coast North America  
( $m^3/s/100m$  coastline)



Standard Positions of Upwelling Index Calculations



- There were some high frequency upwelling and downwelling fluctuations in mid-high latitudes in Oct 2010.

- Anomalous downwelling (upwelling) presented in high (mid) latitudes.

Fig. NP2. Total (top) and anomalous (bottom) upwelling indices at the 15 standard locations for the western coast of North America. Upwelling indices are derived from the vertical velocity of the NCEP's global ocean data assimilation system, and are calculated as integrated vertical volume transport at 50 meter depth from each location to its nearest coast point ( $m^3/s/100m$  coastline). Anomalies are departures from the 1982-2004 base period pentad means.


- Area below (above) black line indicates climatological upwelling (downwelling) season.

- Climatologically upwelling season progresses from March to July along the west coast of North America from 36°N to 57°N.




# Monthly Chlorophyll Anomaly

MODIS Aqua Chlorophyll a Anomaly for October, 2010



- Chlorophyll anomaly pattern persisted from Sep to Oct 2010
- Negative (positive) chlorophyll anomaly presented along the coast north (south) of 33°N.
- The chlorophyll anomalies were generally consistent with the upwelling anomalies in Oct 2010.



MODIS Aqua Chlorophyll a Anomaly for September, 2010






# North America Western Coastal Upwelling

Monthly Coastal Upwelling for West Coast North America  
( $m^3/s/100m$  coastline)



Standard Positions of Upwelling Index Calculations



- Upwelling had been above-normal during the winter of 2006/07, 2007/08, 2008/09.
- But, upwelling was below-normal during the winter of 2009/10.
- Upwelling was relatively strong in spring and summer 2010, and relatively weak since mid Sep 2010.



Fig. NP2. Total (top) and anomalous (bottom) upwelling indices at the 15 standard locations for the western coast of North America. Upwelling indices are derived from the vertical velocity of the NCEP's global ocean data assimilation system, and are calculated as integrated vertical volume transport at 50 meter depth from each location to its nearest coast point ( $m^3/s/100m$  coastline). Anomalies are departures from the 1982-2004 base period pentad means.

- Area below (above) black line indicates climatological upwelling (downwelling) season.
- Climatologically upwelling season progresses from March to July along the west coast of North America from 36°N to 57°N.

# Tropical Indian Ocean

# Evolution of Indian Ocean SST Indices

## Indian Ocean Dipole Mode Indices




- Eastern (SETIO) pole SSTA slightly decreased to about 1C above-normal; while western (WTIO) pole SST decreased to near-normal.
- DMI was below-normal since May 2010, and strengthened since Sep 2010.

**Fig. I1a. Indian Ocean Dipole region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the SETIO [90°E-110°E, 10°S-0] and WTIO [50°E-70°E, 10°S-10°N] regions, and Dipole Mode Index, defined as differences between WTIO and SETIO. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.**

# Recent Evolution of Equatorial Indian SST ( $^{\circ}\text{C}$ ), 0-300m Heat Content ( $^{\circ}\text{C}$ ), 850-mb Zonal Wind (m/s) and OLR ( $\text{W}/\text{m}^2$ ) Anomalies

2 $^{\circ}\text{S}$ –2 $^{\circ}\text{N}$  Average, 3 Pentad Running Mean

5 $^{\circ}\text{S}$ –5 $^{\circ}\text{N}$  Average  
(3 Pentad Running Mean)




- Weakly negative (positive) SSTA presented in the western (central-eastern) Indian Ocean since Jul 2010, consistent with the negative IOD conditions.
- Heat content anomaly transitioned to negative near 75E in Oct 2010, which is not consistent with SSTA.
- Easterly (westerly) wind anomalies presented west (east) of 75E in Oct 2010. The low level wind divergence, consistent with the suppressed convection between 60E-90E, may be cause of the eastward expansion of the negative HC300A.

**Fig. 13.** Time-longitude section of anomalous pentad sea surface temperature (left), upper 300m temperature average (heat content, middle-left), 850-mb zonal wind (U850, middle-right) averaged in 2 $^{\circ}\text{S}$ -2 $^{\circ}\text{N}$  and Outgoing Long-wave Radiation (OLR, right) averaged in 5 $^{\circ}\text{S}$ -5 $^{\circ}\text{N}$ . SST are derived from the NCEP OI SST, heat content from the NCEP's global ocean data assimilation system, and U850 from the NCEP CDAS. Anomalies for SST, heat content and U850/OLR are departures from the 1971-2000, 1982-2004, 1979-1995 base period pentad means respectively.

# Recent Evolution of 10°S Indian SST (°C), 0-300m Heat Content (°C), 850-mb Zonal Wind (m/s)

12°S–8°S Average, 3 Pentad Running Mean

5°S–5°N Average  
(3 Pentad Running Mean)



- Positive SSTa persisted in whole basin and intensified east of 90E in Sep-Oct 2010, contributing to strengthening of negative IOD.
- Positive HC300A propagated westward since Jun 2009, reached the western boundary in Sep 2010, and persisted, negative HC300A in the central and positive HCA in the eastern Ocean intensified in Sep-Oct 2010.
- Low-level wind anomaly was mostly westerly since Aug 2010.


**Fig. I4.** Time-longitude section of anomalous pentad sea surface temperature (left), upper 300m temperature average (heat content, middle-left), 850-mb zonal wind (U850, middle-right) averaged in 12°S–8°S and Outgoing Long-wave Radiation (OLR, right) averaged in 5°S–5°N. SST are derived from the NCEP OI SST, heat content from the NCEP's global ocean data assimilation system, and U850 from the NCEP CDAS. Anomalies for SST, heat content and U850/OLR are departures from the 1971–2000, 1982–2004, 1979–1995 base period pentad means respectively.

# Tropical Indian: SST Anom., SST Anom. Tend., OLR, Sfc Rad, Sfc Flx, 925-mb & 200-mb Wind Anom.

- The tropical Indian Ocean warming weakened in the north and intensified in the south in Oct 2010.

- The SST tendency is partially consistent with net surface heat flux anomalies, except SE Indian Ocean, where dynamics processes likely dominated the warming tendency.

- Convection is enhanced over the N and SE Indian Ocean and Maritime Continent, and suppressed in the central Indian Ocean.





**Fig. 12. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1979-1995 base period means except SST anomalies are computed with respect to the 1971-2000 base period means.**

# **Tropical Atlantic Ocean**



# Evolution of Tropical Atlantic SST Indices




- Positive SSTA in the tropical North and South Atlantic (TNA and TSA) persisted in Oct 2010.
- Meridional Gradient Mode (TNA-TSA) has been above-normal since Feb 2010.
- ATL3 SST has been positive since Dec 2009 and strengthened since Sep 2010.

**Fig. A1a. Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the TNA [60°W-30°W, 5°N-20°N], TSA [30°W-10°E, 20°S-0] and ATL3 [20°W-0, 2.5°S-2.5°N] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.**



# Tropical Atlantic:




- Positive SSTA in the tropical N. Atlantic persisted, but slightly decreased around 10N and increased along Equator.
- Convection was suppressed over the tropical North Atlantic, except the Caribbean Sea regions.
- Relative humidity was mostly below-normal in the tropical Atlantic.
- Easterly wind shear anomaly and above-normal TCHP presented in the tropical Atlantic.

# **North Atlantic Ocean**




# North Atlantic: SST Anom., SST Anom. Tend., OLR, SLP, Sfc Rad, Sfc Flx

- Negative NAO continued in Oct 2010 (next slide), consistent with SLP anomalies.
- Corresponding to the negative NAO was a tripole SSTA pattern with slightly weakening (strengthening) of SSTA in high (mid) latitudes.
- SSTA tendency was not consistent with surface heat flux anomalies.



**Fig. NA1.** Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), **Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sea surface pressure anomalies (middle-right), sum of net surface short- and long-wave radiation anomalies (bottom-left), sum of latent and sensible heat flux anomalies (bottom-right).** SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, sea surface pressure and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1979-1995 base period means except SST anomalies are computed with respect to the 1971-2000 base period means.




# NAO and SST Anomaly in North Atlantic



- NAO Index=-0.9 in Oct 2010.
- NAO has been persistently below-normal since Oct 2009, which contributed to the development and maintenance of negative (positive) SSTA in mid-latitude (tropical) North Atlantic.
- Strong warming presented in the high latitudes North Atlantic Since May 2010.
- Positive SSTA in the Atlantic hurricane MDR has been above-normal since Oct 2009 and slightly weakened since Jun 2010, consistent with the delayed impacts of El Nino and seasonal cycle.
- The combination of persistent negative NAO phase and decay phase of El Nino in spring 2010 results in the strong positive SSTA in MDR, which is similar to 2005.

**Fig. NA2.** Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N (<http://www.cpc.ncep.noaa.gov>). Time-Latitude section of SST anomalies averaged between 80°W and 20°W (bottom). SST are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.

# NAO and SST Anomaly in North Atlantic




- Strong high latitude warming since May 2010.
- Mid-latitude North Atlantic SST has been below-normal since May 2009 and become positive since Jul 2010.
- SST in the Atlantic hurricane MDR has been above-normal since Jul 2009, intensified significantly during Feb-May 2010, and slightly weakened since Jun 2010.

**Fig. NA2.** Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N (<http://www.cpc.ncep.noaa.gov>). Time-Latitude section of SST anomalies averaged between 80°W and 20°W (bottom). SST are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.

# **CFS SST Predictions and Ocean Initial Conditions**

# CFS Niño3.4 SST Predictions from Different Initial Months

## Niño3.4 SST anomalies (K)



- Forecasts from Feb-Apr I.C. show warm biases, and delayed the transition from the warm and neutral phases to the cold phase of ENSO.


- The latest forecast from Oct 2010 I.C. suggests that strong La Niña event will reach its peak in winter of 2010-11, and may last through the spring 2011.

**Fig. M1.** CFS Niño3.4 SST prediction from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1971-2000 base period means.




# U. S. Winter Outlook

(Issued by Climate Prediction Center, NCEP/NWS/NOAA, Oct 21, 2010)



- Warmer winter in the regions from the east part of SW to west part of SE.
- Cooler winter in California, Pacific NW, and Northern Palins.

- Drier winter in the regions from SW to SE.
- Wetter winter in NW, Lower Great Lakes, Ohio Valley, and north part of Tennessee Valley.




Details in “[http://www.noaanews.noaa.gov/stories2010/20101021\\_winteroutlook.html](http://www.noaanews.noaa.gov/stories2010/20101021_winteroutlook.html)”



# CFS DMI SST Predictions from Different Initial Months

## Indian Ocean Dipole SST anomalies (K)



**DMI = WTIO - SETIO**  
**SETIO = SST anomaly in [90°E-110°E, 10°S-0]**  
**WTIO = SST anomaly in [50°E-70°E, 10°S-10°N]**

- Larger inter-ensemble member spread from Feb-Apr 2010 I.C. suggests the forecasts are less reliable. Relatively less spread in the forecasts since May 2010 I.C.
- Forecasts from Sep-Oct 2010 I.C. suggest the current negative IOD will return to normal conditions in winter 2010/11, but regains its strength in summer 2011.


**Fig. M2. CFS Dipole Model Index (DMI) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1971-2000 base period means.**

# CFS Tropical North Atlantic (TNA) SST Predictions

## from Different Initial Months

Tropical N. Atlantic SST anomalies (K)

TNA is the SST anomaly averaged in the region of [60°W-30°W, 5°N-20°N].



- Cold biases were from Feb-Aug I.C.  
- Less spread in the forecasts since Apr 2010 I.C.


- Latest forecasts suggested that positive SSTA in the tropical North Atlantic would decay in following months of 2010, and become near-normal in spring 2011. That is consistent with the delayed impact of ENSO on North Atlantic.

**Fig. M3. CFS Tropical North Atlantic (TNA) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1971-2000 base period means.**

# CFS Pacific Decadal Oscillation (PDO) Index Predictions

## from Different Initial Months

standardized PDO index



PDO is the first EOF of monthly ERSSTv3b anomaly in the region of [110°E-100°W, 20°N-60°N].  
CFS PDO index is the standardized projection of CFS SST forecast anomalies onto the PDO EOF pattern.

- Forecasts from Oct 2009-Jun 2010 I.C. show warm biases and delayed transition from above-normal to below-normal of the PDO index.

- Latest forecasts suggested that the below-normal PDO will last through the Northern Hemisphere spring and summer 2011.

Fig. M4. CFS Pacific Decadal Oscillation (PDO) index predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1971-2000 base period means.

# Overview

- **Pacific/Arctic Ocean**

- ENSO cycle: La Niña conditions persisted with  $\text{NINO3.4} = -1.6^\circ\text{C}$  in Oct 2010.
- NOAA/NCEP Climate Forecast System (CFS) predicted a strong La Niña, to last through the Northern Hemisphere spring 2011.
- PDO has been below-normal since Jul 2010, with  $\text{PDOI} = -1.2$  in Oct 2010.
- Arctic sea ice extent increased seasonally after mid-Sep 2010, and was still well below-normal.

- **Indian Ocean**

- The tropical Indian Ocean Basin warming weakened in the north and slightly strengthened in the south in Oct 2010.
- Dipole Mode index has been below-normal since May 2010, and strengthened to be about  $-0.9^\circ\text{C}$  in Oct 2010.

- **Atlantic Ocean**

- NAO index has been persistently below-normal since Oct 2009, and it was  $-0.9$  in Oct 2010.
- Strong positive SSTA ( $>2.5^\circ\text{C}$ ) presented in the high latitudes since Sep 2010.
- SST in the tropical North Atlantic (TNA) has increased steadily from Dec 2009 to May 2010, gradually weakened from Jun to Aug 2010, and persisted in Sep-Oct 2010.
- The active hurricane season in 2010 was well predicted by NOAA. Multiple factors, including the current La Nina, the delayed impacts of the 2009/10 El Nino, persistent negative NAO, and decadal trend, seem all have contributed to the above normal hurricane activity.

# Backup Slides

# Data Sources and References

- **Optimal Interpolation SST (OI SST) version 2 (Reynolds et al. 2002)**
- **SST 1971-2000 base period means (Xue et al. 2003)**
- **NCEP CDAS winds, surface radiation and heat fluxes**
- **NESDIS Outgoing Long-wave Radiation**
- **NDBC TAO data (<http://tao.noaa.gov>)**
- **PMEL TAO equatorial temperature analysis**
- **NCEP's Global Ocean Data Assimilation System temperature, heat content, currents (Behringer and Xue 2004)**
- **Aviso Altimetry Sea Surface Height**
- **Ocean Surface Current Analyses – Realtime (OSCAR)**

Please send your comments and suggestions to [Yan.Xue@noaa.gov](mailto:Yan.Xue@noaa.gov). Thanks!