

Global Ocean Monitoring: Recent Evolution, Current Status, and Predictions

Prepared by
Climate Prediction Center, NCEP/NOAA

April 11, 2022

<http://www.cpc.ncep.noaa.gov/products/GODAS/>

This project, to deliver real-time ocean monitoring products, is implemented
by CPC in cooperation with NOAA's Global Ocean Monitoring and Observing Program (GOMO)



- **Overview**
- **Recent highlights**
 - Pacific/Arctic Ocean
 - Indian Ocean
 - Atlantic Ocean
- **Global SSTA Predictions**

•Pacific Ocean

- NOAA “ENSO Diagnostic Discussion” on 10 Mar 2022 stated “La Niña is favored to continue into the Northern Hemisphere summer (53% chance during June-August 2022), with a 40-50% chance of La Niña or ENSO-neutral thereafter.”
- La Niña condition persisted with Niño3.4 = -1.0°C in Mar 2022.
- Positive SSTAs continued in the North Pacific.
- The PDO has been in a negative phase since Jan 2020 with PDOI = -1.1 in Mar 2022.

•Arctic Ocean

- “Average Arctic sea ice extent for March 2022 was 14.59 million square kilometers (5.63 million square miles), ranking ninth lowest in the satellite record.”

•Indian Ocean

- SSTAs were small in the tropical Indian Ocean in Mar 2022.


•Atlantic Ocean

- SSTs were near average in the tropical Atlantic Ocean in Mar 2022.
- NAO switched to a positive phase in Dec 2021 with NAOI= 0.4 in Mar 2022.


Global Oceans

Global SST Anomaly (°C) and Anomaly Tendency

MAR 2022 SST Anomaly (°C)
(1991–2020 Climatology)



MAR 2022 – FEB 2022 SST Anomaly (°C)




- Negative SSTAs persisted in the central and southeastern tropical Pacific.
- Positive SSTAs persisted in part of the North Pacific.
- Weak SSTAs were evident across the tropical Atlantic and Indian Oceans.

- Positive (negative) SSTA tendencies were observed in the far-eastern (central and east-central) equatorial Pacific.
- Positive (negative) SSTA tendencies were evident in the southern and eastern (western) North Pacific.
- SSTA tendencies were small in the equatorial Atlantic Ocean and central equatorial Indian Ocean.

SSTAs (top) and SSTA tendency (bottom). Data are derived from the OI SST analysis, and anomalies are departures from the 1991-2020 base period means.


Longitude-Depth Temperature Anomaly and Anomaly Tendency in 2°S-2°N

MAR 2022 Eq. Temp Anomaly (°C)
(GODAS, Clima. 91-20)



- Positive (negative) temperature anomalies were observed along the thermocline in the western and far-eastern (central) equatorial Pacific.
- Positive (negative) temperature anomalies were observed along the thermocline in the western (eastern) equatorial Atlantic Ocean.

MAR 2022 - FEB 2022 Eq. Temp Anomaly (°C)




- Temperature anomaly tendency was positive (negative) along the thermocline in the western and far-eastern (central) Pacific.
- Positive temperature anomaly tendency was evident in the eastern Atlantic Ocean.

Equatorial depth-longitude section of ocean temperature anomalies (top) and anomaly tendency (bottom). Data is from the NCEP's GODAS. Anomalies are departures from the 1991-2020 base period means.


TAO, GODAS, & CFSR monthly mean subsurface temperature anomalies along the Equator during the last 3 months

Ocean Temperature Anomaly in 2S–2N (°C)



TAO (Clim: 1993–2020) GODAS (Clim: 1991–2020) CFSR (Clim: 1991–2020)







- Basic features in the tropical Pacific associated with La Nina evolution are consistent between AVISO and GODAS.
- Positive (negative) anomalous tendencies were present in the eastern (central) equatorial Pacific.
- There are some differences in details between AVISO & GODAS with a lot of small-scale variabilities in AVISO.

Tropical Pacific Ocean and ENSO Conditions


Evolution of Pacific Niño SST Indices



- All Niño indices were negative in Mar 2022, with Niño3.4 = -1.0C.
- Compared with Mar 2021, the east-central and southeastern equatorial Pacific was cooler in Mar 2022.
- The indices may have slight differences if based on different SST products.

Niño region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the specified region. Data are derived from the OI SST analysis, and anomalies are departures from the 1991-2020 base period means.



Comparison of ERSSTv5 & Olv2.1 Niño3.4 Index



- Sometimes, ERSSTv5 is warmer or cooler than Olv2.1.
- For both the extreme positive and negative ($>\pm 1.5^{\circ}\text{C}$) Niño3.4, ERSSTv5 is mostly warmer than Olv2.1.
- During last a couple months, ERSSTv5 was slightly cooler than Olv2.1.

Evolution of Pacific Niño SST Indices


Monthly Tropical Pacific SST Anomaly




- Relative Niño3.4 index is defined as the conventional Niño3.4 index minus the SSTA averaged in the whole tropics (0° - 360° , 20° S- 20° N), in order to remove the global warming signal. Also, to have the same variability as the conventional Niño3.4 index, the relative Niño3.4 index is renormalized (van Oldenborgh et al. 2021: ERL, 10.1088/1748-9326/abe9ed).

[Relative Niño3.4 data updated monthly at:
https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt](https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt)


MAR 2022 SST Anom. (°C)




30MAR2022 - 02MAR2022 SSTA Anom. (°C)




MAR 2022 OLR Anom. (W/m²)




MAR 2022 SW + LW + LH + SH (W/m²)



925mb Wind Anom. (m/s)




200 mb Wind Anom. (m/s)



SSTAs (top-left), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right; positive means heat into the ocean), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.


Current Status of the Pacific Meridional Mode (PMM)

PMM Index (SST based): Dots denote MAR values




PMM Distrib.: ALL_MON = 58.3%

PMM Distrib.: MAR = 60%




SST anomaly for MAR 2022



Lagged regressions of seasonally averaged SST and surface wind anomalies on NPMM SST time series calculated from a Maximum Covariance Analysis.


Lagged Regressions



Amaya, D. J., 2019: *The Pacific meridional mode and ENSO: A review.* *Curr. Climate Change Rep.*, 5, 296–307, 10.1007/s40641-019-00142-x.



<https://www.aos.wisc.edu/~dvimont/MModes/PMM.html>

Evolution of Equatorial Pacific Surface Zonal Current Anomaly (cm/s)



- Anomalous westward currents were observed in both OSCAR and GODAS since Feb 2022.

Oceanic Kelvin Wave (OKW) Index




- Upwelling Kelvin waves were initiated in May, Aug, & Nov 2021, leading to the subsurface cooling in the eastern equatorial Pacific and the development of the 2021/22 La Niña.
- A downwelling Kelvin wave initiated in Dec 2021 led to the weakening of 2021/22 La Niña.
- Dipole-like variations with positive (negative) anomalies in the west (east) were observed since Feb 2022.


(OKW index is defined as standardized projections of total anomalies onto the 14 patterns of Extended EOF1 of equatorial temperature anomalies (Seo and Xue, GRL, 2005).)

Evolution of Pentad D20 and Taux anomalies along the equator


Depth 20°C Pentad Anomaly, ending Apr 05 2022
(2°S–2°N), 12-Pentads Running Mean



Zonal Wind Stress Pentad Anomaly, ending Apr 05 2022
(2°S–2°N), 3-pentad running mean




Equatorial Pacific Ocean Temperature Pentad Mean Anomaly



- Positive (negative) ocean temperature anomalies were present in the central (east-central) Pacific with little propagation.
- The anomalies of GODAS were overall larger than TAO.

Equatorial Sub-surface Ocean Temperature Monitoring



- The equatorial Pacific has been in a recharge phase since Nov 2021.

- Projection of ocean temperature anomalies onto EOF1 and EOF2; EOF1: Tilt/dipole mode (ENSO peak phase); EOF2: WWV mode.


- Recharge/discharge oscillation (ENSO transition phase); Recharge process: heat transport from outside of equator to equator; Negative -> positive phase of ENSO

- For details, see: Kumar & Hu (2014) DOI: 10.1007/s00382-013-1721-0.

Warm Water Volume (WWV) and Niño3.4 Anomalies

- As WWV is linked to ENSO variability (Wyrtki 1985; Jin 1997), it is useful to monitor ENSO in a phase space of WWV and Niño3.4 (Kessler 2002).
- Increase (decrease) of WWV indicates recharge (discharge) of the equatorial oceanic heat content.
- In the WWV index definition, it is the average of ocean temperature anomaly along the whole equatorial Pacific, which sometimes have no coherent variations.





[NINO3.4, WWV] Phase Space



Phase diagram of Warm Water Volume (WWV) and Niño3.4 indices. WWV is the average of depth of 20°C in [120°E-80°W, 5°S-5°N] calculated with the NCEP's GODAS. Anomalies are departures from the 1991-2020 base period means.

North Pacific & Arctic Oceans







Pacific Decadal Oscillation (PDO) Index



- The PDO has been in a negative phase since Jan 2020 with PDOI = -1.1 in Mar 2022.
- Statistically, ENSO leads PDO by 3-4 months, through teleconnection via atmospheric bridge, with El Niño (La Niña) associated with positive (negative) PDO Index.

• PDO is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly SST anomalies onto the 1st EOF pattern.


North Pacific & Arctic Ocean: SSTA, SSTA Tend., OLR, SLP, Sfc Rad, Sfc Flx Anomalies




SSTA (top-left; OI SST Analysis), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) (middle-left; NOAA 18 AVHRR IR), sea surface pressure (middle-right; NCEP CDAS), sum of net surface short- and long-wave radiation (bottom-left; positive means heat into the ocean; NCEP CDAS), sum of latent and sensible heat flux (bottom-right; positive means heat into the ocean; NCEP CDAS). Anomalies are departures from the 1991-2020 base period means.

North America Western Coastal Upwelling

Pentad Coastal Upwelling for West Coast North America
($\text{m}^3/\text{s}/100\text{m}$ coastline)



Standard Positions of Upwelling Index Calculations











- Both the anomalous coastal upwelling and downwelling were weak since mid-Mar 2022.

(top) Total and (bottom) anomalous upwelling indices at the 15 standard locations for the western coast of North America. Derived from the vertical velocity of the NCEP's GODAS and are calculated as integrated vertical volume transport at 50-meter depth from each location to its nearest coast point ($\text{m}^3/\text{s}/100\text{m}$ coastline). Anomalies are departures from the 1991-2020 base period pentad means.


- Area below (above) black line indicates climatological upwelling (downwelling) season.
- Climatologically upwelling season progresses from March to July along the west coast of North America from 36°N to 57°N.

North Pacific SST, OLR, and uv925 anomalies






CFSv2 North Pacific SSTA Predictions

CFSv2 Predicted SST Anomaly (40 Member Mean; °C)




- The CFSv2 predicts that the current SST warm state will continue.



- The seasonal maximum extent of 14.88 million square kilometers was observed on Feb 25.
- Average Arctic sea ice extent for Mar 2022 was 14.59 million square kilometers, ranking ninth lowest in the satellite record.
- The downward linear trend in Mar sea ice extent over the 44-year-satellite record is 2.5% per decade relative to the 1981 to 2010 average.
- Based on the linear trend, since 1979, Mar has seen a loss of 1.74 million square kilometers (672,000 square miles). This is equivalent to about the size of Alaska.



NCEP/CPC Arctic Sea Ice Extent Forecast

Arctic sea ice extent (SIE) forecast
Experimental CFSv2 initialized March 21–25, 2022




Indian Ocean

Evolution of Indian Ocean SST Indices



- Overall, SSTs were near average in the tropical Indian Ocean in Mar 2022.

Indian Ocean region indices, calculated as the area-averaged monthly mean SSTA (OC) for the SETIO [90°E–110°E, 10°S–0] and WTIO [50°E–70°E, 10°S–10°N] regions, and Dipole Mode Index, defined as differences between WTIO and SETIO. Data are derived from the OI SST analysis, and anomalies are departures from the 1991–2020 base period means.





- SSTAs were small in the tropical Indian Ocean.
 - SSTA tendencies along the equator were generally consistent with heat flux anomalies.

SSTAs (top-left), SSTA tendency (top-right), OLR anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.

Tropical and North Atlantic Ocean




Evolution of Tropical Atlantic SST Indices



- SSTAs were small in the tropical Atlantic and positive in the southern Atlantic.
- The Atlantic Niño event ended in Dec 2021.

Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the TNA [60°W-30°W, 5°N-20°N], TSA [30°W-10°E, 20°S-0] and ATL3 [20°W-0, 2.5°S-2.5°N] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the OI SST analysis, and anomalies are departures from the 1991-2020 base period means.


NAO and SST Anomaly in North Atlantic




- NAO switched to a positive phase in Dec 2021 with NAOI= 0.4 in Mar 2022.
- The positive SSTAs in the mid-high latitudes of the North Atlantic Ocean were evident since 2021.

Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N. Time-latitude section of SSTAs averaged between 80°W and 20°W (bottom). SST are derived from the OI SST analysis, and anomalies are departures from the 1991-2020 base period means.

SSTAs in the North Atlantic & MDR



SSTA averaged in MDR (10N–20N, 80W–20W; °C)




- SST in MDR was above average during the last two years.
- SSTA was positive (negative) in the western (eastern) MDR in the last month.


ENSO and Global SST Predictions

Individual Model Forecasts: ENSO will return to neutral in spring or summer


EC: IC= 1Apr 2022




JMA: Updated 11Apr 2022




Australian BOM: Updated 9 Apr 2022



UKMO: Updated 11Apr 2022







Notable warm bias for summer/fall in its Spring IC runs since 2017



Eric Blake - NOAA Federal <eric.s.blake@noaa.gov>

NMME forecasts from different initial conditions



Excessive Momentum and False Alarms in Late-Spring ENSO Forecasts





Fig. 3 False alarm years (1984, 1987, 1991, 2001, 2003, 2011, 2013, 2014, and 2017) in which the forecast probability of the wrong sign of the 3-month tendency exceeded 80% for April–June starts. The black curves are observed monthly values of the Niño 3.4 index with 1-month prior tendencies highlighted in the same color as the corresponding forecast. The colored curves are forecast values with heavy lines for the North American multimodel ensemble mean and light lines for North American multimodel ensemble members. Note the differing vertical scales.

Tippett, M. K., L'Heureux, M. L., Becker, E. J., & Kumar, A. (2020). Excessive momentum and false alarms in late-spring ENSO forecasts. *Geophysical Research Letters*, 47, e2020GL087008. <https://doi.org/10.1029/2020GL087008>




NINO3.4 SST anomalies (K)



- The latest CFSv2 forecasts indicate that La Niña will persist in summer – autumn 2022.


CFS Niño3.4 SST prediction from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means.

IRI/CPC Niño3.4 Forecast: March 2022




- **ENSO Alert System Status: La Niña Advisory**
- Synopsis: *La Niña is favored to continue into the Northern Hemisphere summer (53% chance during June-August 2022), with a 40-50% chance of La Niña or ENSO-neutral thereafter.*

Three Year Evolution of All Double Dip La Niña Winters




Three-year history of sea surface temperatures in the Niño-3.4 region of the tropical Pacific for 8 previous double-dip La Niña events. The color of the line indicates the state of ENSO for the third winter (red: El Niño, darker blue: La Niña, lighter blue: neutral). The black line shows the current event. Monthly Niño-3.4 index is from CPC using ERSSTv5.

<https://www.climate.gov/news-features/blogs/enso/march-2022-la-ni%C3%B1a-update-three-bean-salad>




EEMD of Niño3.4 Index (Black Dot: Intraseasonal-Interseasonal; Red Solid: Interannual; Green Dash: Interdecadal; Shading: Raw)




Monthly mean Niño3.4 index during (a) Jan 1988-Apr 1990, (b) Jan 1998-Apr 2000, (c) Jan 1999-Apr 2001, (d) Jan 2007-Apr 2009, (e) Jan 2010-Apr 2012, and (f) Jan 2020-May 2021. The shading, black dot, red solid, and green dash lines represent raw data, EEMD components at intraseasonal-interseasonal, interannual, and interdecadal and longer time scales, respectively. The unit is $^{\circ}\text{C}$.

- The strength of all the strong La Niña events is determined by the in-phase amplification of all time scale variations.
- Their decay in the boreal spring and early summer is mainly controlled by the intraseasonal-interseasonal variation.

2-year La Niña may be predictable 18 to 24 months in advance





IC: -WWV



IC: El Niño

Temporal evolution of potential prediction utility (PPU) of the Nino-3.4 index for “*perfect CESM1*” in forecasts initialized a during the discharge phase (c) and at the peak of the preceding El Niño event (d). The dashed black line indicates January of the second year (Jan0).

- A 1800-year long control simulation of CESM1 & forecasts with the perfect model approach.
- A strong thermocline discharge or a strong El Niño can lead to La Niña conditions that last 2 years.
- Forecasts initialized with strong thermocline discharge or strong peak El Niño amplitude show higher predictability than those with initial conditions of weaker magnitude.
- **2-year La Niña may be predictable 18 to 24 months in advance under specific initial conditions.**




Niño-3.4 SST anomalies ($^{\circ}\text{C}$) during January 2014–March 2015. Black, green, red, purple and blue curves are for SST anomalies in observations, CTL, SE, SE3 and SEpTNWP, respectively. For forecasts, solid curves represent the ensemble mean. Shaded areas in light green and light red represent ensemble spread for CTL and SE. The dots in red, purple and blue curves means that the ensemble mean SST anomalies in SE, SE3 and SEpTNWP are significantly different from those in CTL at the 95% confidence level.

For 2014/15 Nino3.4, “Our experiments show that 40% of the amplitude error at the peak phase could be attributed to the lack of prediction of negative SST anomalies in the southeastern Pacific.” (Zhu et al. 2016)

Zhu, J., A. Kumar, B. Huang, M. A. Balmaseda, Z.-Z. Hu, L. Marx, and J. L. Kinter III, 2016: The role of off-equatorial surface temperature anomalies in the 2014 El Niño prediction. *Sci. Rep.*, 6, 19677. DOI: 10.1038/srep19677.


standardized PDO index



- CFSv2 predicts a negative phase of PDO in 2022.

CFS Pacific Decadal Oscillation (PDO) index predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means. PDO is the first EOF of monthly ERSSTv3b anomaly in the region of [110°E-100°W, 20°N-60°N]. CFS PDO index is the standardized projection of CFS SST forecast anomalies onto the PDO EOF pattern.


Indian Ocean Dipole SST anomalies (K)



- CFSv2 predicts a negative phase of IOD in summer-autumn 2022.

CFS Dipole Model Index (DMI) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1991-2020 base period means.

Tropical N. Atlantic SST anomalies (K)




- Latest CFSv2 predictions call for slightly above normal SST in the tropical North Atlantic in 2022.

CFS Tropical North Atlantic (TNA) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means. TNA is the SST anomaly averaged in the region of [60w-30w, 5n-20n].

CFSv2 Atlantic SSTA Predictions

CFSv2 Predicted SST Anomaly (40 Member Mean; °C)



- Latest CFSv2 predictions call slightly above or near average SST in the tropical North Atlantic in the next 8 months.

Acknowledgement

- ❖ Drs. Jieshun Zhu, Caihong Wen, and Arun Kumar: reviewed PPT, and provide insightful suggestions and comments
- ❖ Dr. Pingping Xie provided the BASS/CMORPH/CFSR EVAP package
- ❖ Dr. Wanqiu Wang provides the sea ice forecasts and maintains the CFSv2 forecast archive

Please send your comments and suggestions to:

Arun.Kumar@noaa.gov

Jieshun.Zhu@noaa.gov

Caihong.Wen@noaa.gov

Zeng-Zhen.Hu@noaa.gov


- **Weekly Optimal Interpolation SST (OI SST) version 2 (Reynolds et al. 2002)**
- **Extended Reconstructed SST (ERSST) v5 (Huang et al. 2017)**
- **Blended Analysis of Surface Salinity (BASS) (Xie et al. 2014)**
- **CMORPH precipitation (Xie et al. 2017)**
- **CFSR evaporation adjusted to OAFlux (Xie and Ren 2018)**
- **NCEP CDAS winds, surface radiation and heat fluxes (Kalnay et al. 1996)**
- **NESDIS Outgoing Long-wave Radiation (Liebmann and Smith 1996)**
- **NCEP's GODAS temperature, heat content, currents (Behringer and Xue 2004)**
- **Aviso altimetry sea surface height from CMEMS**
- **Ocean Surface Current Analyses – Realtime (OSCAR)**
- **In situ data objective analyses (IPRC, Scripps, EN4.2.1, PMEL TAO)**
- **Operational Ocean Reanalysis Intercomparison Project**
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora_body.html
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html

Backup Slides

Monthly SSS Anomaly Evolution over Equatorial Pacific

NOTE: Since June 2015, the BASS SSS is from in situ, SMOS and SMAP; before June 2015, The BASS SSS is from in situ, SMOS and Aquarius.

- Hovemoller diagram for equatorial SSS anomaly (5° S- 5° N);
- In the equatorial Pacific Ocean, west of 140° E, negative SSS signal continues; positive SSS signal continues between 140° E and 170° W; neutral or likely negative signal continues east of 150° W.



Global Sea Surface Salinity (SSS): Anomaly for February 2022

New Update: The NCEI SST data used in the quality control procedure has been updated to version 2.1 since May 2020;

Positive SSS anomaly continues in the western equatorial Pacific Ocean and SPCZ caused reduced precipitation in these areas. Positive SSS anomaly continues between 10° N and 40° N across the North Atlantic Ocean. Negative SSS anomaly continues in the equatorial Atlantic Ocean where enhanced precipitation is observed. Negative SSS anomaly continues in the Bay of Bengal, while positive anomaly appears in the Arabian Sea.


SSS : Blended Analysis of Surface Salinity (BASS) V0.Z

(a CPC-NESDIS/NODC-NESDIS/STAR joint effort)

<ftp.cpc.ncep.noaa.gov/precip/BASS>


Precipitation: CMORPH adjusted satellite precipitation estimates

Evaporation: Adjusted CFS Reanalysis



Global Sea Surface Salinity (SSS): Tendency for February 2022


ITCZ precipitation is enhanced across both the Pacific and the Atlantic ocean, with a slight meridional shift towards north over the majority of the Pacific and towards south over the Atlantic. Precipitation over the SPCZ is weakened. At the meantime, evaporation over the northwest Pacific storm track region is reduced substantially. These changes in the oceanic fresh water, combined with the effects of oceanic transportation and mixed layer processes, created a slight freshened (saltier) tendency of SSS along the Pacific / Atlantic ITCZ (SPCZ) regions, respectively.



Pentad SSS Anomaly Evolution over Equatorial Pacific


Figure caption:

Hovemoller diagram for equatorial (5° S- 5° N) 5-day mean SSS, SST and precipitation anomalies. The climatology for SSS is Levitus 1994 climatology. The SST data used here is the OISST V2 AVHRR only daily dataset with its climatology being calculated from 1985 to 2010. The precipitation data used here is the adjusted CMORPH dataset with its climatology being calculated from 1999 to 2013.




Equatorial Pacific SST ($^{\circ}\text{C}$), HC300 ($^{\circ}\text{C}$), u850 (m/s) Anomalies


2 $^{\circ}\text{S}$ –2 $^{\circ}\text{N}$ Average, 3 Pentad Running Mean




MAR 2022 SST Anom. ($^{\circ}\text{C}$)




30MAR2022 - 02MAR2022 SST Anomaly ($^{\circ}\text{C}$)




MAR 2022 TCHP Anom. (KJ/cm^2)




MAR 2022 OLR Anom. (W/m^2)




MAR 2022 200mb Wind Anom. (m/s)




MAR 2022 200mb - 850mb Wind Shear Anom. (m/s)




MAR 2022 SW + LW Anom. (W/m^2)




LH + SH Anom. (W/m^2)




925mb Wind Anom. (m/s)




MAR 2022 700 mb RH Anom. (%)




MAR 2022 SST Anom. ($^{\circ}\text{C}$)




30MAR2022 - 02MAR2022 SST Anom. ($^{\circ}\text{C}$)




MAR 2022 OLR Anom. (W/m^2)




MAR 2022 SLP Anom. (hPa)



MAR 2022 SW + LW (W/m^2)




MAR 2022 LH + SH (W/m^2)



Ocean Mixed-Layer Heat Budget

- Observed SSTA tendency ($dSSTA/dt$; bar) and total heat budget (RHS; black line) were positive recently.

- Dynamical terms (Q_u , Q_v , Q_w+Q_{zz}) were either small negative or positive and heat-flux term (Q_q) were negative recently.



Huang, B., Y. Xue, X. Zhang, A. Kumar, and M. J. McPhaden, 2010 : The NCEP GODAS ocean analysis of the tropical Pacific mixed layer heat budget on seasonal to interannual time scales, *J. Climate.*, 23, 4901-4925.

Q_u : Zonal advection; Q_v : Meridional advection;

Q_w : Vertical entrainment; Q_{zz} : Vertical diffusion

Q_q : $(Q_{net} - Q_{open} + Q_{corr})/pcph$; $Q_{net} = SW + LW + LH + SH$;

Q_{open} : SW penetration; Q_{corr} : Flux correction due to relaxation to OI SST