



# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

Update prepared by  
Climate Prediction Center / NCEP  
January 3, 2011



# Outline

- Overview
- Recent Evolution and Current Conditions
- MJO Index Information
- MJO Index Forecasts
- MJO Composites



# Overview

- The MJO remained weak during the last seven days.
- The majority of dynamical model MJO forecasts indicate continued no MJO activity during the upcoming week, with background La Niña conditions and other subseasonal coherent tropical variability expected to dominate the pattern of tropical convection.
- There is some spread in model forecasts for the evolution of the MJO during the Week-2 period, but most maintain weak activity.

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies


Westerly anomalies strengthened across the eastern Indian Ocean but contracted in coverage.



Easterly anomalies continued across the western and central equatorial Pacific and shifted to the west.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )



Westerly anomalies (orange/red shading)  
represent anomalous west-to-east flow

Easterly anomalies (blue shading)  
represent anomalous east-to-west flow

Easterly anomalies have persisted in the west-central Pacific since June (black box) consistent with the development of La Niña conditions.


Enhanced westerly anomalies (red dotted oval) occurred across the eastern Pacific during early-to-mid July, partly associated with MJO activity.

The MJO strengthened in October as evidenced by weak westerly anomalies and a weakening of the easterlies across the central Pacific during mid-October. (blue box).

In mid-December, easterly anomalies weakened just west of the Date Line due to a combination of weak MJO activity and extratropical interactions.




# OLR Anomalies – Past 30 days




Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During early December, enhanced convection (blue circle) continued over parts of the eastern Indian Ocean, Australia and along the SPCZ.



Enhanced convection continued over the Maritime continent and Australia during mid December while suppressed convection (red circle) strengthened across the western Indian Ocean.




During mid-to-late December, enhanced convection weakened but remained over northern South America, continued over Australia, and expanded across the Maritime continent. Drier-than-average conditions continued across parts of the Indian Ocean and central Pacific Ocean.



# Outgoing Longwave Radiation (OLR)

## Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

From late-July into September, generally enhanced (suppressed) convection prevailed across the western Maritime continent (Date Line) (red box). Considerable intraseasonal variability was evident during the period but the MJO did not play a large role.

As the MJO strengthened in late September into October, enhanced convection developed near 60E and shifted eastward followed by suppressed convection near 20E during early-mid October.


Weak MJO activity was again experienced during late November into December. An area of enhanced convection propagated eastward from the Indian Ocean to the Maritime continent followed by suppressed convection thereafter.



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation  
Negative anomalies (green shading) indicate favorable conditions for precipitation

Time ↓



Eastward propagation was evident during mid-July associated with the MJO.

Eastward propagation in August and September was mainly associated with higher frequency coherent tropical variability rather than the MJO (red box).

The MJO strengthened during late September as anomalies increased and eastward propagation was seen through mid-October.


During late November and early December, some eastward propagation associated with the MJO is evident in velocity potential anomalies.



# IR Temperatures (K) / 200-hPa

## Velocity Potential Anomalies


Positive anomalies (brown contours) indicate unfavorable conditions for precipitation  
Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows anomalous upper-level divergence over South America and the Maritime continent. Anomalous convergence is present across the central Pacific and western Indian Ocean.



# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )



Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies


Red shades: Westerly anomalies

Westerly anomalies continued across most of the equatorial Pacific with easterly anomalies across the tropical Atlantic Ocean weakening. Easterly anomalies across the Indian Ocean strengthened.

The off-equatorial cyclone in the northern hemisphere has weakened.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )



Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since early July. Eastward propagation of westerly anomalies in August and September were not associated with the MJO.


In early October, westerly anomalies strengthened considerably associated with MJO activity and an eastward extension of these anomalies is evident.

During November, westerly anomalies increased episodically from 140W to 80W.

Easterly anomalies were enhanced across the Indian Ocean with evidence of tropical intraseasonal variations.



# Weekly Heat Content Evolution in the Equatorial Pacific



From January through March 2010, heat content anomalies remained above-average for much of the period.

From December 2009 – February 2010 two ocean Kelvin waves contributed to the change in heat content across the eastern Pacific (last two dashed black lines).

During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Niña.

Currently, negative heat content anomalies extend across the central and eastern Pacific with positive anomalies in the western Pacific.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index did not indicate significant MJO activity during the past week.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.



# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

**RMM1 and RMM2 values for the most recent  
40 days and forecasts from the ensemble Global  
Forecast System (GEFS) for the next 15 days**

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts


The GFS forecasts were not updated  
today due to a computer-related issue.  
The figure will be updated later today  
(1/3) or tomorrow (1/4).




# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

## Spatial map of OLR anomalies for the next 15 days



## Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The GEFS ensemble mean forecast indicates moderate anomalies in some areas but these are more related to a combination of La Niña and other modes of subseasonal tropical variability.




# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


MJO activity is not forecast during the period due to a weak initial signal.






# MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)




850-hPa Wind Anomalies (Nov-Mar)





# U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>