



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
January 24, 2011**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained active during the past week with the enhanced convective phase located in the western Pacific.**
- **Model MJO index forecasts indicate a weakening signal during Week-1. However, some models indicate a continued eastward propagation. Consequently, uncertainty is high during the Week-2 period.**
- **The MJO is expected to contribute to enhanced rainfall across the SPCZ (Week-1), and parts of Africa (Week-1). Anomalous convection associated with the ongoing La Nina can be expected during Week-2.**
- **For the U.S., the forecast MJO phase is consistent with below normal temperatures across parts of the eastern U.S. during Week-1 and Week-2. Forecast MJO phases are also consistent with extension of the Pacific jet stream during the Week-2 period, but this is currently at odds with current numerical guidance. Uncertainty is high and forecast confidence low for circulation changes across North America.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies continued across the eastern Indian Ocean and western Maritime continent.

Westerly anomalies extended south of the equator into the western Pacific east of Australia during the five to ten days.


Easterly anomalies continued across the western and central equatorial Pacific but have weakened since early January.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

Easterly anomalies have persisted in the west-central Pacific since June (black box) consistent with the development of La Nina conditions.

The MJO strengthened in October as evidenced by weak westerly anomalies and a weakening of the easterlies across the central Pacific during mid-October. (blue box).


In mid-December, easterly anomalies weakened just west of the Date Line due to a combination of weak MJO activity and extratropical interactions.

Easterly anomalies have recently weakened near the Date Line due to increased MJO activity.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During late December, enhanced convection (blue circle) continued over parts of the eastern Indian Ocean, Australia and along the SPCZ while suppressed convection (red circle) was evident across the western Indian Ocean and parts of eastern Africa.

Enhanced convection continued over the Maritime continent and Australia during late December and early January.

During early-to-mid January, enhanced convection continued over the Maritime continent and Australia and intensified along the SPCZ once again. Enhanced convection developed in southeast Africa by mid January.






# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
8-Aug-2010 to 23-Jan-2011 + 14 days

Time  
↓



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

As the MJO strengthened in late September into October, enhanced convection developed near 60°E and shifted eastward followed by suppressed convection near 20°E during early-mid October.

Weak MJO activity was again experienced during late November into December. An area of enhanced convection propagated eastward from the Indian Ocean to the Maritime continent followed by suppressed convection thereafter.

MJO strengthened by mid-January as enhanced convection shifted east from the Maritime Continent, while suppressed convection propagated east across the Indian Ocean.


Longitude



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



Eastward propagation in August and September was mainly associated with higher frequency coherent tropical variability rather than the MJO (red box).

The MJO strengthened during late September as anomalies increased and eastward propagation was seen through mid-October.

During late November and early December, some eastward propagation associated with the MJO is evident in velocity potential anomalies.

Most recently, anomalous upper level divergence has developed across parts of the Pacific in part associated with a strengthening MJO signal.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows anomalous upper-level divergence over the Maritime continent and parts of the Pacific with anomalous upper-level convergence over the Indian Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Cyclonic circulations (C) are evident north and south of the equator.




Westerly anomalies strengthened over the western Indian Ocean.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since early July. Eastward propagation of westerly anomalies in August and September were not associated with the MJO.


In early October, westerly anomalies strengthened considerably associated with MJO activity and an eastward extension of these anomalies is evident.

Most recently, easterly anomalies have shifted east towards the Date Line.



# Weekly Heat Content Evolution in the Equatorial Pacific

Time  
↓



From January through March 2010, heat content anomalies remained above-average for much of the period.

From December 2009 – February 2010 two ocean Kelvin waves contributed to the change in heat content across the eastern Pacific (last two dashed black lines).

During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Currently, negative heat content anomalies extend across the central and eastern Pacific with positive anomalies in the western Pacific.

Longitude





# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution

- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months





The MJO index indicates a strengthening signal during the past week.





# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present. Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS forecasts continued eastward propagation followed by a weakening signal during week-1. Uncertainty increases dramatically during week-2.







# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

**Spatial map of OLR anomalies for the next 15 days**

**Time-longitude section of (7.5 S-7.5 N) OLR anomalies for the last 180 days and for the next 15 days**



The GEFS ensemble mean forecast indicates suppressed convection continuing over the Indian Ocean, while enhanced convection develops across the eastern Maritime Continent and western Pacific Ocean.




# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

MJO activity is forecast to continue at least through this week with suppressed convection shifting east across the Maritime Continent and western Pacific.









# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)



## 850-hPa Wind Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>





# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>