



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
January 10, 2012**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remained active during the past week with its signal primarily centered over the Pacific Ocean. The activity appears weaker than that observed during autumn 2011.**
- **La Nina and additional subseasonal tropical variability are substantially affecting the MJO signal and the anomalous convection in the global Tropics.**
- **At the current time, most dynamical model MJO index forecasts do not indicate a strong MJO signal over the upcoming period, but some MJO signal is likely to continue.**
- **The MJO is likely to continue to contribute to above (below) average rainfall across the South Pacific (portions of the Indian Ocean) mainly during Week-1.**
- **The recent evolution of the MJO is consistent with above-average precipitation for the west coast as indicated by some forecast guidance during the Week-2 period.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly wind anomalies continued over the equatorial Indian Ocean, but weakened during the past five days.

Easterly anomalies continued over the west Pacific.


Westerly wind anomalies strengthened over the east Pacific Ocean during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



In early October, MJO activity weakened the persistent easterly anomalies across the central Pacific (first dashed line).

An equatorial Rossby wave imparted westerly anomalies across parts of the western Pacific and Maritime continent during late October and early November (thin solid line).


MJO activity continued into December (altering dashed and dotted lines), but most recently westerly (easterly) wind anomalies across the Indian Ocean (western Pacific) have become more stationary.

Longitude



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During early December, enhanced convection (blue circle) developed near the Philippines, the Maritime continent and the South Pacific Convergence Zone (SPCZ). Convection across the Indian Ocean became mixed and suppressed convection (red circle) continued near the Date Line.


A large area of enhanced convection was evident from the northern Indian Ocean to the SPCZ during late December, while suppressed convection continued near the Date Line.

A large area of enhanced convection stretching from the Maritime Continent to the SPCZ continued into the beginning of January. Suppressed convection developed over the equatorial Indian Ocean.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
24-Jul-2011 to 8-Jan-2012 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

Beginning in mid-September, enhanced convection shifted from southern Asia to the western Pacific while suppressed convection developed during late September across India and also shifted eastward to the western Pacific.

MJO activity continued into the first half of December when OLR anomalies decreased and eastward propagation was not clear.

During late December, eastward propagation of OLR anomalies is indicated once again with suppressed convection indicated across the Indian Ocean during the first week in January.

Longitude

Time
↓




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



During July and August very fast eastward propagation was evident at times and mainly associated with higher frequency sub-seasonal coherent tropical variability not associated with MJO activity.

Beginning in the second half of September into December, alternating negative (dashed lines) and positive (dotted lines) anomalies were evident and associated with MJO activity during the period.


Eastward propagation of anomalies became less coherent during December as compared to autumn 2011.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The latest large scale velocity potential pattern shows anomalous upper-level divergence across the Maritime continent and parts of the South Pacific, while anomalous upper-level convergence is evident across the Indian Ocean.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Large scale cyclonic circulations are evident both north and south of the equator near the Date Line and upper-level westerly wind anomalies persist across the east Pacific (red boxes).


During the past five days, westerly anomalies have spread east into Africa.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) during much of the period prior to mid-September.

Alternating westerly (dashed lines) and easterly (dotted lines) anomalies are evident from mid-September into December associated with the MJO.

Westerly anomalies over the Pacific have substantially strengthened during early-to-mid December and some propagation is evident during the past few weeks.



Weekly Heat Content Evolution in the Equatorial Pacific



Since the beginning of January 2011, positive heat content anomalies shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011. Much of the Pacific basin now indicates above- or near-normal integrated heat content.

Since late July, negative heat content anomalies are evident across the equatorial central and eastern Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index decreased in amplitude with little eastward propagation during the past week. The enhanced phase remains across the Pacific.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present. Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecasts a nearly stationary signal with the enhanced phase persisting across the Pacific during the next two weeks.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates enhanced convection across the western Pacific and suppressed convection continuing in the Indian Ocean primarily during the next two weeks.







Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The CA forecast shows enhanced convection over parts of the western Pacific. Suppressed convection shifts east from the Indian Ocean to the Maritime Continent.




MJO Composites – Global Tropics

850-hPa Wind Anomalies (Nov-Mar)



Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>