



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
March 26, 2012**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained active during the past week with the enhanced phase crossing the Pacific. Enhanced convection was evident across portions of the western Pacific and the Americas.**
- **Dynamical model MJO index forecasts show continued MJO activity, although with slower eastward propagation. Statistical models indicate more consistent eastward propagation.**
- **Based on the latest observations and forecast models, the MJO is forecast to remain active and impact the Western Hemisphere over the next 1-2 weeks.**
- **The MJO is forecast to contribute to enhanced convection across parts of the Pacific and the Americas during the period. Suppressed convection remains favored for much of the Indian Ocean and Maritime Continent during the period.**
- **Upcoming phases of the MJO favor and are consistent with an increase in chances of colder air entering the central and eastern continental U.S. beginning during the Week-2 period. MJO composites favor below-normal temperatures for parts of the central and southeast U.S. and above-average precipitation for parts of the southeast U.S. beginning in Week-2.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**





# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies shifted eastward and increased in coverage during the past five days in proximity to the Maritime continent region.


Easterly anomalies persisted over the Central Pacific during the past five days, but have decreased in magnitude and coverage.

Westerly wind anomalies increased over the eastern Pacific during the past five days.



# 850-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



In early October, MJO activity weakened the persistent easterly anomalies across the central Pacific (first dashed line).

MJO activity continued into December (altering dashed and dotted lines), but then westerly (easterly) wind anomalies across the Indian Ocean (western Pacific) became more stationary.

In early February, westerly anomalies extended to 140E and were associated with MJO activity.


Most recently, westerly anomalies associated with the MJO have propagated to the western Pacific, near the Date Line (black box).

Longitude



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During mid February to March, suppressed convection was observed over the central and Western Pacific. Enhanced convection was observed across Africa and the Indian Ocean.


Enhanced convection was evident from Africa to the Maritime Continent during early March. Suppressed convection was observed for the west/central Pacific and parts of the South America.

During early to mid-March, the MJO contributed to enhanced (suppressed) convection across the Maritime Continent and Western Pacific (central Pacific, South America and Africa).



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
9-Oct-2011 to 25-Mar-2012 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

MJO activity was evident during October, November and early December as alternating areas of enhanced (dashed lines) and suppressed (dotted lines) convection shifted eastward.

The MJO once again strengthened during late January as enhanced convection shifted eastward across the Maritime continent. The MJO has continued into March, with enhanced convection beginning across the Western hemisphere and suppressed convection building across the Indian Ocean.

**Longitude**


**Time**  
↓



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



Beginning in the second half of September and lasting until December, alternating negative (dashed lines) and positive (dotted lines) anomalies were evident and associated with MJO activity during the period.

Eastward propagation of anomalies became less coherent during late December and early January and anomalies weakened.

The MJO strengthened in late January and eastward propagation has been evident through mid-March. Most recently, enhanced divergence (convergence) is centered over the Pacific (Indian Ocean).






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows a wave-1 structure with the strongest upper-level divergence across the Pacific and Americas and with upper-level convergence mainly over the Indian Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




During the most recent five days, westerly anomalies over the central Pacific have shifted slightly eastward, to Africa and the western Indian Ocean.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Alternating westerly (dashed lines) and easterly (dotted lines) anomalies are evident from mid-September into December associated with the MJO.

In December, westerly anomalies strengthened over the Pacific.

Eastward propagation was again more clearly evident during late January and February, continuing until the present time. Upper-level easterly anomalies are evident near 140E, with the westerly anomalies shifting eastward, over the Americas and Africa.



# Weekly Heat Content Evolution in the Equatorial Pacific



Time



An oceanic Kelvin wave (dashed line) shifted eastward during March and April 2011.

Since late July, negative heat content anomalies are evident across the equatorial central and eastern Pacific.

In February and March 2012, negative heat content anomalies weakened in the central and eastern equatorial Pacific.

Longitude



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index shows continued MJO activity during the past week centered mainly across the central Pacific.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

**light gray shading:** 90% of forecasts  
**dark gray shading:** 50% of forecasts

The ensemble GFS forecasts MJO activity to continue during the next two weeks, although eastward propagation early in the period is slow.








# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates enhanced (suppressed) convection across the western Pacific (Indian Ocean) during most of the period.







# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The CA forecast shows suppressed convection shifting from the Indian Ocean across the Maritime Continent during the period with enhanced convection impacting the Americas and Africa.




# MJO Composites – Global Tropics

## 850-hPa Wind Anomalies (Nov-Mar)



## Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>