



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
April 23, 2012**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained active during the past week with the enhanced phase centered across Africa, however, some measures indicate that the MJO may be weakening and becoming less coherent.**
- **Dynamical model MJO index forecasts indicate very weak MJO activity during the period, a stark contrast to the recent months of strong, coherent activity.**
- **Disruption of the signal is likely related to interference from other types of coherent subseasonal variability as well as a weakening of the MJO itself so it is too early, however, to say that the recent MJO activity is ending.**
- **At the current time, the MJO is not expected to contribute substantially to areas of anomalous tropical convection during the period.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies are now evident across most areas of central Africa and the Indian Ocean except for parts of the equatorial Indian Ocean.




Easterly anomalies continued across the western Pacific and ended across most of the Maritime continent during the past five days.

Westerly anomalies weakened over the eastern Pacific during the past five days.



# 850-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

MJO activity continued into December as indicated by altering dashed and dotted lines. Later during other portions of December and January, westerly (easterly) wind anomalies across the Indian Ocean (western Pacific) became more stationary.


During the first half of February, the MJO contributed to increased westerly anomalies near 140E and across the eastern Pacific while decreasing easterly anomalies in the central Pacific.

MJO activity continued into April, with westerly anomalies associated with the MJO located near the Date Line and western hemisphere early in the month. Most recently, easterly anomalies have returned to the Date Line during mid-April.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**




Late in March, enhanced convection shifted eastward into the western Pacific while suppressed convection continued over east Africa and developed across the Indian Ocean.

During early April, enhanced convection intensified over portions of the eastern Pacific, Americas and western Africa with suppressed convection shifting eastward to include parts of the Maritime continent and Australia.

During mid-April, enhanced convection continued across portions of South America, Africa and the Indian Ocean, north and south of the equator. Suppressed convection was evident near the Date Line in the western Pacific.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

(Courtesy of CAWCR Australia Bureau of Meteorology)

MJO activity was evident in November and early December and briefly again in late December and early January as alternating areas of enhanced (dashed lines) and suppressed (dotted lines) convection shifted eastward.

Strong MJO activity once again developed during late January and has continued into April. The enhanced convection is now entering areas in the eastern hemisphere such as Africa and the western Indian Ocean.




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



Beginning in the second half of September and lasting until December, alternating negative (dashed lines) and positive (dotted lines) anomalies were evident and associated with MJO activity during the period.

Eastward propagation of anomalies became less coherent during late December and early January and anomalies weakened.

The MJO strengthened in late January and eastward propagation has been evident through mid-April. The period of this MJO activity has been considerably longer than the activity observed during late 2011.


Longitude



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows enhanced divergence mainly across Africa, the Indian Ocean and the Maritime continent. Enhanced convergence is generally indicated over the Pacific.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies shifted eastward from Africa and the Indian Ocean to parts of the Maritime continent and western Pacific (red boxes).



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Alternating westerly (dashed lines) and easterly (dotted lines) anomalies associated with the MJO are evident from October into December.

In December, westerly anomalies strengthened over the central Pacific.

Eastward propagation was again more clearly evident during late January and February, continuing into April when westerly anomalies shifted eastward, over the Americas, Africa and the western Indian Ocean.



# Weekly Heat Content Evolution in the Equatorial Pacific



Time



From July 2011 through February 2012, heat content was below average in the central and eastern equatorial Pacific.

Recent heat content anomalies have returned to near zero across much of the central and eastern Pacific.

Longitude



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index continued to show eastward propagation to the Indian Ocean during the past week. In recent days the index has decreased in amplitude and eastward propagation has slowed.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS MJO index forecasts show weak MJO activity during the period.






# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates small anomalies during the next ten days with suppressed convection strengthening across the eastern Indian Ocean later in Week-2.






# Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The CA forecast shows enhanced convection slowly shifting to the Maritime Continent during the period with weak suppressed convection impacting parts of the Americas and Africa.






# MJO Composites – Global Tropics

## 850-hPa Wind Anomalies (Nov-Mar)



## Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>