



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
August 6, 2012**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained active during the past week with the enhanced phase propagating east across the Pacific Ocean.**
- **Dynamical model MJO index forecasts indicate considerable spread, although most show an eastward propagating signal over the next two weeks.**
- **Based on the latest observations, the MJO is forecast to remain active with the enhanced phase impacting the eastern Pacific, the Americas, the Atlantic and Africa during the period.**
- **The MJO is expected to contribute to enhanced convection and elevated chances for tropical cyclone development across the east Pacific (Week-1 / Week-2) and Atlantic (Week-2 / Week-3). Suppressed convection is expected to shift northeast from the Maritime Continent to the west Pacific.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies strengthened and expanded eastward into the west Pacific.

Westerly wind anomalies continue across the eastern tropical Atlantic and western Africa.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

During the first half of February, the MJO (alternating black dashed and dotted lines) contributed to increased westerly anomalies near 140E and across the eastern Pacific while decreasing easterly anomalies in the central Pacific. MJO activity continued into April, with westerly anomalies associated with the MJO located near the Date Line and western hemisphere early in the month.

Anomalies were generally persistent across much of the global tropics during April and May.


Strong westerly anomalies developed across the eastern Pacific in mid-June and shifted westward (black solid line) and likely were associated with a robust equatorial Rossby wave as it progressed west across the Pacific.

Most recently, westerly anomalies have developed between 120-140E.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



Enhanced convection was observed over the Indian Ocean, Maritime Continent, and much of the North American monsoon region.

During mid-July, anomalous convection shifted east across the Maritime Continent into the western Pacific with suppressed convection developing across the Indian Ocean. Convection remained enhanced over Africa.


By early August, enhanced (suppressed) convection persisted across the west Pacific (Indian Ocean). Suppressed convection developed across the east Pacific, Mexico, and Central America.





# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
19-Feb-2012 to 5-Aug-2012 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

**Strong MJO activity (alternating dashed and dotted lines) was evident during February and continued into mid-April.**

**Anomalies became less coherent during most of April and May.**

**In late May through July, eastward propagation of both enhanced and suppressed convection is evident across the eastern hemisphere. Atmospheric Kelvin wave activity also played a large role in the pattern of anomalous convection across the Pacific and western Hemisphere during this period.**

**Longitude**




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



Longitude

The MJO strengthened in late January as indicated by alternating negative (dashed lines) and positive (dotted lines) anomalies with eastward propagation. The activity continued into mid-April.

Beginning in late April, anomalies became weaker and less coherent than earlier in the year.

Eastward propagation was once again evident from late May through July associated with the MJO as well as atmospheric Kelvin wave activity, which at times resulted in fast eastward propagation of observed anomalies.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern reflects anomalous upper-level divergence across the west and central Pacific with anomalous upper-level convergence stretching from the Atlantic to the Indian Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies persisted north of the equator in the central Pacific with weak easterly anomalies closer to the equator (blue box). Westerly anomalies also shifted eastward across Africa to the Indian Ocean (red box).



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




The MJO strengthened once again in late January as indicated by alternating westerly (dotted lines) and easterly (dashed lines) anomalies. This activity continued to mid-April.

Anomalies were less coherent during much of April and May.

Some evidence of faster moving, subseasonal variability can be seen during much of June and July. Most recently, easterly anomalies have become strong over the Maritime continent.



# Weekly Heat Content Evolution in the Equatorial Pacific



From July 2011 through February 2012, heat content was below average in the central and eastern equatorial Pacific.

From March into July 2012, heat content anomalies became positive and increased in magnitude across eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.





# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index continued to show a somewhat weak amplitude with some eastward propagation during the past week.





# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS forecasts show a strengthening MJO signal with eastward propagation to Africa and the Indian Ocean over the next two weeks.







# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates negative anomalies (enhanced convection) shifting east to the east Pacific, Mexico and the Caribbean. Positive anomalies (suppressed convection) spreads east from the Indian Ocean (Week-1) to the western Pacific (Week-2).




# Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



This forecast indicates enhanced convection developing across the eastern Pacific and Central America during the period. Suppressed convection is forecast to slowly expand east to the western Pacific.









# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>





# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>