



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
September 24, 2012**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The latest observations indicate that the MJO continued to weaken as many indicators are now less coherent as compared to the last couple of weeks. The enhanced phase remained over the western Pacific.**
- **The majority of dynamical model MJO index forecasts, however, indicate a larger amplitude MJO signal emerging across the western Pacific during the upcoming two week period. Model spread and uncertainty remain somewhat high, similar to last week.**
- **Based on the latest observations, the MJO is forecast to remain weak, but there is some potential for a more organized signal to develop across the western Pacific by the beginning of Week-2.**
- **The MJO may contribute some to enhanced convection across the western Pacific and perhaps the eastern Pacific and Central America during Week-2. Suppressed convection is favored for parts of India, the eastern Indian Ocean and Maritime continent throughout the period. Also, the chances for tropical cyclogenesis remain elevated across the western Pacific.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



The monsoon flow across the Indian Ocean and southern Asia weakened during the last five days.


Winds across the central Pacific have been near average during the past five days.

Westerly anomalies persisted over the tropical Atlantic and western Africa.



850-hPa Zonal Wind Anomalies ($m s^{-1}$)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



The MJO (dashed line) contributed to westerly anomalies near the Date Line and across the western hemisphere during April. Anomalies became more persistent in most areas during the remainder of April and May.

Strong westerly anomalies developed across the eastern Pacific in mid-June and shifted westward (black solid line) and contribute to weakening the trade winds.


Easterly anomalies persisted near 80E for much of August and September.

During September, westerly anomalies developed near 140E and the easterlies have decreased near the Date Line.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



In late August into early September, enhanced convection was observed over many areas of the northern Indian Ocean, southern Asia and the western Pacific. Enhanced convection was also evident across the eastern Pacific and Caribbean Sea.


During early-to-mid September, enhanced convection continued for southern Asia and parts of the western Pacific while strong suppressed convection developed across the eastern equatorial Indian Ocean. Widespread suppressed convection was observed over the Americas.

Suppressed convection continued across the eastern equatorial Indian Ocean and the Americas while enhanced convection over the western Pacific generally shifted northward.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
8-Apr-2012 to 23-Sep-2012 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

From late May into September, eastward propagation of both enhanced and suppressed convection is evident across the eastern hemisphere (alternating dashed and dotted lines).

Atmospheric Kelvin wave activity also played a large role in the pattern of anomalous convection across the Pacific and western Hemisphere during much of this period, especially June and July.

During early September, the strongest enhanced convection shifted eastward to the western Pacific, just west of Date Line.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Longitude

The MJO strengthened in late January and continued into mid-April as indicated by alternating negative (dashed lines) and positive (dotted lines) anomalies with eastward propagation.

Beginning in late April, anomalies became weaker and less coherent than earlier in the year.

Eastward propagation was once again evident from late May into September associated with the MJO, as well as atmospheric Kelvin wave activity, which at times resulted in fast eastward propagation of observed anomalies.


In mid-September, anomalies decreased and eastward propagation less clear.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern reflects anomalous upper-level convergence across the Americas and the Atlantic with smaller areas of anomalous upper-level divergence over Africa, the Indian ocean and parts of the Pacific.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies (blue boxes) decreased considerably during the past five days over the eastern Pacific and Americas.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



The MJO contributed to alternating westerly (dotted lines) and easterly (dashed lines) anomalies into mid-April.


Anomalies were less coherent during much of April and May as the MJO was weak.

Westerly anomalies shifted eastward across the Pacific during July and early August.

Westerly anomalies increased markedly across the eastern Pacific in early September and have persisted.



Weekly Heat Content Evolution in the Equatorial Pacific



From July 2011 through February 2012, heat content was below average in the central and eastern equatorial Pacific.

From March into July 2012, heat content anomalies became positive and increased in magnitude across eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

Positive anomalies decreased across the eastern Pacific during late August and September.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index amplitude remained low during the past week with no eastward propagation.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecasts the MJO signal to increase in amplitude with some eastward propagation during the next two weeks.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates enhanced (suppressed) convection developing across the western Pacific (Indian Ocean) mainly during Week-2.






Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The forecast indicates weak anomalous convection during the next two weeks.







MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>