



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
January 7, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO strengthened considerably during the past week with some eastward propagation across the Maritime Continent.**
- **Dynamical and statistical model MJO index forecasts are in good agreement that the MJO will propagate through phases 5 and 6 over the next two weeks.**
- **Based on the latest observations and MJO index forecasts, the MJO is forecast to propagate through phases 5 and 6 (to the Western Pacific) during the next two weeks.**
- **The MJO may contribute to enhanced (suppressed) convection across the Maritime Continent and western equatorial Pacific (Indian Ocean) during the period.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies shifted east in the Indian Ocean.

Easterly anomalies also shifted east over the past several days, consistent with an active MJO.


Westerly anomalies continued over the eastern tropical Pacific during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

Easterly anomalies persisted near 80E for much of August to October (red box).

During September, westerly anomalies developed near 140E and persisted into October.

Easterly anomalies developed west of the Date Line during late October in the west Pacific and have persisted.


Westward propagation (shaded areas, sloping down and to the left) during much of November and early December are primarily due to equatorial Rossby wave activity.

Recently, there has been some eastward propagation of anomalies.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



Early to mid-December featured weakly organized convection over the Indian Ocean and a large area of suppressed convection over the entire West Pacific.


During mid- to late December, convection began to organize over the Indian Ocean, more closely resembling MJO activity. Suppressed convection from the Date Line west appeared less organized, partly due to tropical cyclone activity.

In late December and early January convection shifted east across the Maritime Continent, while suppressed convection intensified near the Date Line.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
22-Jul-2012 to 6-Jan-2013 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

From late July into September, eastward propagation of both enhanced and suppressed convection is evident across the eastern hemisphere (alternating dashed and dotted lines).

The MJO was active during October into November with enhanced convection developing over Africa during mid-October and shifting eastward to the western Pacific by mid-November.

During late November and much of December, convective anomalies were disorganized, in part a consequence of continued weak MJO activity. Suppressed convection persisted near the Date Line during early December, along with some westward propagation of these anomalies.

Recently however, renewed eastward propagation is indicative of more robust MJO activity.

Longitude

Time






200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

Eastward propagation was evident from July into September associated with the MJO (alternating dashed and dotted lines), as well as atmospheric Kelvin wave activity, which at times resulted in fast eastward propagation of observed anomalies.

In mid-September, anomalies decreased and eastward propagation became less clear.

In early October, upper-level divergence (convergence) increased over the Pacific (Indian Ocean) and has shifted eastward throughout October and early November.


During December, anomalies were weaker, with less coherent eastward propagation. Other subseasonal variability (atmospheric Kelvin and equatorial Rossby waves) were also evident during December. Recently, however, more robust eastward propagating upper-level divergent anomalies are evident, largely associated with MJO.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern closely resembles a wave-1 structure, indicative of organized MJO activity. Upper-level convergence is strongest over the Americas, Atlantic Ocean, and Africa. Upper-level divergence is evident over parts of eastern Indian Ocean and Maritime Continent.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies (blue boxes) have been located farther north across the Northern mid-latitudes. A retracted jet over East Asia, during the latter period, is consistent with observed MJO activity.


Recently, strong westerly anomalies have developed over the eastern equatorial Pacific, also consistent with MJO.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westerly anomalies shifted eastward across the Pacific during July and early August.

Westerly anomalies prevailed across the eastern Pacific and Americas for much of September and October, but were replaced by easterly anomalies during mid-October.


Westerly anomalies shifted east to the eastern Pacific in early November, but have alternated between easterly and westerly anomalies since this period. An area of stronger, easterly anomalies was located over the equatorial eastern Pacific, but has now reversed.

Longitude



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



From December 2011 through February 2012, heat content was below average in the central and eastern equatorial Pacific.

From March into July 2012, heat content anomalies became positive and increased in magnitude across eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

Positive anomalies decreased across the eastern Pacific during late August and September.

An oceanic Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November. Recently, negative anomalies, in the central and eastern Pacific, east of the Date Line, have increased in magnitude.

Longitude



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates rapid strengthening and eastward propagation over the past week. However, projection of other modes onto the index remains evident.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The bias-corrected ensemble GFS forecasts indicate an eastward propagation of a strong MJO signal through phases 5 and 6 over the next two weeks.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates eastward propagation toward the western Pacific, consistent with MJO. Drier-than-average conditions are forecast for the Indian Ocean, parts of Africa and South America.







Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





This statistical forecast is in good agreement with the GFS based forecast, with enhanced convection moving into the western Pacific and suppressed convection moving into the Indian Ocean.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>