



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
May 27, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO has remained largely incoherent during the past week.**
- **Dynamical model MJO index forecasts are in good agreement for a generally weak MJO signal during the first part of Week-1, in part related to interference from other tropical sub-seasonal variability, but also from a weakened MJO. The models indicate potential renewed organization of an enhanced phase centered across Africa later in Week-1 and Week-2.**
- **Based on recent observations and model MJO forecasts, the MJO is forecast to become better organized during Week-2 with the enhanced phase centered across the eastern Western Hemisphere and Africa.**
- **Enhanced rainfall is favored during Week-1 across parts of the western and southeastern Indian Ocean as well the eastern Pacific and much of Central America. Suppressed rainfall is favored across an area extending from the central Indian Ocean along the Equator to just west of the Date Line. Conditions for tropical cyclogenesis remain favorable for parts of the East Pacific and Caribbean.**
- **A more organized MJO during Week-2 favors enhanced (suppressed) convection in the western and central Indian Ocean and parts of Africa (West Pacific). The Caribbean is likely to remain active with elevated chances of cyclogenesis lingering into Week-2.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



The wind field has become less coherent across the Indian Ocean and western Maritime Continent.


Easterly anomalies diminished rapidly over the central and western Pacific.

Westerly anomalies continued over the East Pacific during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

Westward propagation (dashed/solid lines sloping down and to the left) of anomalies during much of November and early December were primarily due to equatorial Rossby wave activity as the MJO was then generally weak.

During late December the MJO strengthened (alternating dotted/dashed lines).


During March and early April, anomalies indicate signs of being influenced by equatorial Rossby wave activity with less eastward propagation evident.

The MJO strengthened during early May with eastward propagation of low-level wind anomalies noted. A weaker MJO combined with interference from other sub-seasonal modes has limited eastward propagation recently.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During late April and the beginning of May, convection rapidly developed across the Indian Ocean, while the west-central Pacific experienced suppressed convection.


During early to mid-May, enhanced (suppressed) convection prevailed across the Indian Ocean and Maritime Continent (central/eastern Pacific Ocean).

Most recently the OLR field has become less coherent, dominated by small-scale variability.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
8-Dec-2012 to 25-May-2013 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

During much of December, convective anomalies were disorganized.

The MJO was again a dominant mode of variability across the Tropics from January into March as indicated by the alternating dashed and dotted lines.

Near the end of March, the anomalies show signs of influence from other modes of tropical variability. However, MJO activity reemerged in early April across the Indian Ocean.

During early May, anomalous OLR increased significantly across the Indian Ocean. The MJO signal broke down during the middle of the month.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



Time



Longitude

After some MJO activity at the start of November, anomalies decreased with less coherent eastward propagation during most of November and December. Other modes of subseasonal variability were more prevalent during this period.

As the MJO strengthened in late December, (alternating dashed and dotted lines), anomalies increased in magnitude with more robust eastward propagation indicated during late 2012 to April 2013.

Anomalies became less coherent at times during this period as the influence from other modes of variability are evident in the depicted anomalies, namely during late January into early February, before reorganizing in late February and early March.


The velocity potential anomalies were more coherent only briefly during early May and have since broken down.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern has remained largely incoherent, at least in terms of MJO-related anomalies. Upper-level divergence is observed across the East Pacific just north of the Equator associated with an ongoing active ITCZ. Upper-level convergence is strongest over the West Pacific north of the Equator.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Large-scale tropical anomalies have dissipated.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Eastward propagation of westerly wind anomalies associated with the MJO is evident beginning in late December and continuing into April 2013. Some propagation of easterly anomalies is evident during late January and early February.

During March and early April, anomalies were influenced by westward moving features over the central and western Pacific.

Westerly anomalies shifted east of the Date Line during early May. The rapid phase speed suggests the influence of a higher-frequency Kelvin wave.



Weekly Heat Content Evolution in the Equatorial Pacific



From March into August 2012, heat content anomalies became positive and increased in magnitude across the eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

An oceanic Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates a weak signal during the past week to ten days.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The bias-corrected ensemble GFS indicates a weak signal early, emerging in Phase 1 later.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts suppressed (enhanced) convection developing across the eastern Indian Ocean and Maritime Continent (east Pacific, central America, and western Caribbean Sea) during the next two weeks.





Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





This statistical forecast indicates small anomalies across the global tropics.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>