



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
July 8, 2013**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **A coherent MJO signal propagated eastward over the Western Hemisphere into the western Indian Ocean during the previous week.**
- **Dynamical and statistical model MJO index forecasts exhibit a wide variance of solutions, with some maintaining a coherent MJO and others weakening the signal.**
- **Based on recent observations and model MJO forecasts, the MJO is forecast to propagate eastward over the Indian Ocean, with continued influence from other modes of tropical variability.**
- **For Week-1, enhanced convection is favored across the northern and central Indian Ocean, western and central Africa, western Mexico, and the southeastern United States. Suppressed convection is favored over the western Pacific. Tropical cyclogenesis is possible over the central North Atlantic.**
- **During Week-2, enhanced convection is favored over the eastern Indian Ocean and the Maritime Continent, while odds for suppressed convection are enhanced across the eastern Pacific. Conditions are expected to become increasingly favorable for tropical cyclone formation over the western North Pacific.**

**Additional potential impacts across the global tropics and a discussion for the U.S. are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies persisted over the equatorial and northern Indian Ocean with diminished intensity.

Weak easterly anomalies developed over the Maritime Continent.


Westerly anomalies persisted over the eastern Pacific and Atlantic basins with diminished intensity.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



During early January the MJO strengthened (alternating dotted/dashed lines).

During March and early April, anomalies indicate signs of being influenced by equatorial Rossby wave activity with less eastward propagation evident.


The MJO strengthened during early May, with the signal becoming more incoherent later in the month.

The MJO strengthened again in late June, with eastward propagation of low-level westerly wind anomalies noted. More recently, other sub-seasonal modes have limited eastward propagation.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During early June, a more coherent pattern of enhanced (suppressed) convection developed over the Indian Ocean and Maritime Continent (central and eastern Pacific and Africa).

The enhanced convective anomalies propagated eastward over the Maritime Continent and the western Pacific during mid to late June.


During late June and early July, enhanced (suppressed) convective anomalies propagated eastward into the eastern Pacific, western Atlantic, and Africa (Indian Ocean and western Pacific). Enhanced convective anomalies persisted over the maritime continent.



# Outgoing Longwave Radiation (OLR)

## Anomalies (2.5°N-17.5°N)

Time  
↓



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO was a dominant mode of variability across the Tropics from January into March as indicated by the alternating dashed and dotted lines.

Near the end of March, the anomalies show signs of influence from other modes of tropical variability. However, MJO activity reemerged in early April across the Indian Ocean.

During early May, OLR decreased significantly (stronger negative anomalies) across the Indian Ocean. The MJO signal quickly broke down.

Recently, enhanced convection propagated eastward from the Maritime Continent and western Pacific into the eastern Pacific and Caribbean. Suppressed convection increased over the Indian Ocean and western Pacific.

Longitude




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



Longitude

The MJO strengthened in late December, (alternating dashed and dotted lines) and anomalies increased in magnitude with more robust eastward propagation indicated during late 2012 to April 2013.

Anomalies became less coherent at times during late January and early February as the influence from other modes of variability are evident in the depicted anomalies. Some reorganization is evident in late February and early March.

The velocity potential anomalies were more coherent only briefly during early to mid-May.

Recently, the signal is more coherent and consistent with a canonical MJO footprint, although other modes are still apparent.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern is consistent with the MJO convectively-active phase over Africa and the Indian Ocean, surrounded by regions of large scale suppressed convection over the Pacific and western Atlantic basins. Other modes of tropical variability remain evident as well, including a developing tropical cyclone over the eastern Pacific Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly (easterly) upper-level wind anomalies increased over the western and central Indian Ocean (Gulf of Guinea).



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Eastward propagation of westerly wind anomalies associated with the MJO continued into April 2013. Some propagation of easterly anomalies is evident during late January and early February.


During March and early April, anomalies were influenced by westward moving features over the central and western Pacific.

Westerly anomalies shifted east of the Date Line during both early May and early June. The rapid phase speed suggests the influence of higher-frequency Kelvin waves. Recently, easterly anomalies propagated eastward over the Western Hemisphere, with influence from westward propagating features evident over the Maritime Continent.



# Weekly Heat Content Evolution in the Equatorial Pacific

Time  
↓



Longitude

Through August 2012, heat content anomalies became positive and increased in magnitude across the eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

An oceanic Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Positive anomalies increased over the eastern and central Pacific during June 2013.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The eastward propagation of the MJO signal slowed as the convectively active phase emerged over the western Indian Ocean, indicating interactions with other modes of intraseasonal variability.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS indicates some eastward propagation later in Week-1, with the signal becoming incoherent during Week-2.









# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts enhanced (suppressed) convection over the Indian Ocean (western Pacific) during Week-1. Convective anomalies are forecast to decrease during Week-2 as the signal becomes less coherent.




# Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



This statistical forecast indicates a slightly more coherent MJO, with eastward propagation of the convectively active phase over the Maritime Continent during Week-2.







# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>