



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
July 22, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO signal propagated over the Maritime Continent during the previous week, while the influence of other modes of variability is apparent in the global tropical circulation.**
- **Dynamical model MJO forecasts indicate a weak or incoherent MJO pattern during the upcoming two weeks. A few statistical models forecast eastward propagation of an MJO signal.**
- **Based on recent observations and statistical forecasts, the MJO is forecast to maintain coherent and propagate eastward over the western Pacific during the upcoming two weeks.**
- **Enhanced convection is favored over India, the Bay of Bengal, and the Maritime Continent during Week-1, while suppressed convection is more likely over the equatorial Indian Ocean. Enhanced monsoon activity is anticipated over northwestern Mexico and the southwestern U.S., while suppressed convection is favored over the Gulf of Mexico.**
- **Tropical cyclone formation is favored over the eastern Pacific basin during both Week-1 and Week-2. Some dynamical models indicate an unusual westward extent to the potential tropical cyclone path. Consistent with the MJO phase, tropical cyclogenesis is also possible over the western North Pacific and South China Sea during the upcoming two weeks.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Low-level easterly wind anomalies increased over the equatorial Indian Ocean.


Easterly anomalies persisted over the western Pacific.

Westerly wind anomalies increased over the eastern Pacific.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

During February, MJO activity was apparent in the low-level wind pattern (alternating dashed and dotted lines).

During March and early April, anomalies indicate signs of being influenced by equatorial Rossby wave activity with less eastward propagation evident.


The MJO strengthened during early May, with the signal becoming more incoherent later in the month.

The MJO strengthened again in late June, with eastward propagation of low-level westerly wind anomalies noted. Recently, other modes of tropical intraseasonal variability are apparent in the wind field.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



The MJO enhanced phase propagated eastward over the Western Hemisphere during late June, with enhanced convection apparent over the eastern Pacific.


During early July, the MJO enhanced phase propagated over the Indian Ocean, promoting increased convection along and just north of the equator.

During mid-July, the MJO signal propagated over the Maritime Continent. Other modes of convective variability are apparent in the OLR field, including enhanced convection over North America.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°N-17.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
3-Feb-2013 to 21-Jul-2013 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO was a dominant mode of variability across the Tropics from January into March as indicated by the alternating dashed and dotted lines.

Near the end of March, the anomalies show signs of influence from other modes of tropical variability. However, MJO activity reemerged in early April across the Indian Ocean.

During early May, OLR decreased significantly (stronger negative anomalies) across the Indian Ocean. The MJO signal quickly broke down.

The MJO signal emerged during June and early July, as indicated by the alternating dashed and dotted lines. Other modes of variability are apparent as well, such as a Rossby wave over the Maritime continent and a Kelvin Wave over the central Pacific.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

The MJO strengthened in late December, (alternating dashed and dotted lines) and anomalies increased in magnitude with more robust eastward propagation indicated during late 2012 to April 2013.

Anomalies became less coherent at times during late January and early February as the influence from other modes of variability are evident in the depicted anomalies. Some reorganization is evident in late February and early March.

The velocity potential anomalies were more coherent only briefly during early to mid-May.


Recently, the signal is more coherent and consistent with a canonical MJO footprint, although other modes are still apparent.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern remains somewhat consistent with the MJO pattern, though other modes of variability are also apparent in the upper-level circulation, such as the convectively active phase of a Kelvin Wave over the eastern Pacific.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly wind anomalies increased (decreased) over the western (central) Pacific. Easterly anomalies weakened over the equatorial Atlantic.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Eastward propagation of westerly wind anomalies associated with the MJO continued into April 2013. Some propagation of easterly anomalies is evident during late January and early February.


During March and early April, anomalies were influenced by westward moving features over the central and western Pacific.

During June and early July, eastward propagation of upper-level zonal wind anomalies was observed. More recently, influence from other modes of tropical intraseasonal variability produced a less coherent upper-level zonal wind anomaly pattern.



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



Longitude

Through August 2012, heat content anomalies became positive and increased in magnitude across the eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

An oceanic Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Positive anomalies increased over the central Pacific during June and early July 2013.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the previous week, a coherent MJO pattern was not apparent on the Wheeler and Hendon phase diagram due to the influence of other modes of convective variability.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS indicates an incoherent MJO signal during both Week-1 and Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts little or no anomalous convection. Suppressed (enhanced) convection is forecast over the western Pacific (eastern Pacific, Atlantic, and equatorial Indian Ocean) during Week-2.




Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



This statistical forecast indicates persistent anomalous convection over the western Pacific, with suppressed convection spreading over the northern Indian Ocean during Week-2.







MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>