



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
October 28, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remained weak during the past week with other types of tropical intraseasonal variability influencing anomalous tropical convection.**
- **Dynamical model MJO index forecasts and statistical guidance indicate a continued weak signal during the next two weeks.**
- **Based on recent observations, statistical tools, and dynamical forecasts, the MJO is forecast to remain weak. Other types of subseasonal tropical variability are likely to influence the pattern of tropical convection.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>





850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies developed over the central Indian Ocean.

Westerly anomalies intensified and expanded over the western North Pacific.

Easterly anomalies developed near the Date Line during the past five days.


Westerly anomalies intensified over the Eastern Pacific during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



The MJO strengthened during June and continued to be significant until mid-July with fast eastward propagation.


During late July through mid-August, other types of subseasonal variability strongly contributed to the observed anomalies. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During mid-October, a Rossby wave is evident from 160E to 100E with westerly anomalies persisting across the Western Hemisphere.

Recently the pattern is less coherent, exhibiting evidence of high frequency variability.



OLR Anomalies – Past 30 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During late September and early October, suppressed convection expanded over the Indian Ocean while enhanced convection was evident across the west Pacific, associated with tropical cyclone activity.


Suppressed convection persisted across the equatorial Indian Ocean during early October, while the tropical cyclones remained active across the west Pacific and the Bay of Bengal.

During mid-October, enhanced convection developed across the equatorial Indian Ocean and persisted across the western North Pacific. Suppressed convection was observed across parts of the Maritime Continent.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°N-17.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
12-May-2013 to 27-Oct-2013 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO strengthened once again during June and continued into early July.

MJO was active during September, with the enhanced phase propagating east over the west Pacific Ocean, and the suppressed phase strengthening over the Indian Ocean.

Tropical cyclone activity contributed to the persistence of enhanced convection across the west Pacific as well as a weakened suppressed phase further west.

Longitude

Time
↓




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

The MJO was active for through early May 2013 period as shown by generally alternating positive (brown) and negative (green) anomalies with clear eastward propagation.

The MJO was less coherent during much of May, then strengthened once again during June and the first half of July before weakening by the end of the month.

The MJO was not active during late July and much of August, but strengthened during September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.


Recently, the upper-level velocity potential has exhibited little MJO-related variability.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern remained incoherent during the past week as other modes of tropical intraseasonal variability were apparent. The largest positive anomalies (upper-level convergence) are observed over eastern Asia and the Atlantic Ocean, while upper-level divergence covers parts of Africa, the western Equatorial Pacific, and the Eastern Pacific.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly upper-level zonal wind anomalies diminished over the Indian Ocean and intensified over the Maritime Continent.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

Eastward propagation of wind anomalies associated with the MJO (dotted and dashed lines) continued into May 2013.

The MJO strengthened during June and its influence continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.


During August, westerly wind anomalies were generally persistent just west of the Date Line.

Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Westerly wind anomalies spread eastward near the Maritime Continent. The pattern is inconsistent with robust MJO activity.



Weekly Heat Content Evolution in the Equatorial Pacific



An oceanic downwelling Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November 2012.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Positive anomalies have persisted across most of the basin since June 2013, with evidence of an oceanic downwelling Kelvin wave in late August and September.

The development of an oceanic Kelvin wave is observed during early October.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicated little to no propagation of a weak signal during the past week. Movement in the index is likely reflecting other modes of tropical variability.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS indicates a strengthening signal during the next two weeks, although the signal is not consistent with MJO related activity and likely results from other modes projecting onto the MJO diagnostic tools.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts suppressed convection intensifying across the Maritime Continent and west Pacific during the next two weeks, while enhanced convection is predicted across the east Pacific, South America, and Africa.




Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The constructed analog MJO forecast continues the weak signal, with no anomalies predicted.







MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>