



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
November 25, 2013**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained incoherent during the previous week, with influence from other types of coherent tropical subseasonal variability dominating the anomalous convection and circulation pattern.**
- **Dynamical model MJO index forecasts indicate little signal over the next week as other types of subseasonal variability continue to strongly influence the tropical circulation. There is considerable spread among the dynamical model forecasts during Week-2. Statistical forecasts suggest only a weak MJO signal.**
- **Based primarily on the latest observations and some dynamical model guidance, the MJO is not forecast to contribute significantly to anomalous tropical convection at the current time.**

**Additional potential impacts across the global tropics and a discussion for the U.S. are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Weak easterly anomalies persisted across parts of the equatorial Indian Ocean.


Easterly anomalies persisted over the western Pacific and Maritime Continent north of the equator, while westerly anomalies expanded across the remainder of the Maritime Continent and northern Australia.

Westerly anomalies continued over the eastern Pacific during the past five days, and increased over the equatorial Atlantic.



# 850-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

The MJO strengthened during June and continued until mid-July with fast eastward propagation.

During late July through mid-August, the MJO was weak. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.


During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.

During the first half of November, MJO activity was not coherent as multiple westward moving features dominated the pattern of low level zonal wind anomalies.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During late October, enhanced convective activity was observed over the western and eastern Pacific north of the equator, associated with tropical cyclone activity. Enhanced (suppressed) convection was observed over South Asia (South China Sea and eastern Indian Ocean).


During early November, enhanced convection associated with tropical cyclones continued over the western Pacific. Enhanced (suppressed) convection was observed over the western Indian Ocean (near the Date Line)

During mid-November, enhanced (suppressed) convection was observed over the Bay of Bengal, Maritime Continent, Australia, and the Gulf of Mexico (western Pacific, Caribbean, eastern Brazil).



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°N-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
9-Jun-2013 to 24-Nov-2013 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

**The MJO strengthened once again during June and continued into early July.**

**MJO was active during late August and September with the enhanced phase propagating eastward over the western Pacific Ocean, while the suppressed phase strengthened over the Indian Ocean.**

**Tropical cyclone activity contributed to the persistence of enhanced convection across the West Pacific as well as a weakened suppressed phase further west.**

**Recently, eastward propagating convective anomalies across the western Indian Ocean were interacting with westward moving modes over the Maritime Continent.**


**Longitude**



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



The MJO was active (alternating dashed and dotted lines) during June and early July before weakening at the end of the month.

The MJO was not active during late July and much of August, but strengthened during late August and September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.

During late October and early November, some coherent eastward propagation on the fast side of the MJO envelope of phase speeds was observed, but the signal rapidly broke down during mid-November.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern remains incoherent from an MJO perspective, with areas of large scale upper level divergence (green contours) over South America, the Atlantic, Africa, the Maritime Continent, and Australia contrasting with areas of large scale upper level convergence (brown contours) over the eastern Pacific and Indian Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly upper-level zonal wind anomalies continued over the western Pacific, while easterly anomalies developed over the eastern Pacific with anomalous cross-equatorial flow.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude


The MJO strengthened (alternating dotted and dashed lines) during June and its influence continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.

During August, westerly wind anomalies were generally persistent just west of the Date Line. Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Most recently, anomalies of alternating sign have continued over the eastern Pacific, due in part to extratropical Rossby waves breaking into the Tropics.



# Weekly Heat Content Evolution in the Equatorial Pacific



An oceanic downwelling Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November 2012.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Evidence of oceanic downwelling Kelvin waves are seen in late August and October/November.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO signal as measured by the RMM index has exhibited no coherence during the previous two weeks.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS indicates little MJO signal during the Week-1 period, with an emerging signal over the Indian Ocean towards the end of Week-2.









# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts anomalous convection over the western Pacific and Maritime Continent during Week-1, with enhanced (suppressed) convection over the Indian Ocean (western Pacific) during late Week-2.





# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The constructed analog MJO forecast indicates some eastward propagation of enhanced convective anomalies over the Maritime Continent and western Pacific.



# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>