



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
January 12, 2015**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO remained active during the past week with the enhanced phase centered over the West Pacific.**
- **Dynamical model MJO index forecasts and statistical forecasts generally indicate a continued MJO signal during the next two weeks, although most indicate a decrease in amplitude.**
- **Based on the latest observations and several forecast tools, the MJO is forecast to remain active during the next two weeks with the enhanced convective phase shifting into the Western Hemisphere.**
- **The MJO is expected to contribute to enhanced (suppressed) rainfall across parts of the Pacific (Indian) Ocean during Week-1. An elevated threat for tropical cyclone development exists for the Mozambique Channel early in Week-1.**
- **During Week-2, the MJO is expected to favor enhanced rainfall across parts of the South Pacific, while suppressed rainfall shifts east from the Indian Ocean to the Maritime Continent.**

**A forecast map of potential impacts across the global Tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies expanded east across the Maritime Continent and West Pacific during the past five days.

Easterly anomalies strengthened across the East Pacific.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

From late July to August, an envelope of westerly wind anomalies shifted eastward across the Pacific associated with weak MJO activity (dashed line). Embedded within this envelope were frequent and strong westward moving high frequency features (red arrows) over the eastern and central Pacific (western Pacific, Maritime Continent, and Indian Ocean).


A westerly wind burst was observed near the Date Line during mid-October

MJO activity was observed beginning in late November into December and a second stronger event has evolved during late December and early January 2015. Westerly anomalies have shifted east to 160E.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During mid-December, suppressed convection remained near the Date Line, while convection became enhanced over the Indian Ocean.


During late December, the pattern was more coherent, with enhanced convection shifting east from the Indian Ocean to the Maritime Continent. Suppressed convection was observed over Brazil.

A coherent pattern continued into the New Year with enhanced (suppressed) convection shifting east across the Maritime Continent (Indian Ocean). Suppressed convection persisted over Brazil.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
27-Jul-2014 to 11-Jan-2015 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

The MJO became more organized during July and August, as enhanced and suppressed convection phases shifted eastward from the Indian Ocean to the Pacific Ocean during this period (dashed/dotted lines).

The pattern became less coherent with respect to canonical MJO activity by September and the MJO remained weak till late November (red box).

The MJO strengthened in late November with alternating areas of enhanced and suppressed convection moving from the Indian Ocean to the Date Line through early January.

**Longitude**

**Time**  
↓




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



A slow eastward progression of negative anomalies was observed during the summer across the Indo-Pacific warm pool and central-eastern Pacific (red box).

The pattern became more organized during July as the MJO strengthened at this time (dashed and dotted lines) as a more coherent “Wave-1” canonical MJO-like structure developed and shifted eastward with time.

The MJO weakened and remained incoherent through September and October.

During November the MJO strengthened as indicated by eastward propagation of anomalies with the enhanced phase entering the west-central Pacific by early December. Rapid eastward propagation has continued, likely due to Kelvin wave activity, and strong upper level divergence is now centered over the West Pacific.

Longitude






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The upper-level anomalous velocity potential spatial pattern is coherent with an active MJO. Upper-level divergence is now centered over the West Pacific with upper-level convergence extending from South America to the Indian Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Easterly anomalies diminished over the Indian Ocean, while strong mid-latitude interactions are evident over the eastern Pacific.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




A slow, eastward progression of westerly anomalies is evident over the Maritime Continent and western Pacific during August. Some westward propagation is noticeable during September and early October.

Easterly wind anomalies persisted east of the Date Line from late October through early December.

During early January, easterly anomalies shifted from 90E to 140E.



# Weekly Heat Content Evolution in the Equatorial Pacific



A strong downwelling event began in January 2014 and propagated across the Pacific reaching the South American coast by May 2014.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies decreased during June and July associated with an upwelling Kelvin wave (dotted line).

Warm anomalies increased across much of the Pacific basin due to another moderate downwelling Kelvin wave traversing the Pacific during October and November 2014. The upwelling phase is now evident in the central Pacific.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index depicts an eastward propagating strong amplitude MJO signal during the past week.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a continued MJO signal shifting east across the Pacific Ocean during Week-1. The signal decreases in amplitude in Week-2.









# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The GEFS mean MJO index based OLR anomaly forecast depicts a weakening of anomalous convection during the next two weeks.




# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The constructed analog forecast depicts a continued eastward propagation of anomalous convection which is typical during an ongoing MJO.



# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>