



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
February 16, 2015**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **Observations and MJO monitoring indices, such as the CPC MJO index based on upper-level velocity potential, indicate a fast-moving MJO signal with the enhanced phase centered over the Western Hemisphere. Other indices indicate little to no MJO signal.**
- **The dynamical forecast models generally indicate a weak signal during Week-1, with a highly uncertain strengthening signal during Week-2. Statistical tools indicate little to no signal, which is typical for those tools when the initial state is weak.**
- **During the next two weeks there are increased chances for tropical cyclone formation near Australia and across the central, South Indian Ocean. Other modes of variability are expected to contribute to potentially enhanced convection over the Maritime Continent and near Australia through Week-2.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Easterly (westerly) anomalies intensified over the Timor Sea (western North Pacific).

Easterly anomalies developed near the Date Line while westerly anomalies intensified slightly over the eastern Pacific.



850-hPa Zonal Wind Anomalies ($m s^{-1}$)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



In early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.


During November and December, easterly anomalies were persistent from 120E to near the Date Line. Westerly anomalies replace those easterly anomalies during January. Easterly anomalies disrupted the signal during early February. Westerly anomalies returned to the Western Pacific during late January.

Most recently, the strongest anomalies are over the eastern Pacific, with weak easterly anomalies near the Date Line.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During mid to late January, enhanced (suppressed) convection was mostly east (west) of the Date Line. MJO activity had mostly waned by this time.


From late January to February, enhanced convection developed over the West Pacific. Below average convection was measured over northern Australia, the Maritime Continent, and eastern Brazil.

Convection remained above (below) average over the West Pacific (Maritime Continent and northern Australia). Suppressed convection persisted over Hawaii, with slightly enhanced convection over northwest South America.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
30-Aug-2014 to 15-Feb-2015 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

Some MJO activity was evident during August, as enhanced and suppressed convection phases shifted eastward from the Indian Ocean to the Pacific Ocean during this period (dashed/dotted lines).

The pattern became less coherent with respect to canonical MJO activity by September and the MJO remained weak till late November (red box).

The MJO strengthened in late November with alternating areas of enhanced and suppressed convection moving from the Indian Ocean to the Date Line through January.

Recently, enhanced convection has been almost stationary near the West Pacific.

Longitude


Time
↓



200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



Time
↓

Longitude

The pattern became more organized during late July as the MJO strengthened. This is observed as a coherent “Wave-1” canonical MJO-like structure that shifted eastward with time.

The MJO weakened and remained incoherent through September and October.

Beginning in November the MJO strengthened as indicated by eastward propagation of alternating anomalies into January 2015. At times, the signal has been dominated by faster-moving Kelvin wave variability, but from late December through mid-January the signal has been more consistent with canonical MJO activity.


Since mid-January, the signal has broken down, with other modes of variability dominating the upper-level velocity potential anomaly pattern.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The spatial pattern of upper-level velocity potential anomalies has become incoherent with MJO activity. The pattern has a North-South oriented structure, over the Eastern Hemisphere, and indications of other modes of variability over the Western Hemisphere.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies over the Maritime Continent have weakened while westerly anomalies over Africa and the Indian Ocean, and the East Pacific have intensified.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

A slow, eastward progression of westerly anomalies is evident over the Maritime Continent and western Pacific during August (black dashed line). Some westward propagation is noticeable during September and early October over the eastern Pacific.


Easterly wind anomalies persisted east of the Date Line from late October through early December (red box).

During late December through the present, westerly anomalies increased in coverage and intensity from 120W to 80W, similar to September and October 2014.

Westerly anomalies have also been persistent over the Indian Ocean.



Weekly Heat Content Evolution in the Equatorial Pacific



A strong downwelling event began in January 2014 and propagated across the Pacific reaching the South American coast by May 2014.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies decreased during June and July associated with an upwelling Kelvin wave (dotted line).

Warm anomalies increased across much of the Pacific basin due to another moderate downwelling Kelvin wave traversing the Pacific during October and November 2014. The upwelling phase is now evident in the central and eastern Pacific.

Warm anomalies began increasing again near the Date Line during late January and early February.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index over the past week has shown no coherent signal and westward movement of convection.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS ensemble RMM Index forecasts do not depict a signal consistent with MJO activity.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The GEFS RMM Index based OLR anomalies forecast depicts westward propagation of the signal during Week-1, with a signal emerging over the eastern Indian Ocean during Week-2.




Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The statistical forecast depicts some eastward propagation of an MJO signal during the upcoming two week period. The initial signal is weak, so confidence in this method is lower than usual.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>