



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
March 2, 2015**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **Recent observations and MJO indices depict little to no signal consistent with canonical MJO activity, with a weak anomaly pattern across multiple diagnostic tools.**
- **Other modes of tropical convective variability, including Kelvin Waves and Equatorial Rossby Waves, continue to influence the pattern.**
- **The dynamical forecast models indicate a strengthening MJO signal during Week-1, with its enhanced convective phase propagating across the West Pacific during Week-2. Statistical models favor a continued weak signal.**
- **Although there is uncertainty on how strong the MJO signal becomes during the next two weeks, the MJO may contribute to enhanced convection shifting eastward from the Maritime Continent to the West Pacific during the next two weeks.**

**Additional potential impacts across the global tropics and a discussion for the U.S. are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies shifted north to the equatorial Indian Ocean.


Westerly anomalies expanded westward across the Pacific to the Date Line, associated with Equatorial Rossby Wave activity.

Westerly anomalies persisted over the equatorial eastern Atlantic



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

In early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.


During November and December, easterly anomalies were persistent from 120E to near the Date Line. Westerly anomalies replace those easterly anomalies during January. Easterly anomalies disrupted the signal during early February. Westerly anomalies returned to the Western Pacific during late January.

The largest anomalies (westerly) are observed east of the Date Line, and are associated with an ERW.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



**From the end of January to early February, enhanced convection developed over the West Pacific, partly associated with tropical cyclone activity. Suppressed convection overspread the southern Maritime Continent and northern Australia, and persisted over Hawaii.**


**During mid-February, OLR anomalies were relatively small although suppressed convection persisted across much of northern Australia.**

**During late February, convection increased across northeast Brazil, parts of the Maritime Continent, and much of Australia. Suppressed convection was observed across the eastern Indian Ocean.**



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
14-Sep-2014 to 1-Mar-2015 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

The OLR anomaly pattern became less coherent with respect to canonical MJO activity during September and the MJO remained weak until mid-November (red box).

The MJO strengthened in late November with alternating areas of enhanced and suppressed convection moving from the Indian Ocean to the Date Line through January.

Enhanced convection persisted just west of the Date Line during late January and early February as the MJO signal broke down.

More recently, convective anomalies were generally small as the MJO signal remains incoherent.


**Longitude**



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



The MJO was incoherent from mid-September through October.

Beginning in November the MJO strengthened as indicated by eastward propagation of alternating anomalies into January 2015. At times, the signal was dominated by faster-moving variability on the Kelvin Wave time scale, but from late December through mid-January the signal was more consistent with canonical MJO activity.

Since mid-January, the signal has broken down, with other modes of variability dominating the upper-level velocity potential anomaly pattern.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The spatial pattern remains inconsistent with canonical MJO activity, with no robust Wave-1 pattern evident.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




An Upper-level anticyclone shifted west to the northeast Pacific over the middle latitudes by the end of February, while westerly anomalies remained enhanced across the eastern equatorial Pacific.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westward propagating features are noticeable during September and early October over the eastern Pacific.


Easterly wind anomalies persisted east of the Date Line from late October through early December (red box).

During late December through the present, westerly anomalies increased in coverage and intensity from 120W to 80W, similar to September and October 2014. Westerly anomalies also became more persistent over the Indian Ocean.

More recently, westward propagation of westerly anomalies is evident over the eastern Pacific.



# Weekly Heat Content Evolution in the Equatorial Pacific



A strong downwelling event began in January 2014 and propagated across the Pacific reaching the South American coast by May 2014.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies decreased during June and July associated with an upwelling Kelvin wave (dotted line).

Warm anomalies increased across much of the Pacific basin due to another moderate downwelling Kelvin wave traversing the Pacific during October and November 2014. The upwelling phase was evident in the central and eastern Pacific during January.

Warm anomalies associated with another downwelling KW are evident over the central Pacific.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index during the past weeks has shown no coherent signal.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GFS ensemble RMM Index forecasts depict an increase in the amplitude of a MJO signal during Week-1 with eastward propagation during Week-2 .









# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The GEFS RMM Index based OLR anomalies forecast depicts larger anomalies during Week-1, with enhanced convection shifting east across to the Date Line by the end of Week-2.




# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The statistical forecast depicts a weak signal during the next two weeks.



# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>