

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
9 November 2015

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

MJO indices indicate a continued moderate strength signal with some eastward propagation across the Indian Ocean during the past week.

During Week-1, the MJO is expected to destructively interfere with ENSO, but weaken by Week-2.

Dynamical and statistical models indicate a weakening MJO signal during Week-2. The ongoing El Niño is likely to return as the major contributor to anomalous convection during the next two weeks.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies weakened over the West Pacific but strengthened over the Indian Ocean.

Westerly anomalies weakened across the central and eastern Pacific.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

The red box highlights the persistent low-frequency westerly wind anomalies associated with ENSO. Some transient variability is observed as well.

A robust MJO event was observed in late June through mid-July, constructively interfering with the background state.

The background ENSO remains the primary signal, but tropical cyclone activity across much of the Pacific continues to influence the pattern.

An eastward shift in the pattern was observed in late October, related to activity in the MJO time band.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Enhanced convection continued across parts of the central and eastern Pacific, while suppressed convection remained centered on the Maritime Continent during early October.

Enhanced convection waned slightly over the Central Pacific, while suppressed convection expanded slightly over the West Pacific. Enhanced convection also intensified over the Indian Ocean during mid-October.

During late October and early November, suppressed convection expanded eastward to near Hawaii, expanded over the Indian Ocean, and weakened near the Maritime Continent. Enhanced (suppressed) convection also waned over the East Pacific (South America).



Outgoing Longwave Radiation (OLR) Anomalies (5°N-5°S)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)


Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Since April, the ongoing El Niño is observed (red box) as a tendency toward a dipole of anomalous convection extending from the Maritime Continent (suppressed) to the East Pacific (enhanced).

During June and early July, the MJO become active, interfering with the ENSO signal at times.

Since July, the MJO has remained weak, with strong El Niño conditions and tropical cyclone activity dominating the pattern.

A couplet of enhanced/suppressed convection intensified over the western Indian Ocean/Maritime Continent early in October, but seems to be weakening as some of the signal moves eastward.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

The MJO strengthened in March as seen in the upper-level velocity potential anomalies.


The ongoing ENSO state is highlighted by the red box, showing anomalous divergence over the central and eastern Pacific. This pattern has only been temporarily interrupted by strong Kelvin wave/MJO activity at times.

During June and early July, a high-amplitude MJO event was observed, constructively interfering with the El Niño signal in early July.


From July through early October, a generally stationary pattern, reflective of El Niño conditions, was observed.

During late October, there was an eastward shift in the pattern, and recently, the pattern is much noisier.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The upper-level velocity potential anomaly pattern resembles a wave-2 pattern, with weaker anomalies than in prior weeks. The pattern is noisier than in prior weeks, but still resembles an ongoing El Nino.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation


200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Upper-level westerly anomalies shifted eastward to near the Date Line.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow


Easterly anomalies have persisted over the central and eastern Pacific associated with El Niño since mid-April (red box).

During June, these easterly anomalies were interrupted by robust atmospheric Kelvin wave/MJO activity.

During August, some westward propagation of westerly anomalies from the Maritime Continent to the Indian Ocean was evident.

From July to early October, generally stationary pattern was observed, consistent with an ongoing El Niño.

Recently, an eastward shift in the pattern is evident, with westerly anomalies strengthening near 160E.



Weekly Heat Content Evolution in the Equatorial Pacific


Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

During November, positive subsurface temperature anomalies increased and shifted eastward in association with the downwelling phase of a Kelvin wave. During November - January, the upwelling phase of a Kelvin wave shifted eastward.

Following a strong westerly wind burst in March, another downwelling phase of a Kelvin wave propagated eastward, reaching the South American coast during May.

Reinforcing downwelling events have followed, resulting in persistently above-normal heat content from the Date Line to 90W.

Recently, another downwelling phase of a Kelvin wave developed in the Pacific Ocean associated with anomalous low-level westerlies.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


The MJO index indicated a slight weakening of the signal but continued eastward propagation during the past week.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.


Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 08 Nov 2015
OLR




The GEFS MJO index-based OLR forecast depicts enhanced convection over the Indian Ocean, with the area of suppressed convection diminishing across the central Pacific Ocean.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:09-May-2015 to 08-Nov-2015
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (08 Nov 2015)




The constructed analog model depicts enhanced (suppressed) convection shifting eastward to the Maritime Continent (Americas/Africa).

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:09-May-2015 to 08-Nov-2015
The unfilled contours are CA forecast reconstructed anomaly for 15 days



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>