

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
18 January 2016

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

The MJO remained active during the past week and completed a global circulation since early December. The MJO signal recently weakened following the constructive interference with the ongoing El Nino during early January.

Dynamical model forecasts of the MJO index generally support a weaker signal during the next two weeks which may be related in part to increasing destructive interference with the ongoing El Nino.

The MJO is expected to play at least a minor role in anomalous convection throughout the global tropics with the ongoing El Nino remaining a major contributor.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Westerly anomalies shifted east across the Indian Ocean, while easterly anomalies persisted across the Maritime Continent.



A strong westerly wind burst continued over the Central Pacific Ocean at the Date Line.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

The red box highlights the persistent low-frequency westerly wind anomalies associated with ENSO.

An eastward shift in the pattern was observed in late October, related to subseasonal activity.

Renewed MJO activity during December produced an eastward propagation of westerly anomalies from the Indian Ocean

During early January, a strong westerly wind burst near the Date Line is related to constructive interference with the ongoing El Niño.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid to late December, the MJO signal destructively interfered with the ongoing El Niño with enhanced convection briefly affecting the Maritime Continent and northern Australia.

Suppressed convection returned to the Maritime Continent by the end of December as the MJO signal began to constructively interfere with the ongoing El Niño. Suppressed (enhanced) convection persisted across Brazil (Central Pacific).

During early to mid-January, the MJO signal continued to propagate east with enhanced convection diminishing at the Date Line. Suppressed convection expanded from the Maritime Continent and northern Australia to include the SPCZ.



Outgoing Longwave Radiation (OLR) Anomalies (5°N-5°S)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Since April, the ongoing El Niño is observed (red box) as a tendency toward a dipole of anomalous convection extending from the Maritime Continent (suppressed) to the East Pacific (enhanced).

During June and early July, the MJO became active, interfering with the ENSO signal at times. From August through October, the MJO remained weak, although some subseasonal activity did modulate the pattern of tropical convection during October.

During December, the MJO became active again, with the enhanced phase propagating from the Indian Ocean to the west-central Pacific during the month. Constructive interference between the MJO and ENSO signals resulted in widespread convection across the central Pacific.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


The ongoing ENSO state is highlighted by the red box, showing anomalous divergence over the central and eastern Pacific. This pattern has only been temporarily interrupted by strong Kelvin wave/MJO activity at times.

During June and early July, a high-amplitude MJO event was observed, constructively interfering with the El Niño signal in early July.


From July through early October, a generally stationary pattern, reflective of El Niño conditions, was observed. During late October, there was an eastward shift in the pattern associated with subseasonal activity.

Renewed MJO activity was observed during December and early January, yielding a robust signal in the upper levels. The MJO signal and the El Niño signal constructively interfered during early January.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A Wave-2 pattern is evident, likely due to influence from other modes of variability. Enhanced upper-level divergence (convergence) is inferred over the Pacific, Brazil, and the Indian Ocean (West Pacific and Atlantic).

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s⁻¹)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Upper-level westerly anomalies persisted over the South Asian and Maritime Continent.

Upper-level meridional divergence is apparent near and east of the Date Line, although slightly weakened from last week.



200-hPa Zonal Wind Anomalies (m s⁻¹)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Easterly anomalies have persisted over the central and eastern Pacific since June associated with El Niño (red box).

During late October, a temporary eastward shift in the westerly anomalies was evident across the Pacific.

Eastward propagation of upper-level zonal wind anomalies was apparent over the Maritime Continent and West Pacific during late December and early January, consistent with MJO activity.



Weekly Heat Content Evolution in the Equatorial Pacific


Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Following a strong westerly wind burst in March, a strong downwelling phase of a Kelvin wave propagated eastward, reaching the South American coast during May.

Reinforcing downwelling events have followed, resulting in persistently above-normal heat content from the Date Line to 80W throughout the period.

An expansion of below average heat content over the western Pacific is evident since spring and this area has increased since November 2015.

More recently, some warming is evident from near 170W and eastward.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


Although the MJO continued to propagate east during the past week, the amplitude of this index decreased.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts


The GFS ensemble MJO index forecast depicts a weaker signal during early part of Week-1 with a re-emerging signal by Week-2.

Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days




The GEFS MJO index-based OLR forecast depicts a weakening of anomalies early in Week-1 followed by a re-intensification of enhanced (suppressed) convection across the eastern Indian Ocean (West Pacific).

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:18-Jul-2015 to 17-Jan-2016
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (17 Jan 2016)




The constructed analog model depicts an eastward propagation of moderate strength OLR anomalies.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S–7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:18-Jul-2015 to 17-Jan-2016
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)




Precipitation Anomalies (Nov-Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>