

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
11 July 2016

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

The MJO remained weak during the last 7 days, with the RMM based index and CPC Index both indicating that the MJO is not active. Some velocity potential plots indicate a continued signal.

Forecast models suggest some strengthening, with the signal located anywhere from the Atlantic Ocean to the Indian Ocean, depending on the model. The signals in the various models are impacted by tropical cyclone activity.

Enhanced convection and tropical cyclone activity is expected over the eastern Pacific for the next 2 weeks, aligned with the MJO remnants and low-level signals. The upper-level signal is likely to propagate eastward and move across the Indian Ocean and Maritime Continent.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies persisted over much of the central Pacific while diminishing over the western Pacific.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

The red box highlights the persistent low-frequency westerly wind anomalies associated with the 2015-2016 El Niño background state.

Several fast-propagating intraseasonal events modulated the El Niño base state, and are marked using long (short) dashed lines for the enhanced (suppressed) phase of the intraseasonal signal.

During April, the wind field became less coherent as El Niño conditions weakened. During June, westerly anomalies generally prevailed across the Indian Ocean and Pacific, with the exception of a brief transition to easterlies beginning mid-month.

Recently, westerly anomalies prevailed over the central Pacific. Anomalies were relatively weak overall.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Enhanced convection returned to the eastern Indian Ocean and western Maritime Continent during mid June, while suppressed convection persisted over the western equatorial Pacific.

In late June, enhanced convection remained across the eastern Indian Ocean and pushed into the Maritime Continent. Convection remained suppressed further east to the DL, but to less so relative to earlier in June.

During late June and early July, enhanced convection moved northward over South Asia and spread eastward to the eastern Pacific, while suppressed convection moved in over the Maritime Continent and the western Atlantic.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)


The 2015-2016 El Niño background state is observed (red box) as a dipole of anomalous convection extending from the Maritime Continent to the East Pacific. The signal weakened steadily through boreal Spring.

Alternating periods of constructive and destructive interference with El Niño are evident. A fast eastward propagating signal raced across the Pacific during February.

During early May, an eastward-propagating convective envelope associated with the MJO developed east of the Prime Meridian. This OLR signal weakened over the Pacific.

During mid-June, some enhanced convection was evident over Maritime Continent.

Recently, signals have been mixed, with tropical cyclone activity evident in the patterns.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

The 2015-16 El Niño background state is highlighted by the red box, showing anomalous divergence over the central and eastern Pacific.


MJO activity was evident in February and March, alternatively constructively and destructively interfering with the ENSO background state.

The upper-level velocity potential pattern became less coherent as the El Niño waned during April.


During May and June, an eastward propagating signal was evident in the upper-level velocity potential field. This signal was more coherent in time and space than the low-level MJO indicators.

Anomalous divergence is evident over the Americas and Africa, with anomalous convergence over the Maritime Continent.

200-hPa Velocity Potential Anomaly: 5N-5S



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The large scale upper-level velocity potential anomaly pattern exhibits Wave-3 structure, not consistent with robust MJO activity.


Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies expanded over the central and western Pacific.

200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Easterly anomalies have persisted over the central and eastern Pacific from June 2015 to May 2016 associated with El Niño (red box). Corresponding westerly anomalies persisted over the Maritime Continent.

During May, westerly anomalies expanded eastward to the Date Line as El Niño weakened. Faster propagating modes were evident in the upper-level wind field.

The upper-level wind field became less coherent during late May and early June.

Most recently easterly anomalies are prevalent from the Maritime Continent to the eastern Pacific, with westerly anomalies over the Americas.



Weekly Heat Content Evolution in the Equatorial Pacific


Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Reinforcing downwelling events were observed during the second half of 2015, resulting in persistently above-normal heat content from the DL to 80W throughout the period.

An eastward expansion of below average heat content over the western Pacific is evident since January, with negative anomalies beginning to spread east of the Date Line.

In the last three months, there has been a rapid eastward expansion of below-average oceanic heat content across the central and eastern Pacific. Negative anomalies now extend across the equatorial Pacific.

A small area of positive SST anomalies is evident near 135E.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past week, the RMM index indicated some strengthening of the MJO signal over the Africa.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts


During the next two weeks, the GFS ensemble indicates a complete breakdown of any remaining MJO signal, with westward propagation. This is likely related to tropical cyclone activity.

Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast


Spatial map of OLR anomalies for the next 15 days



The GEFS OLR forecast based on the RMM Index depicts a stagnant signal with amplification through 10 days, then a weakening signal.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (10 Jul 2016)




The Constructed Analog (CA) model predicts a stationary and weak signal through Week-2.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:09-Jan-2016 to 10-Jul-2016
The unfilled contours are CA forecast reconstructed anomaly for 15 days



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>