

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
26 December 2016

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- The CPC velocity potential-based MJO index indicates a more coherent MJO recently, but the RMM index continues to depict a MJO signal of low amplitude since the end of November.
- Background La Niña conditions continue to influence the pattern of anomalous tropical convection.
- Dynamical model forecasts of the RMM index generally maintain a weak MJO signal during the next two weeks. Therefore, its contribution if any to the extratropical circulation is unclear.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Anomalous cyclonic flow near Hawaii has promoted a surge of tropical Pacific air into the western U.S.

Anomalous westerlies (easterlies) persisted over the eastern Indian Ocean and Maritime continent (equatorial central Pacific Ocean), consistent with the low-frequency state.



850-hPa Zonal Wind Anomalies (m s⁻¹)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

In July and early August, high frequency, eastward-propagating modes were observed crossing the Pacific.

During September and October, persistent westerly (easterly) anomalies were evident over the eastern Indian Ocean and western Maritime Continent (central Pacific). These anomalies are low frequency in nature, and reflect the La Niña base state as well as a negative phase of the Indian Ocean Dipole.

During the past month, westerly anomalies continued to persist across the Indian Ocean (red box).



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

The low-frequency signal continued to influence the pattern of anomalous convection during late November, with subseasonally-linked drying observed in the western Indian Ocean yielding a tripole of anomalous convective activity for the Indo-Pacific region.

During early December, the aforementioned tripole persisted across the Indo-Pacific region. Elsewhere in the Pacific, anomalous convection shifted east to Hawaii.

During mid-December, the low-frequency state continued to influence anomalous convection across the Indo-Pacific region, while convection increased along the SPCZ. Negative OLR anomalies extended from Hawaii to the West Coast of the United States.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°S - 17.5° S)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection shifted slowly east from the eastern Indian Ocean to the Maritime Continent has been evident since July (green box). Low-frequency suppressed convection, tied to building La Niña conditions, has been apparent near the Date Line since August (black box).

A fast eastward propagating convective envelope was evident during early September.

From late November into early December, an eastward propagating mode (likely a Kelvin wave) was apparent across the Indian Ocean and West-Central Pacific.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


Through early August, an eastward propagating signal was evident, with multiple periods of variability apparent.

During August, the intraseasonal signal became less coherent, with a weaker and somewhat more stationary anomaly field in place. By late August and early September, there was renewed propagation of the intraseasonal signal.


From mid-September to late October, the low frequency signal dominated the pattern. During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum.

During December, a more stationary pattern is evident with positive (negative) anomalies observed across the Indian Ocean (Maritime Continent and West Pacific).

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The pattern of upper-level velocity potential anomalies has become more Wave-1 spatially with upper-level convergence strengthening over the Indian Ocean and upper-level divergence over the Western Hemisphere.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation


200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Enhanced upper-level westerlies extended from the tropics of the Pacific Ocean to the mid-latitudes of western North America.



Anomalous upper-level convergence (divergence) persisted over the Indian Ocean (Maritime Continent).

200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

During July, some eastward propagation was evident, although the spatial consistency implies higher frequency variability than expected with the MJO.

During September, eastward propagation of westerly anomalies was broadly consistent with organized MJO activity.

During November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident.

Since late November, easterly anomalies have re-emerged across the Indian Ocean consistent with the passage of subseasonal activity and the re-alignment of the low frequency base state (red box).




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Downwelling events were observed through early February, resulting in persistently above-normal heat content from the Date Line to 80W over that period.

An eastward expansion of below average heat content over the western Pacific is evident through April, with widespread negative anomalies building across the Pacific over the course of the spring and summer months.

The strongest negative anomalies now persist in the East Pacific as the oceanic heat content anomalies have become slightly positive at the Date Line.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


The MJO index remained low in amplitude since the end of November.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GFS ensemble forecast depicts a continued incoherent MJO signal during the next two weeks.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 25 Dec 2016
OLR




The prediction for OLR anomalies over the next two weeks based on the GEFS RMM forecast indicates enhanced convection developing over the Indian Ocean by Week-2.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cint:4Wm⁻²) Period:25-Jun-2016 to 25-Dec-2016
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (25 Dec 2016)




The prediction for OLR anomalies based on the constructed analog RMM forecast indicates eastward propagation of suppressed convection from the Indian Ocean to the Maritime Continent.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S- 7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5° S, 7.5° N] (cont:4 Wm^{-2}) Period:25-Jun-2016 to 25-Dec-2016
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>