

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
6 February 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- The MJO is active, with both the RMM based and CPC velocity potential based MJO indices depicting robust eastward propagation of an enhanced (suppressed) convective envelope over the West Pacific (Africa and the western Indian Ocean).
- The MJO is transitioning from constructively interfering with the low frequency state which favors convection over the Maritime Continent, to destructively interfering with the background of suppressed Central Pacific convection.
- Dynamical model RMM index forecasts indicate continued propagation, with strengthening as the signal moves across the Pacific and into the Western Hemisphere, before possible weakening late in Week-2.
- The MJO is anticipated to influence the evolution of the global tropical convective pattern. Impacts are less certain as the signal looks to destructively interfere with the base state in the Central Pacific.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Westerly anomalies shifted eastward across the Maritime Continent.



Easterly anomalies extended further eastward across the Central Pacific in the past 5 days.

850-hPa Zonal Wind Anomalies (m s⁻¹)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Since late August, persistent westerly (easterly) anomalies were evident over the eastern Indian Ocean and western Maritime Continent (central and western Pacific) as shown by the red (blue) box at right. These anomalies are low frequency in nature, associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During mid-January, Rossby Wave activity was evident, with destructive interference on the base state evident through 100E.

Over approximately the last week the low frequency state has re-emerged as the dominant signal, with easterlies strengthening across the Central Pacific.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During early/mid-January, the low frequency state with enhanced (suppressed) convection over the Maritime Continent (Indian Ocean and Central Pacific) dominated. Kelvin wave activity was apparent across the southern Indian Ocean.

In mid-/late January, enhanced convection associated with the subseasonal state shifted over the Atlantic and Africa, while the low frequency background state persisted.

The low frequency pattern continued in late January/early February as the subseasonal signal constructively interfered. A break in convective suppression northeast of Madagascar was associated with what would become Tropical Storm Carlos.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°S - 17.5° S)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern Indian Ocean and the Maritime Continent has been evident since July (green box), with suppressed convection over the Indian Ocean and near the antimeridian (black boxes).

A fast eastward propagating convective envelope was evident during early September. Another intraseasonal event occurred during late November and early December. Each briefly interfered with the background state.

Since late January, another eastward propagating event has interacted with the low frequency background as it shifted from the Indian Ocean to the Maritime Continent.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


In late August and early September, intraseasonal activity was apparent, before reversion to the low frequency pattern associated with the negative IOD and La Niña through late October.

During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum.


After a break in intraseasonal activity during early December, a signal emerged over the Maritime Continent and has continued propagating through the present.

The aforementioned intraseasonal signal constructively interfered with the low frequency state over the Maritime Continent near the beginning of February.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



Upper-level velocity potential anomalies reveal a robust wavenumber-1 pattern, with enhanced (suppressed) convection over the Maritime Continent through Central Pacific (Americas, Atlantic, Africa, and Indian Ocean). Convection generally aligns well with circulation expectations.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation


200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Anomalous divergence is evident over the Maritime Continent, with anomalous westerlies strengthening north of the equator in the Central Pacific.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow


After a quiet August, eastward propagation of westerly anomalies was broadly consistent with organized MJO activity during September.

In November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident.

In late November, easterly anomalies re-emerged across the Indian Ocean and Maritime Continent, consistent with the passage of sub-seasonal activity and the re-alignment of the low frequency base state.

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere.

More recently, the low frequency state has re-emerged as destructive interference with the intraseasonal signal waned.




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June, with widespread negative anomalies building across the Pacific over the course of boreal spring and summer.

More recently, upper-ocean heat content anomalies have been low amplitude, consistent with the forecast transition to ENSO-neutral conditions.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past week, continued eastward propagation of an MJO signal was evident over the Indian Ocean and Maritime Continent. As of yesterday, the index has entered the West Pacific.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast

RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days


light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS depicts strengthening and continued eastward propagation of the intraseasonal signal over the Western Pacific and into the Western Hemisphere over the next two weeks.


The signal is forecast to weaken by both the ensemble mean and individual members over the Western Hemisphere late in Week-2.

Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast


Spatial map of OLR anomalies for the next 15 days



The GEFS prediction for RMM Index-based OLR anomalies over the next two weeks depicts eastward propagation of robust OLR anomaly envelopes.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (05 Feb 2017)




The statistical (Constructed Analog) RMM-based OLR anomaly prediction is very similar spatially to that of the GEFS, with weaker amplitude initially, but intensifying during Week-2.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:06-Aug-2016 to 05-Feb-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>