

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
15 May 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- Both RMM-based and CPC velocity potential-based MJO indices indicate an continued signal with some eastward propagation.
- Dynamical model RMM index forecasts generally indicate some eastward propagation, through the next week. Some models significantly weaken the signal. All of the models imply enhanced convection over the Indian Ocean and Maritime Continent, regardless of predicted amplitude of the MJO.
- The continued propagation of the MJO across the Indian Ocean is favored in this week's outlook, with the MJO playing a role in the overall pattern of tropical convection. Other modes of variability are also likely to influence the pattern over the Maritime Continent and Indian Ocean.
- During Week-1, the MJO may play a role in a tropical cyclone over the Bay of Bengal. During Week-2, the convective signal from the MJO is likely to be centered over the Maritime Continent.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies moved northward to over Indian and Southeast Asia, while westerly anomalies built in, along the equator, over the Maritime Continent

Westerly anomalies consolidated and strengthened over the East Pacific.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies, shown by the red (blue) box at right, were associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During late January, Rossby wave activity was evident, with destructive interference on the base state evident through 100E.

During February, MJO activity also destructively interfered with the base state. During mid-March and early April, the low frequency state seemed to reemerge, with some intraseasonal variability evident in late March.

Recently, the anomaly pattern seems similar to the prior, low-frequency pattern, though over the Indian Ocean, the signal is opposite. The pattern shows some westward moving features near the Maritime Continent.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid-April, suppressed convection persisted over much of the East Pacific and Indian Oceans. Enhanced convection was evident over the western Pacific.

During late April, suppressed convection persisted over much of the Indian Ocean. Over the Pacific north (south) of the equator suppressed (enhanced) convection developed in long, west-east oriented patterns.

During early May, suppressed convection remained over much of the Indian Ocean. Enhanced convection developed over the Americas and Atlantic Ocean. Tropical cyclone activity impacted the pattern over the South and East Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S - 7.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern Indian Ocean and the Maritime Continent has been evident from July through mid-February (green box), with suppressed convection over the Indian Ocean and near the Date Line (black boxes).

Intraseasonal events in January through mid-March have served to alter the low frequency states. Particularly, with the suppressed phase reversing the low frequency enhanced convective signal over the Maritime Continent in late February. The MJO signal weakened by mid-March.

During April, patterns were generally weak. By early May, enhanced convection generally spread from the Americas to Africa, and over the Maritime Continent.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum. The pattern, during December and January, was more related to seasonal variability.

A signal emerged over the Maritime Continent and continued propagating through early March, creating alternating periods of constructive and destructive interference with the base state.


During March, a low frequency signal favoring enhanced (suppressed) convection over the Maritime Continent (Indian Ocean) once again became the primary component of the anomaly field. Kelvin wave activity has been apparent during April, primarily east of the Date Line.

The eastward propagation with Kelvin waves continued into May, with some consolidation evident over the East Pacific. Some of the low-frequency signal is interfering.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The spatial distribution of the upper-level VP anomaly field depicts a large area of enhanced ascent, north of the equator, from the Americas to the Maritime Continent. This is likely the result of the interference of intraseasonal and low-frequency signals.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s⁻¹)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

A wave train is evident over the North Pacific during early May, with mid-latitude influence near the Date Line.

During the second week of May, generally westerly (easterly) anomalies spread over Africa and the Indian Ocea (Pacific Ocean and Americas).



200-hPa Zonal Wind Anomalies (m s⁻¹)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

In November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident. In late November, easterly anomalies re-emerged across the Indian Ocean and Maritime Continent, consistent with the passage of sub-seasonal activity and the re-alignment of the low frequency base state.

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere.

Easterly anomalies returned to the East Pacific during late April. During early May, easterly (westerly) anomalies also developed over the Central Pacific (Maritime Continent). Some westward moving features are evident, as a result, the anomalies over the Central Pacific shifted westward.




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June, with widespread negative anomalies building across the Pacific over the course of boreal spring and summer.

The anomaly field has weakened across the central and eastern Pacific, with positive anomalies persisting over much of the Pacific, except for a small portion of the eastern Pacific.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past several days, the amplitude of the RMM-based MJO index has decreased, though still maintaining an eastward propagation.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS depicts a gradual decrease in the amplitude of the MJO index over the next week, with the potential emergence of a signal late in Week-2.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 14 May 2017
OLR




The GEFS RMM-based OLR anomaly forecast depicts a gradual weakening of a fairly stationary anomaly field over the next week, with a slight strengthening near the end of the period.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:12-Nov-2016 to 14-May-2017
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (14 May 2017)




The statistical RMM-based OLR anomaly prediction indicates more robust eastward propagation of the anomaly pattern, with the enhanced phase of the MJO emerging over the West Pacific by the end of the period.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:12-Nov-2016 to 14-May-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>