

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
17 July 2017

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO showed signs of organization over the past week, with the enhanced phase approaching the Maritime Continent.
- The various dynamical models continue to exhibit fairly poor agreement, though there has been some trend toward the ECMWF solution that supports more coherent MJO activity. Likewise, statistical guidance now shows more canonical MJO evolution.
- The MJO is expected to propagate eastward across the Maritime Continent over the next week, with typical warm-season precipitation impacts expected from the Indian Ocean to the West Pacific, including parts of Southeast Asia.
- The forecast MJO evolution would support reduced tropical cyclone (TC) activity over the East Pacific basin. This is somewhat at odds with dynamical guidance which shows recurring tropical cyclogenesis over the next two weeks.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies



Westerly anomalies persisted over the equatorial Indian Ocean during both periods.

Generally low-amplitude anomalies were observed over the Pacific during the recent period.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies, shown by the red (blue) box at right, were associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During late January, Rossby wave activity was evident, with destructive interference on the base state evident through 100E.

In February, MJO activity also destructively interfered with the base state. During mid-March and early April, the low frequency state reemerged, with some intraseasonal variability evident in late March.

Equatorial flow was fairly weak throughout June, with easterlies favored between 120E and the Date Line and also emerging across the western Indian Ocean. During July, there appeared to be some organization of the pattern around a wave-1 structure, but with westward-moving variability still evident.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid- to late June, suppressed convection was observed over much of the Indian and Pacific Ocean basins. A surge of tropical moisture into southeastern North America is also evident.

Convective signals were fairly weak in late June and early July, though enhanced convection was observed over parts of Southeast Asia.

By early to mid-July, the enhanced convective pattern became more organization over Southeast Asia and the Maritime Continent. Some low-frequency tendency is noted over the latter during the past month.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern IO and the Maritime Continent was evident from July 2016 through early April 2017 (green box), with suppressed convection near the Date Line (right black box). The remainder of the IO generally had suppressed convection during this period (left black box), with the exception of an MJO-related wet period from mid-Feb to early March.

From mid-April through present, convective anomalies were generally weak. In mid-May, enhanced convection was noted over the Indian Ocean with some eastward propagation.

During the past week convective signals increased in amplitude over the Maritime Continent.





# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

A signal emerged over the Maritime Continent and continued propagating through early March, creating alternating periods of constructive and destructive interference with the base state.

During March, a low frequency signal favoring enhanced (suppressed) convection over the Maritime Continent (Indian Ocean) once again became the primary component of the anomaly field.


Kelvin wave activity was apparent from April through early June, as seen in the rapidly propagating eastward signals. Over the past month anomalies have been somewhat stationary with enhanced (suppressed) convection over the Maritime Continent (East Pacific). The recent maximum observed near 100E is among the strong signals observed over that region during the past six months.

200-hPa Velocity Potential Anomaly: 5N-5S  
5-day Running Mean





# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The latest pattern of upper-level velocity potential anomalies reveals a somewhat coherent pattern, with enhanced (suppressed) convection centered over Southeast Asia, the Maritime Continent, and the far West Pacific (Africa and parts of the Western Hemisphere, with the exception of tropical wave activity).

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation


# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Upper-level wind anomalies continue to be weak throughout the tropics, though a somewhat coherent pattern has been observed recently with easterlies (westerlies) over the Indian Ocean (Maritime Continent and West Pacific).



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Easterly anomalies returned to the East Pacific during late April and persisted with some period of high-frequency interference.

During early to mid-June, easterly anomalies were most prominent across the global tropics, in part due to mid-latitude influences.

Recently, westerly anomalies were observed over the West Pacific, consistent with the organized enhanced convection ongoing west and northwest of that region.




# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June 2016, with negative upper-ocean heat content anomalies persisting through the end of 2016.

During the current year, positive anomalies have developed and generally persist over the entire basin. The anomalies are generally weak, though a small pocket of warmer than normal heat content has developed between 150-140W. Much of this heat is in the uppermost 100 meters, with anomalously cool water below this.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past week, the RMM index showed increasing amplitude into Phase 3, with some subtle eastward propagation.



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.





# GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS forecast has trended somewhat toward the ECMWF solution, and now depicts some weak but coherent MJO propagation during the next two weeks.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 15 Jul 2017  
OLR




The GEFS RMM-based OLR anomaly forecasts the canonical MJO evolution at very low amplitude.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of ( $7.5^{\circ}$  S- $7.5^{\circ}$  N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [ $7.5^{\circ}$ S, $7.5^{\circ}$ N] (cont:4Wm<sup>-2</sup>) Period:13-Jan-2017 to 15-Jul-2017  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (15 Jul 2017)




The constructed analog RMM-based OLR anomaly is fairly similar to the GEFS today, but with a slightly faster eastward propagation.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:13-Jan-2017 to 15-Jul-2017  
The unfilled contours are CA forecast reconstructed anomaly for 15 days




# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (May - Sep)



Precipitation Anomalies (May - Sep)






# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>