

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
6 November 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- The CPC velocity potential-based and RMM-based MJO indices indicate that the MJO has weakened since late October.
- Dynamical model RMM-index forecasts differ, but the preferred model solution indicates a continued weakening of the MJO.
- The MJO is expected to have less of an impact on anomalous convection throughout the global tropics compared to mid and late October. The base state (potentially emerging La Niña conditions) and a robust equatorial Rossby wave are expected to play the largest roles.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly (easterly) anomalies developed over the northern Indian Ocean (Southeast Asia and West Pacific) Ocean during the past five days.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Low-frequency easterly anomalies (blue box) have largely persisted over the west-central Pacific throughout the last 180 days.

Equatorial zonal wind anomalies were of low amplitude in June. During July, a slight eastward shift in the low-frequency pattern is noted, related to short-lived MJO activity.

During August and September, the low-frequency envelope of easterly anomalies became re-established from 140E to just east of the Date Line.

During October, a robust MJO event became established, with eastward propagation of westerly anomalies over the Maritime Continent and Central Pacific. The MJO destructively interfered with the low-frequency state and resulted in small westerly anomalies by the end of October.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During early to mid-October, enhanced convection associated with a strengthening MJO developed over the eastern Indian Ocean and western Maritime Continent. Suppressed convection remained entrenched over the central and eastern Pacific.

Enhanced convection, related to a robust MJO shifted east across the Maritime Continent and West Pacific during mid to late October, while suppressed convection developed over Africa and the Indian Ocean.

During late October, the MJO rapidly propagated east across the Western Hemisphere with convection increasing once again across Africa. The low frequency state continued to support suppressed convection across the equatorial central Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S - 7.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid-July, there was a burst of enhanced convection over the Maritime Continent, due to interactions between a short-lived intraseasonal signal and the low-frequency state.

Multiple modes of variability including tropical cyclones contributed to the pattern of anomalous convection during August and September. Weak MJO activity was present during August and early September.

An MJO signal emerged over the Maritime Continent during early October and propagated east rapidly during the past month. Since the beginning of November, the enhanced phase of the MJO shifted east of the Prime Meridian.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

During July, enhanced convection strengthened over the Maritime Continent as the low-frequency signal constructively interfered with an easterly propagating signal. This eastward propagating signal appears more or less intact with a period in line with canonical MJO phase speeds.


A signal on the MJO timescale is evident in this field during late August and September.

Another MJO event developed near the Maritime Continent during early October, with a large upper-level footprint near 120E and robust eastward propagation. The leading edge of the enhanced phase shifted over Africa and into the Indian Ocean at the beginning of November.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A wave-1, upper-level velocity potential pattern reflects canonical MJO activity, with the enhanced (suppressed) phase centered over the western Indian Ocean (Pacific Ocean).

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly


Blue shades: Easterly anomalies

Red shades: Westerly anomalies

The upper-level zonal wind pattern also reflects a robust Wave-1 asymmetry characteristic of a strong MJO event.

The envelope of westerly anomalies shifted east across the Indian Ocean during the past five days.

Midlatitude influences from both hemispheres are evident near the Date Line.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

During early to mid-June, easterly anomalies were most prominent across the global tropics, in part due to mid-latitude influences.

Starting in July, the anomaly patterns propagated eastward associated with weak MJO activity and atmospheric Kelvin waves.

During September, fast-moving eastward propagation of anomalies continued, consistent with additional atmospheric Kelvin Waves. A slower signal was evident over the eastern Maritime Continent and west Pacific before decaying near 160 W.


Westerly (easterly) anomalies expanded across much of the Pacific (Indian Ocean) during early October, with an eastward propagation of those two components through later October.



Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Negative upper-ocean heat content anomalies have slowly increased across the equatorial Pacific east of the Date Line.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


The amplitude of the RMM-index decreased during the final week of October as the MJO propagated towards the Indian Ocean.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The bias-corrected GEFS forecast indicates the MJO continuing its fast eastward propagation across the Indian Ocean to the Maritime Continent during the next week.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 05 Nov 2017
OLR




OLR anomalies based on the GEFS RMM-index forecast depict enhanced convection shifting east from the Indian Ocean to the West Pacific during the next two weeks.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:06-May-2017 to 05-Nov-2017
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (05 Nov 2017)




The constructed analog RMM-index forecast features a rapid decay of anomalous by late in Week-1.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:06-May-2017 to 05-Nov-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>