

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
13 November 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- The CPC velocity potential-based and RMM-based MJO indices indicate that the MJO has weakened during early November. The remaining signal is now over the Eastern Indian Ocean/Maritime Continent.
- Dynamical model RMM-index forecasts differ, but the preferred model solution indicates a continued weakening of the MJO.
- The MJO is expected to have less of an impact on anomalous convection throughout the global tropics compared to mid and late October. The base state (La Niña conditions) and higher frequency variability are likely to play larger roles.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Westerly anomalies continued over the eastern Indian Ocean but weakened over the western Indian Ocean.



Easterly anomalies strengthened and expanded in coverage near the Date Line.

Westerly anomalies developed over the eastern Pacific near 120W.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Low-frequency easterly anomalies (blue box) have largely persisted over the west-central Pacific throughout the last 180 days.

Equatorial zonal wind anomalies were of low amplitude in June. During July, a slight eastward shift in the low-frequency pattern is noted, related to short-lived MJO activity.

During August and September, the low-frequency envelope of easterly anomalies re-established from 140E to just east of the Date Line. During October, a robust MJO event developed, with eastward propagation of westerly anomalies.

Recently, MJO activity is beginning to constructively interfere with the low-frequency state.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During early to mid-October, enhanced convection associated with a strengthening MJO developed over the eastern Indian Ocean and western Maritime Continent. Suppressed convection remained entrenched over the central and eastern Pacific.

During late October, the MJO rapidly propagated east across the Western Hemisphere with convection increasing once again across Africa. The low frequency state continued to support suppressed convection across the equatorial central Pacific.

During early November, the remaining MJO signal moved across the Indian Ocean, enhancing convection in the area, while low-frequency signals remained over the Central Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S - 7.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid-July, there was a burst of enhanced convection over the Maritime Continent, due to interactions between a short-lived intraseasonal signal and the low-frequency state.

Multiple modes of variability including tropical cyclones contributed to the pattern of anomalous convection during August and September. Weak MJO activity was present during August and early September. The low-frequency signal emerged more fully in August.

An MJO signal emerged over the Maritime Continent during early October and propagated eastward rapidly during the past month. During early November, the enhanced phase of the MJO shifted across the Indian Ocean, and now more closely aligns with the low-frequency state.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

During July, enhanced convection strengthened over the Maritime Continent as the low-frequency signal constructively interfered with an easterly propagating signal. This eastward propagating signal appears more or less intact with a period in line with canonical MJO phase speeds.


A signal on the MJO timescale is evident in this field during late August and September.

Another MJO event developed near the Maritime Continent during early October, with a large upper-level footprint near 120E and robust eastward propagation. That MJO signal has continued into early November, now centered over the Maritime Continent.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



Generally the upper-level velocity potential pattern reflects a wave-1 structure, though there is some noise from non-MJO modes of variability. The enhanced (suppressed) phase centered over the Maritime Continent (Atlantic).

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly


Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Westerly anomalies over the Maritime Continent were replaced with easterly anomalies.

Westerly anomalies over the eastern Pacific strengthened.

Mid-latitude influences from both hemispheres are evident over the eastern Pacific.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

During early to mid-June, easterly anomalies were most prominent across the global tropics, in part due to mid-latitude influences.

Starting in July, the anomaly patterns propagated eastward associated with weak MJO activity and atmospheric Kelvin waves.

During September, fast-moving eastward propagation of anomalies continued, consistent with additional atmospheric Kelvin Waves. A slower signal was evident over the eastern Maritime Continent and west Pacific.


Westerly (easterly) anomalies expanded across much of the Pacific (Indian Ocean) during later October, and have remained.



Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Negative upper-ocean heat content anomalies remain entrenched in the eastern Pacific, with some strengthening near 100W.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The bias-corrected GEFS forecast indicates a weakening MJO signal significantly.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 12 Nov 2017
OLR




OLR anomalies based on the GEFS RMM-index forecast reflect a weakening MJO signal.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cint:4Wm⁻²) Period:13-May-2017 to 12-Nov-2017
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (12 Nov 2017)




The constructed analog RMM-index forecast depicts a much stronger than signal than the dynamical models, with the enhanced phase moving into the West Pacific.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:13-May-2017 to 12-Nov-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>