

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by: Kyle MacRitchie  
Climate Prediction Center / NCEP  
12 March 2018

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO continued to weaken as it propagated over the Indian Ocean during the past week. The GEFS forecasts that the MJO will continue to weaken over the next couple of weeks.
- Both the GEFS and the constructed analog forecasts suggest that an active MJO could emerge near the end of March over the Maritime Continent.
- Areas of enhanced convection over the Indian Ocean and Maritime Continent, which are partially related to the weak MJO signal, may lead to tropical cyclone development over the next week.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Weak anomalies throughout the Indian Ocean are consistent with low tropical wave activity.

Anomalous easterlies are present along the equator, consistent with La Niña.

Midlatitude waves dominate the Atlantic flow.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Low frequency anomalous easterlies that have been present since before early September reversed sign during February.

During October and early November, a robust MJO event developed, with eastward propagation of westerly and easterly anomalies. This event weakened in early to mid-November.

A new MJO event became organized in December, propagating from the Indian Ocean to the Pacific. The signal crossed the Western Hemisphere in late December, re-emerging over the Indian Ocean in early January. The signal continued to propagate eastward, moving into the central and eastern Pacific and began weakening over the past couple of weeks.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

The suppressed phase of the MJO led to positive OLR anomalies over much of the Maritime Continent and southern Indian Ocean during early February. The enhanced phase destructively interfered with ENSO over the central Pacific and enhanced South Pacific convection.

Positive OLR anomalies weakened but continued to dominate the Indian Ocean and Maritime Continent during mid-February.

Positive OLR anomalies over the southern Indian Ocean and Maritime Continent weakened during the past week. During the same period, positive OLR anomalies strengthened over the central Pacific associated with anomalously low cloud cover and convection.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°S - 17.5° S)


**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

An active MJO formed in early October and circumnavigated the globe by early November.

Another MJO event developed in late November over the eastern Indian Ocean and Maritime Continent that was able to briefly disrupt the La Niña convective suppression near the Date Line. It re-emerged in the Indian Ocean at the end of December and strengthened as it shifted east towards the Date Line at present.

A third MJO event developed during January, leading to a reversal of the La Niña convective signal over the Date Line in early February. This MJO signal has weakened substantially over the past couple of weeks.





# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

An MJO event developed near the Maritime Continent during early October with strong anomalous upper-level winds near 120E. The signal circumnavigated the global tropics and weakened about 30 days later.

MJO activity renewed in November. The intraseasonal signal associated with this MJO event was weaker than the previous episode due to destructive interference from an equatorial Rossby wave.


The signal destructively interfered with the base state through the end of December, crossing the Western Hemisphere into the Indian Ocean for the beginning of January. Since then, it has continued eastward and strengthened.

This MJO event further intensified during January and early February, but has since weakened substantially.





# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A wave-1 pattern is present with enhanced (suppressed) convection generally over the Eastern (Western) Hemisphere. The areas of enhanced convection over the Indian Ocean and Maritime Continent are potential areas of tropical cyclone formation over the next week.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation  
Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )


Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies

Cross-equator flow interactions continued over the eastern Pacific during the last 10 days, along with a highly amplified pattern across the midlatitudes in the Western Hemisphere.

Upper-level wind anomalies were generally weak across the Eastern Hemisphere, with a weak divergent signal over eastern Africa and the western Indian Ocean apparent in the most recent analysis.



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Low-frequency westerly anomalies remained in place east of 140°E starting in October, with the exception of a brief period of easterlies in late October. There is also some recent evidence of easterlies over the far Eastern Hemisphere over the last week or so that appear to have extratropical sourcing.

In mid-December anomalous easterlies developed east of the Date Line, replacing the westerly anomalies that had been generally present since October.

Strong anomalous westerlies that formed in early January just west of the Date Line propagated eastward, consistent with a strong MJO event during this period.

More recently, relatively stationary anomalous westerlies have dominated most of the globe east of 100°E.




# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Negative upper-ocean heat content anomalies have persisted in the central and eastern Pacific since late summer.

A downwelling Kelvin wave associated with the intraseasonal signal weakened the negative anomalies across the east-central Pacific during late January and early February.

Another downwelling Kelvin wave is leading to highly positive, eastward-propagating anomalies near the Date Line.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


A weak MJO continued to weaken as it propagated across the Indian Ocean over the past week.



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.





# GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS forecasts weakening of the MJO signal over the next few days.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 11 Mar 2018  
OLR




OLR anomalies associated with the MJO based on the GEFS show little propagation with a weakening signal over time.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [7.5°S,7.5°N] (cint:4Wm<sup>2</sup>) Period:09-Sep-2017 to 11-Mar-2018  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model  
reconstruction by RMM1 & RMM2 (11 Mar 2018)




The constructed analog is more progressive with the MJO signal, bringing a slightly weakened enhanced envelope to the West Pacific during the next two weeks.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of ( $7.5^{\circ}$  S- $7.5^{\circ}$  N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [ $7.5^{\circ}$ S,  $7.5^{\circ}$ N] ( $\text{cont: } 4 \text{ Wm}^{-2}$ ) Period: 09-Sep-2017 to 11-Mar-2018  
The unfilled contours are CA forecast reconstructed anomaly for 15 days




# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov - Mar)



Precipitation Anomalies (Nov - Mar)






# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>