

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by: Christina Finan  
Climate Prediction Center / NCEP  
8 October 2018

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO remained active during the past week, with the signal maintaining amplitude and continuing eastward propagation into Phase 1.
- Dynamical model guidance anticipates the MJO transitioning to Phase 2 in Week-1, with the enhanced signal spanning the Indian Ocean and suppressed signal continuing over the Maritime Continent and western Pacific.
- The enhanced signal over the Indian Ocean is likely to contribute to increased chances of tropical cyclogenesis for the region. The Eastern Pacific is not forecast to be active as a move to Phase 2 will bring suppressed convection to the region.
- Also of note, the robust westerly wind burst in the Pacific may help push conditions towards El Niño as we head towards boreal Winter.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)


Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies

Low-level anomalous westerlies persisted across much of the tropical Pacific during late September.

The anomalous westerlies have strengthened over the eastern and central Pacific and shifted out of the western Pacific.



# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

The MJO was active during late April and May. Westward moving variability weakened the signal in June. A weak intraseasonal signal re-emerged during mid to late July. During August, the intraseasonal signal weakened, and other modes, including Rossby wave and tropical cyclone activity, influenced the pattern.

Through much of September, Rossby wave activity continued to dominate the Pacific, while westerly anomalies overspread the equatorial Pacific.

During late September, signs of a westerly windburst near 160°E are evident. Westerly anomalies have continued to strengthen over the central and eastern Pacific. This may foreshadow a transition towards El Niño conditions and an increase in the warm water volume availability in the Central Pacific.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

In early September, suppressed convection became more prominent over the Indian Ocean. An uptick in easterly waves resulted in three tropical cyclones forming over the Atlantic.

Easterly wave activity waned by mid-September, with suppressed convection across much of the tropical Atlantic as wind shear increased. Suppressed convection over the Indian Ocean shifted eastward toward the Maritime Continent, as small regions of enhanced convection are seen over the western Indian Ocean.

Most recently, the suppressed Indian Ocean convection continued to shift further east and strengthen over the Maritime Continent, while the enhanced convection emerging in the Western Indian Ocean becomes more widespread as the active phase of the MJO continued through Phase 1.




# Outgoing Longwave Radiation (OLR) Anomalies (5°S - 5°N)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

The MJO was active during the boreal spring, remaining fairly strong throughout April, weakening briefly, and re-strengthening in mid-May. The signal weakened again in June. During July, the MJO signal re-emerged, with some eastward propagation evident in the OLR field.

Kelvin waves, Rossby waves, and tropical cyclones dominated the pattern during August and early September, while the intraseasonal signal remained fairly weak. During mid-September, the suppressed phase of the MJO emerged over the Eastern Indian Ocean and Maritime Continent. Since then, the suppressed signal has propagated further east and enhanced convection has emerged over the western Indian Ocean.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)


Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

There was robust MJO activity through boreal spring along with the decay of La Niña conditions. The enhanced phase of the MJO weakened east of the Date Line during June. Eastward propagation of broad suppressed convection continued into early July.


The upper-level footprint of the MJO re-emerged during mid-July, with a broad divergent signal propagating from the Maritime Continent to the central Pacific.

From mid-July to early September, a somewhat stationary pattern of enhanced (suppressed) convection over the east-central Pacific (Indian Ocean) has emerged, associated with the transition towards El Niño conditions. The suppressed phase of the MJO recently constructively interfered with the base state across the eastern Indian Ocean and western Maritime Continent.





# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A wave-1 pattern is apparent, though slightly noisy in the southern hemisphere. Enhanced (suppressed) convection spans the eastern Pacific to the Indian Ocean (Maritime Continent and West Pacific), associated with the robust MJO event presently across the Western Hemisphere. Enhanced convection over the western Indian Ocean has strengthened.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)


Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies

Mass is converging across the far eastern Pacific associated with convergent flow from the extratropics in both hemispheres.

Anomalous westerlies across much of the Atlantic likely contributed to the downturn in tropical cyclone activity after the active start to September. These westerlies are beginning to break down in the start of October.



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Weak westerly anomalies propagated eastward from the Indian Ocean to the Americas in early May; this pattern broke down in early June.

Anomalous westerlies amplified over the Maritime Continent in mid-June and propagated eastward at MJO-like phase speeds.

During August the intraseasonal pattern weakened, with Rossby wave activity influencing the West Pacific pattern.

Persistent anomalous westerlies continue over the far East Pacific, while easterly anomalies have been more prevalent over the central Pacific. Westerly anomalies have shifted eastward from the Indian Ocean over the Maritime Continent.



# Weekly Heat Content Evolution in the Equatorial Pacific


Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Negative upper-ocean heat content anomalies persisted in the central and eastern Pacific through December.

A downwelling Kelvin wave associated with the intraseasonal signal weakened the negative anomalies across the east-central Pacific during late January and early February.

Several downwelling oceanic Kelvin waves contributed to the eastward expansion of relatively warm subsurface water during February. Positive anomalies have now been observed over most of the basin since April.

The recent westerly wind burst east of New Guinea appears to have triggered another oceanic Kelvin wave and round of downwelling, helping to reinforce the warm water availability for a potential El Niño event.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.


The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast

Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

MJO activity is forecast by the GEFS to continue through Phase 1 into Phase 2 during Week-1, with a slight decline in amplitude.

After Week-2, the GEFS predicts the MJO signal to remain in Phase 2, decay rapidly, and fall back inside the unit circle.






# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 07 Oct 2018  
OLR




The GEFS shows the MJO signal remaining fairly stationary throughout the next two weeks. The enhanced (suppressed) convection is forecast to remain across the Indian Ocean (Maritime Continent and western Pacific), with a possible weakening early in Week-2, followed by re-strengthening.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [7.5°S,7.5°N] (cint:4Wm<sup>-2</sup>) Period:06-Apr-2018 to 06-Oct-2018  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (07 Oct 2018)




The constructed analog forecast closely mirrors the GEFS, though the re-strengthening late in Week-2 is not apparent. Enhanced convection emerging over the Indian Ocean with lower frequency signals across the eastern Pacific and Caribbean.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:07-Apr-2018 to 07-Oct-2018  
The unfilled contours are CA forecast reconstructed anomaly for 15 days




# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (May - Sep)




Precipitation Anomalies (May - Sep)



# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>