

Global Ocean Monitoring: Recent Evolution, Current Status, and Predictions

Prepared by

Climate Prediction Center, NCEP/NOAA

August 10, 2015

<http://www.cpc.ncep.noaa.gov/products/GODAS/>

**This project to deliver real-time ocean monitoring products is implemented
by CPC in cooperation with NOAA's Climate Observation Division (COD)**

Outline

- **Overview**

- **Recent highlights**

- Pacific/Arctic Ocean

- **El Niño conditions**

- **NE Pacific warming**

- Indian Ocean

- Atlantic Ocean

- **Global SST Predictions**

- **Is the amplitude of NINO3.4 going to exceed 2°C in winter 2015/2016?**

Overview

➤ Pacific Ocean

- ❑ El Niño conditions strengthened in July 2015 and the Nino34 index (+1.6°C) exceeded the threshold for a strong El Niño ($\geq 1.5^{\circ}\text{C}$).
- ❑ Most model predictions called for a strong El Niño through the Northern Hemisphere fall-winter 2015.
- ❑ Upper ocean warming associated with the "Blob" has persisted since winter 2013/2014.
- ❑ Positive PDO phase strengthened, with the PDO index increased from +0.7 to +1.5 in July.

➤ Indian Ocean

- ❑ Positive SSTAs dominated the whole Indian Ocean.


➤ Atlantic Ocean

- ❑ NAO switched to negative phase with NAOI = -3.1 in July.
- ❑ NOAA's updated hurricane outlook called for 90% chance of below-normal Atlantic hurricane season.

Global Oceans


Global SST Anomaly ($^{\circ}\text{C}$) and Anomaly Tendency

JUL 2015 SST Anomaly ($^{\circ}\text{C}$)
(1981–2010 Climatology)



- SSTA exceeded $+1.5^{\circ}\text{C}$ across the central and eastern equatorial Pacific Ocean.
- Strong positive SSTA presented in the NE Pacific Ocean and near the Bering Strait.
- Negative SSTA dominated in the subpolar north Atlantic.
- Positive SSTA persisted in the Indian and Southern Oceans.

JUL 2015 – JUN 2015 SST Anomaly ($^{\circ}\text{C}$)




- Positive SSTA tendency continued in the eastern equatorial Pacific.
- Positive(negative) SSTA tendency presented in the north subtropical Pacific (western Pacific).
- A strong cooling (warming) tendency was observed in the north high latitude of Atlantic (north subtropical Atlantic).

Fig. G1. Sea surface temperature anomalies (top) and anomaly tendency (bottom). Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1981-2010 base period means.


Longitude-Depth Temperature Anomaly and Anomaly Tendency in 2°S-2°N

JUL 2015 Eq. Temp Anomaly (°C)
(GODAS, Climo. 81-10)



- Strong positive ocean temperature anomalies persisted in the central-eastern equatorial Pacific.
- Positive temperature anomalies occupied most of the Indian Ocean.
- Negative temperature anomalies dominated the Atlantic Ocean.

JUL 2015 - JUN 2015 Eq. Temp Anomaly (°C)




- A warming (cooling) tendency presented in the western-central (eastern) Pacific.



Fig. G3. Equatorial depth-longitude section of ocean temperature anomalies (top) and anomaly tendency (bottom). Data are derived from the NCEP's global ocean data assimilation system which assimilates oceanic observations into an oceanic GCM. Anomalies are departures from the 1981-2010 base period means.

NOAA's Updated 2015 Atlantic Hurricane Season Outlook

(<http://www.cpc.ncep.noaa.gov/products/outlooks/hurricane2015/>)





Three tropical storms were formed in North Atlantic by Aug. 6.
Ten tropical storms with five hurricanes were formed in E. Pacific by Aug.6.



(<http://weather.unisys.com/hurricane/>)

Tropical Pacific Ocean and ENSO **Conditions**

Evolution of Pacific NINO SST Indices




- Niño 3.4, Niño 3 and Niño 1+2 indices strengthened in July 2015.
- Niño3.4 = 1.6 °C in July 2015 and ranks the second warmest July since 1982.
- Compared with last July, the central-eastern equatorial Pacific and the central and southern American coast were warmer in July 2015.


Fig. P1a. Niño region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the specified region. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1981–2010 base period means.

Last Three Month SST, OLR and 925hp Wind Anom.



MAY 2015 SST Anom. ($^{\circ}\text{C}$)




JUN 2015 SST Anom. ($^{\circ}\text{C}$)




JUL 2015 SST Anom. ($^{\circ}\text{C}$)





MAY 2015 OLR Anom. (W/m^2)
925hp Wind Anom (m/s)



JUN 2015 OLR Anom. (W/m^2)
925hp Wind Anom (m/s)




JUL 2015 OLR Anom. (W/m^2)
925hp Wind Anom (m/s)



- Positive SSTA strengthened and extended from the South American coast line to the central equatorial Pacific in the last three months.
- From May to July, negative OLR anomalies persisted over the central and eastern Pacific and westerly low-level winds prevailed across most of the equatorial Pacific.

Equatorial Pacific SST (°C), HC300 (°C), u850 (m/s) Anomalies


2°S–2°N Average, 3 Pentad Running Mean



- Strength of SST warm anomalies has strengthened since Apr. 2015 .
- A third downwelling kelvin wave was triggered by westerly wind burst in late June/early July.


Evolution of Equatorial Pacific Surface Zonal Current Anomaly (cm/s)

U (15m), cm/s, 2°S–2°N (Shading=Anomaly; Contour=Climatology)



-Anomalous eastward current extended from western Pacific to central-eastern Pacific since late June, partially attributed to the downwelling Kelvin wave.

NINO3.4 Heat Budget



- **Observed SSTA tendency (dT/dt) in NINO3.4 region (dotted black line) was positive since mid-Jan 2015.**

-**All dynamical terms (Q_u , Q_v , Q_w+Q_{zz}) were positive since Feb 2015, and heat flux term (Q_q) was negative.**

Huang, B., Y. Xue, X. Zhang, A. Kumar, and M. J. McPhaden, 2010 : The NCEP GODAS ocean analysis of the tropical Pacific mixed layer heat budget on seasonal to interannual time scales, *J. Climate.*, 23, 4901-4925.

Q_u : Zonal advection; Q_v : Meridional advection;




Q_w : Vertical entrainment; Q_{zz} : Vertical diffusion

Q_q : $(Q_{net} - Q_{open} + Q_{corr})/pcph$; $Q_{net} = SW + LW + LH + SH$;


Q_{open} : SW penetration; Q_{corr} : Flux correction due to relaxation to OI SST³

North Pacific & Arctic Oceans

PDO index



- Positive PDO has persisted 13 months since July 2014 and PDO index = 1.5 in July 2015.



- Pacific Decadal Oscillation is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly SST anomalies onto the 1st EOF pattern.

- The PDO index differs slightly from that of JISAO, which uses a blend of UKMET and OIv1 and OIv2 SST.

North Pacific & Arctic Ocean: SST Anom., SST Anom. Tend., OLR, SLP, 925hp wind, Sfc Rad, Sfc Flx






Fig. NP1. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sea surface pressure and 925hp wind anomalies (middle-right), sum of net surface short- and long-wave radiation anomalies (bottom-left), sum of latent and sensible heat flux anomalies (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, sea surface pressure and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1981-2010 base period means.

North America Western Coastal Upwelling

Pentad Coastal Upwelling for West Coast North America
($\text{m}^3/\text{s}/100\text{m}$ coastline)



Standard Positions of Upwelling Index Calculations




- Anomalous downwelling dominated south of 51°N in July 2015, consistent with the southwesterly wind anomalies.

Fig. NP2. Total (top) and anomalous (bottom) upwelling indices at the 15 standard locations for the western coast of North America. Upwelling indices are derived from the vertical velocity of the NCEP's global ocean data assimilation system, and are calculated as integrated vertical volume transport at 50 meter depth from each location to its nearest coast point ($\text{m}^3/\text{s}/100\text{m}$ coastline). Anomalies are departures from the 1981-2010 base period pentad means.


- Area below (above) black line indicates climatological upwelling (downwelling) season.
- Climatologically upwelling season progresses from Mar to Jul along the west coast of North America from 36°N to 57°N .

Last Three Month SSTA of North Pacific



MAY 2015 SST Anom. ($^{\circ}\text{C}$)




JUN 2015 SST Anom. ($^{\circ}\text{C}$)



JUL 2015 SST Anom. ($^{\circ}\text{C}$)




Temperature anomaly averaged at [150W–135W, 40N–50N]




- Strong positive SSTA persisted in the NE Pacific in the last three months .
- Strong subsurface temperature warming in the NE Pacific [150°w-135°w, 40°-50°N] persisted since 2013 winter.

Last Three Month SSTA , SLP, 925p Wind and Net Heat flux Anomalies


MAY 2015 SST Anom. (°C)




JUN 2015 SST Anom. (°C)




JUL 2015 SST Anom. (°C)




MAY 2015 SLP Anom.(hPa)
925hp Wind Anom(m/s)




JUN 2015 SLP Anom.(hPa)
925hp Wind Anom(m/s)




JUL 2015 SLP Anom.(hPa)
925hp Wind Anom(m/s)




MAY 2015 SW+LW+LH+SH (W/m²)



JUN 2015 SW+LW+LH+SH (W/m²)



JUL 2015 SW+LW+LH+SH (W/m²)






-3.5 -2.5 -1.5 -1 -0.5 0.5 1 1.5 2.5 3.5

-14 -10 -6 -4 -2 2 4 6 10 14

-80 -60 -40 -20 -10 10 20 40 60 80

Arctic Sea Ice

National Snow and Ice Data Center
<http://nsidc.org/arcticseaicenews/index.html>



- Arctic sea ice extent was well below average in July 2015.

Indian Ocean

Evolution of Indian Ocean SST Indices






Fig. I1a. Indian Ocean Dipole region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the SETIO [90°E–110°E, 10°S–0] and WTIO [50°E–70°E, 10°S–10°N] regions, and Dipole Mode Index, defined as differences between WTIO and SETIO. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1981–2010 base period means.

Tropical and North Atlantic Ocean

Evolution of Tropical Atlantic SST Indices

Monthly Tropical Atlantic SST Anomaly

(Bar: 1981–2010 Climatology; Curve: Last 10 YR Climatology)




- Negative SSTA dominated in the North subtropical Atlantic.
- ATL3 was near normal in July.
- Negative dipole index weakened in July 2015.

Fig. A1a. Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies ($^{\circ}\text{C}$) for the TNA [60 $^{\circ}\text{W}$ -30 $^{\circ}\text{W}$, 5 $^{\circ}\text{N}$ -20 $^{\circ}\text{N}$], TSA [30 $^{\circ}\text{W}$ -10 $^{\circ}\text{E}$, 20 $^{\circ}\text{S}$ -0] and ATL3 [20 $^{\circ}\text{W}$ -0, 2.5 $^{\circ}\text{S}$ -2.5 $^{\circ}\text{N}$] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1981-2010 base period means.




Tropical Atlantic:

SST Anom., SST Anom. Tend., TCHP OLR, Sfc Flx, 925-mb/200-mb Winds and RH



- Negative SSTA and TCHP continued in the hurricane Main Development Region (MDR) .
- Above-normal vertical wind shear was observed in MDR in July 2015.

NAO and SST Anomaly in North Atlantic




- NAO switched to strong negative phase with NAOI=-3.1 in July 2015.
- Tripole pattern continued in N. Atlantic Ocean.

Fig. NA2. Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N (<http://www.cpc.ncep.noaa.gov>). Time-Latitude section of SST anomalies averaged between 80°W and 20°W (bottom). SST are derived from the NCEP OI SST analysis, and anomalies are departures from the 1981-2010 base period means.


ENSO and Global SST Predictions

IRI NINO3.4 Forecast Plum


Mid-Jul 2015 Plume of Model ENSO Predictions



Mid-Jul IRI/CPC Plume-Based Probabilistic ENSO Forecast



Early-Jul CPC/IRI Consensus Probabilistic ENSO Forecast



- Majority of models predicted a moderate ($1^{\circ}\text{C} \leq \text{NINO3.4} < 1.5^{\circ}\text{C}$) to strong ($\text{NINO3.4} \geq 1.5^{\circ}\text{C}$) El Niño in winter 2015/2016.


- Models predicted NINO3.4 in consecutive five seasons exceeding $+2^{\circ}\text{C}$ (IRI web site):

Dynamical Models (9/16): NCEP CFSv2, NASA GMAO, AUS/POAMA, ECMWF, UKMO, COLA CCSM3, MetFRANCE, GFDL CM2.1, CMC CANSIP

Statistical Models (1/8): CPC Markov


SST, D20 and 925hp Wind anomalies in July

Nino 3.4 SST Anomaly




1982

JUL 1982 SST Anom. (°C)




JUL 1982 D20 Anom. (m)
925hp Wind Anom(m/s)




1997

JUL 1997 SST Anom. (°C)




JUL 1997 D20 Anom. (m)
925hp Wind Anom(m/s)




2015

JUL 2015 SST Anom. (°C)



JUL 2015 D20 Anom. (m)
925hp Wind Anom(m/s)





Real-Time Multiple Ocean Reanalyses Intercomparison

Jul 1982

Jul 2015

Anomalous Temperature (C) Averaged in 5S-5N: JUL 1982

Anomalous Temperature (C) Averaged in 5S-5N: JUL 2015



(http://origin.cpc.ncep.noaa.gov/products/GODAS/multiora_body.html)



Real-Time Multiple Ocean Reanalyses Intercomparison

Jul 1997

Jul 2015


Anomalous Temperature (C) Averaged in 5S-5N: JUL 1997

Anomalous Temperature (C) Averaged in 5S-5N: JUL 2015



(http://origin.cpc.ncep.noaa.gov/products/GODAS/multiora_body.html)


Anomalous Depth (m) of 20C Isotherm Averaged in [120E-80W, 5S-5N]



97



Anomalous Depth (m) of 20C Isotherm Averaged in [120E-80W, 5S-5N]
Ensemble Mean




- Warm Water Volume (D20 Anom. Ave. in 120E-80W, 5S-5N) in Jun-Jul 2015 is similar to Jun-Jul 1982.


NCEP CFSv2 North Pacific SST Predictions

Ics: 20150729-20150808


Aug 2015




Nov 2015




Sep 2015




Dec 2015




Oct 2015




Jan 2016



PDO



NPAC[150W-135W,40N-50N]




CFS Tropical North Atlantic (TNA) SST Predictions

from Different Initial Months

Tropical N. Atlantic SST anomalies (K)

TNA is the SST anomaly averaged in the region of [60°W-30°W, 5°N-20°N].



- Predictions initiated in Oct-Dec 2014 may be biased by errors in the Atlantic in CFSR.
- Latest CFS2 prediction calls a warming tendency in North Atlantic during summer-winter 2015.

Fig. M3. CFS Tropical North Atlantic (TNA) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1981-2010 base period means.

Overview

➤ **Pacific Ocean**

- ❑ **El Niño conditions strengthened in July 2015 and the Nino34 index (+1.6°C) exceeded the threshold for a strong El Niño ($\geq 1.5^\circ\text{C}$).**
- ❑ **Most model predictions called for a strong El Niño through the Northern Hemisphere fall-winter 2015.**
- ❑ **Upper ocean warming associated with the "Blob" has persisted since winter 2013/2014.**
- ❑ **Positive PDO phase strengthened, with the PDO index increased from +0.7 to +1.5.**

➤ **Indian Ocean**


- ❑ **Positive SSTAs dominated the whole Indian Ocean.**

➤ **Atlantic Ocean**


- ❑ **NAO switched to negative phase with NAOI = -3.1 in July.**
- ❑ **NOAA's updated hurricane outlook called for 90% chance of below-normal Atlantic hurricane season.**

Backup Slides

Tropical Cyclone Heat Potential Anomaly and Tendency (KJ/cm²)



- Strong positive TCHP anomalies presented in the central and eastern equatorial Pacific.
- TCHP anomalies were near normal over the tropical Atlantic Ocean.



TCHP field is the anomalous heat storage associated with temperatures larger than 26°C.

Global Sea Surface Salinity (SSS) Anomaly for July 2015

- NOTE: Since Aquarius terminated operations, the blended SSS analysis is from in situ and SMOS only from June 2015. Please report to us any suspicious data issues!
- Negative / positive SSS anomalies observed over the eastern / western equatorial Pacific, caused by changes in the fresh water fluxes in association with the enhanced / weakened ITCZ over the regions.

- Data used

- SSS :

- Blended Analysis of Surface Salinity (BASS) V0.Y
(a CPC-NESDIS/NODC-NESDIS/STAR joint effort)
(Xie et al. 2014)


- <ftp.cpc.ncep.noaa.gov/precip/BASS>

- Precipitation:

- CMORPH adjusted satellite precipitation estimates

- Evaporation:


- CFS Reanalysis



Global Sea Surface Salinity (SSS)

Tendency for July 2015


- Negative / positive SSS tendency presents over the eastern / western Pacific, consistent with the anomaly changes shown in last slide
- Positive SSS tendency also appears over the NW Atlantic off the North America continent




Global Sea Surface Salinity (SSS)

Anomaly Evolution over Equatorial Pacific

- Hovemoller diagram for equatorial SSS anomaly (**10°S-10°N**);
- Negative SSS anomaly strengthened with the maximum appears around 170°W. At the meantime, a stretch of positive SSS anomaly is developing over the western Pacific from 130°E – 160°E;



Equatorial Pacific Ocean Temperature Pentad Mean Anomaly



- Positive subsurface temperature anomalies in the eastern Pacific increased steadily in the last six pentads.

Tropical Pacific: SST Anom., SST Anom. Tend., OLR, Sfc Rad, Sfc Flx, 925-mb & 200-mb Winds






Fig. P2. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1981-2010 base period means.

Oceanic Kelvin

Standardized Projection on EEOF 1



(OKW index is defined as standardized projections of total anomalies onto the 14 patterns of Extended EOF1 of equatorial temperature anomalies (Seo and Xue, GRL, 2005).)

North Atlantic: SST Anom., SST Anom. Tend., OLR, SLP, Sfc Rad, Sfc Flx

-A dipole like SST pattern presented in North Atlantic.

- SSTA tendency was largely consistent with surface flux anomalies.




Fig. NA1. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sea surface pressure anomalies (middle-right), sum of net surface short- and long-wave radiation anomalies (bottom-left), sum of latent and sensible heat flux anomalies (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, sea surface pressure and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1981-2010 base period means.

Tropical Indian: SST Anom., SST Anom. Tend., OLR, Sfc Rad, Sfc Flx, 925-mb & 200-mb Wind Anom.

- SST warming occupied the whole basin.





Fig. 12. Sea surface temperature (SST) anomalies (top-left), anomaly tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the NCEP OI SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1981-2010 base period means.

CFS Niño3.4 SST Predictions from Different Initial Months

Niño3.4 SST anomalies (K)




- CFSv2 predicts a strong El Niño event through out the fall-winter 2015.

Fig. M1. CFS Niño3.4 SST prediction from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1981-2010 base period means.

CFS Pacific Decadal Oscillation (PDO) Index Predictions

from Different Initial Months

standardized PDO index



PDO is the first EOF of monthly ERSSTv3b anomaly in the region of [110°E-100°W, 20°N-60°N].

CFS PDO index is the standardized projection of CFS SST forecast anomalies onto the PDO EOF pattern.

- Forecast from July IC calls for above-normal PDO throughout northern hemisphere fall-winter 2015.

Fig. M4. CFS Pacific Decadal Oscillation (PDO) index predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1981-2010 base period means.

NCEP CFSv2 & NMME NINO 3.4 forecast






NWS/NCEP/CPC

Last update: Wed Aug 5 2015
Initial conditions: 26Jul2015-4Aug2015



NWS/NCEP/CPC


Last update: Wed Aug 5 2015
Initial conditions: 26Jul2015-4Aug2015



- Latest CFSv2 and NMME forecast called for a significant strong El Niño event (>2°C) throughout the Northern Hemisphere fall-winter 2015.

Be aware that new climatology (1981-2010) was applied since Jan 2011

SST Climatology Diff. ($^{\circ}\text{C}$): (1981–2010) – (1971–2000)



1971-2000 SST Climatology (Xue et al. 2003):

http://www.cpc.ncep.noaa.gov/products/predictions/30day/SSTs/sst_clim.htm

1981-2010 SST Climatology: <http://origin.cpc.ncep.noaa.gov/products/people/yxue/sstclim/>

- The seasonal mean SST in February-April (FMA) increased by more than 0.2°C over much of the Tropical Oceans and N. Atlantic, but decreased by more than 0.2°C in high-latitude N. Pacific, Gulf of Mexico and along the east coast of U.S.
- Compared to FMA, the seasonal mean SST in August-October (ASO) has a stronger warming in the tropical N. Atlantic, N. Pacific and Arctic Ocean, and a weaker cooling in Gulf of Mexico and along the east coast of U.S.

Switch to 1981-2010 Climatology

- **SST from 1971-2000 to 1981-2010**
 - Weekly **OISST.v2**, monthly ERSST.3b
- **Atmospheric fields from 1979-1995 to 1981-2010**
 - NCEP CDAS **winds**, sea level pressure, 200mb velocity potential, surface shortwave and longwave radiation, surface latent and sensible fluxes, relative humidity
 - Outgoing Long-wave Radiation
- **Oceanic fields from 1982-2004 to 1981-2010**
 - GODAS temperature, **heat content**, depth of 20°C, sea surface height, mixed layer depth, tropical cyclone heat potential, surface currents, upwelling
- **Satellite data climatology 1993-2005 unchanged**
 - Aviso Altimetry Sea Surface Height
 - Ocean Surface Current Analyses – Realtime (OSCAR)

Data Sources and References

- **Optimal Interpolation SST (OI SST) version 2 (Reynolds et al. 2002)**
- **NCEP CDAS winds, surface radiation and heat fluxes**
- **NESDIS Outgoing Long-wave Radiation**
- **NDBC TAO data (<http://tao.noaa.gov>)**
- **PMEL TAO equatorial temperature analysis**
- **NCEP's Global Ocean Data Assimilation System temperature, heat content, currents (Behringer and Xue 2004)**
- **Aviso Altimetry Sea Surface Height**
- **Ocean Surface Current Analyses – Realtime (OSCAR)**

Please send your comments and suggestions to Yan.Xue@noaa.gov. Thanks!