

Updating Climate Normals Impact on NMME Forecast

Qin Zhang (NOAA/CPC), Johnna Infanti (NOAA/CPC),
Sara Strey (Innovim/CPC), and Matthew Rosencrans
(NOAA/CPC)

Update NMME New Normals (1991-2020) for SST, Tmp2m and Prate

- GFDL_SPEAR - climo (1991-2020) applied in NMME ensemble mean (February 2021)
- NCAR_CCSM4, CanCM4i and GEM_NEMO - new climos calculated, compared with old climos, confirmed by model centers, ops and research lead (March 2021)
- NASA_GEOS5v2, CFSv2 and NMME - new climos calculated, working through comparisons and approved in April, 2021


- New climos of all models have been used in May's IC forecast, 2021

What NMME Recalculated with New Normals


- Each model's climatology (1991-2020) for 7 month forecast
- Monthly and seasonal Standard Deviation (SD) of each model based on 1991-2020 hindcast
- Anomalies of each models for 1991-2020 based on new climos
- Recalculated anomaly correlation coefficients (ACC) of forecast skills (1991-2020) for each model and NMME
- Three categories (above, normal and below) for probabilistic forecast
- Calibilation probability forecast from new adjustment of probability
anomaly correlation (PAC) for 1991-2020
- Nino3.4 plumes updated with new normals

SST Forecast Skills

1991-2020 climo




1982-2010 climo




Prate Forecast Skill

1991-2020 climo




1982-2010 climo




Tmp2m Forecast Skill

1991-2020 climo




1982-2010 climo




Nino3.4 Forecast

1991-2020 climo



1982-2010 climo



Summary

- Forecast NMME patterns of variables (SST, Tmp2m and Prate) keep similar to difference climos
- The amplitudes over some areas have been impacted by changing new normals to reduce warm and increase cold.
- The NMME forecast skills are increased for some regions for new climos, such as SST in the tropical.