

Attribution of Seasonal Climate Anomalies

January–February–March 2020


Summary of January–February–March 2020 Observed Conditions and Outlooks

- The sea surface temperature (SST) anomalies in equatorial central Pacific Ocean continued to be on the weak warm side; the long lasting strong positive phase of Indian Ocean Dipole (IOD) SST anomaly pattern (since summer 2019) changed to positive SST anomalies over the entire Indian Ocean in JFM2020; positive SST anomalies throughout the NE Pacific weakened further (Slide 4).
- The observed drier than normal precipitation persisted over the region extending from Indonesian Archipelago to southeast equatorial Indian Ocean; the observed positive precipitation anomalies over the equatorial central Pacific was consistent with the underlying positive SST anomalies (Slide 5).
- The large scale pattern of the precipitation anomalies over the tropical areas were generally well predicted in CFSv2 (slide 11) and the multi-model ensemble forecasts from the WMO Lead Center (slides 39-40).
- The large scale structures of the observed positive height anomalies over the tropical and N. sub-tropical region were well captured in the CFSv2 forecasts, while the observed negative height anomalies at high latitudes and polar region were missed leading to missed cold surface temperature anomalies over extreme northern parts of NA including Alaska.
- NA z200, precip, and T2m JFM2020 forecasts had noticeable improvements from the AMIP simulations to the seasonal forecasts, and further improvements in the constructed seasonal forecasts (slides 14-16).


Observed Seasonal Anomalies

Global and North America




Observed Anomaly JFM2020



Observed Anomaly JFM2020





Observed Anomaly JFM2020



JFM2020 CPC Seasonal Outlooks and NMME Forecasts

Temperature


Temp nonEC
HSS=16




Precipitation

CPC



Jan-Feb-Mar 2020



Prec nonEC
HSS=22



NMME





For the rationale behind CPC outlooks see https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2019/201912_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies





Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
 - **Seasonal forecast**: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - **Reconstructed forecast**: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach fr constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).





JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)







JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)







JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)







JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



JFM2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs


Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.



JFM2020 Anomaly Correlation for Individual AMIP Simulation with Observation -- z200(20N–90N)







Observed & AMIP Ensemble Average Anomalies
JFM2020 z200(m) 18 runs/worst 2 runs/best 2 runs




JFM2020 Anomaly Correlation for Individual CFSv2 Forecast
with Observation --- z200 (20N–90N)







Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 z200(m) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 z200(m) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast







JFM2020 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)







Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast







JFM2020 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 T2m(K) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
JFM2020 T2m(K) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast



200mb Height from Linear Model


JFM2020 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N

OPI Prate Anom (mm/day)



OBS (from R1)


Linear Model Response to Tropical Heating




Pattern COR: global=0.11, tropics(30S–30N)=0.15

Seasonal Forecasts from the Constructed Analog Model


CA HGT200 Prd for JFM2020, ICs through Mar2020(m), Lead -3



CA T2m Prd for JFM2020, ICs through Mar2020(K), Lead -3



CA Prec Prd for JFM2020, ICs through Mar2020(mm/day), Lead -3



Seasonal Forecasts from WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME)

<https://www.wmolk.org/>

- LC-LRFMME seasonal forecast are based on forecasts provided by WMO recognized Global Producing Centers (GPCs) for Long-Range Forecasts to the LC-LRFMME. Contribution of all GPCs is acknowledged.
- Seasonal forecasts from GPCs are merged into a multi-model ensemble forecast.
- LC-LRFMME forecasts are based on GPC seasonal forecast systems run during the first week of the month for the next season. For example, forecasts runs in first week of January for the seasonal mean of February-March-April.
- Forecasts in slides 42-45 are from the Lead Center.
- For latest seasonal outlook guidance see
<http://www.wmo.int/pages/prog/wcp/wcasp/LC-LRFMME/index.php>
- *For more information see visit Lead Center website; also see Graham, R., and Co-authors, 2011: New perspectives for GPCs, their role in the GFCS and a proposed contribution to a 'World Climate Watch'. Climate Research, 47, 47-55.*


Simple Composite Map

Melbourne, Montreal, Moscow, Offenbach, Seoul

[Unit : K]

2m Temperature : JFM2020

(issued on Dec2019)



WMO Lead Centre
LRF MME


Simple Composite Map

Melbourne, Montreal, Moscow, Offenbach, Seoul

[Unit : mm]

Precipitation : JFM2020

(issued on Dec2019)




Probabilistic Multi-Model Ensemble Forecast

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Pretoria, Seoul, Tokyo, Toulouse, Washington

2m Temperature : JFM2020

(issued on Dec2019)




WMO Lead Centre
LRF MMRE

Probabilistic Multi-Model Ensemble Forecast

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Pretoria, Seoul, Tokyo, Toulouse, Washington

Precipitation : JFM2020

(issued on Dec2019)



WMO Lead Centre
for
LRF MME

References

- Fan, Y., and Dool H. van den Dool (2008), A global monthly land surface air temperature analysis for 1948-present. *J. Geophys. Res.*, 113, D01103. [doi:10.1029/2007JD008470](https://doi.org/10.1029/2007JD008470).
- Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).
- Reynolds, R. W. et al (2007), Daily high resolution-blended analyses for sea surface temperature. *J. Clim.*, 20, 5473-5496. [doi:10.1175/2007JCLI1824.1](https://doi.org/10.1175/2007JCLI1824.1).
- Saha, S. et al (2010), The NCEP climate forecast system reanalysis. *Bull. Amer. Meteor. Soc.*, 91, 1015-1057. [doi:10.1175/2010BAMS3001.1](https://doi.org/10.1175/2010BAMS3001.1).
- Saha, S. et al (2014), The NCEP climate forecast system version 2. *J. Clim.*, 27, 2185-2208. [doi:10.1175/JCLI-D-12-00823.1](https://doi.org/10.1175/JCLI-D-12-00823.1).
- Xie, P, and P. A. Arkin (1997), Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bull. Amer. Meteor. Soc.*, 78, 2539-2558. doi: [http://dx.doi.org/10.1175/1520-0477\(1997\)078%3C2539:GPAYMA%3E2.0.CO;2](http://dx.doi.org/10.1175/1520-0477(1997)078%3C2539:GPAYMA%3E2.0.CO;2)

Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: NCDC daily OI analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
 - 18 members
- All above seasonal mean anomalies are based on 1999-2010 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)