

# Attribution of Seasonal Climate Anomalies

## April-May-June 2020

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


# Summary of April-May-June 2020 Observed Conditions and Outlooks

- The sea surface temperature (SST) anomalies in the equatorial central Pacific Ocean continued to be on weak warm side (Slide 4) with central/eastern Pacific getting cooler.
- Except over eastern and northern regions of North America, land temperature anomalies everywhere were generally above normal (especially over northern Asia) and was predicted well in MME models (slide 5, 13, 33, 34, and 38).
- In the tropics, the large-scale distribution of negative and positive precipitation anomalies was predicted in CFSv2 and other multi-model ensemble (MME) forecast systems (slide 11, 33, 34, & 39).
- The north-south pattern of high and low height anomalies between 60°N-90°N ([with positive projection on the AO](#)) in previous 3-mon-means became weaker in AMJ2020. The initialized CFSv2 forecasts captured well the large-scale structures of the observed positive height anomalies over the tropical and N. sub-tropical region, while it missed negative anomalies over the NA leading to missed cold surface temperature over the region from Canada to the US mid-east and south-east (slide 12, 13, 15, & 16).
- The monthly forecasts from the shortest 3-day-leads initial conditions captured major features of NA precipitation and T2m anomalies, especially for May2020 T2m and June2020 precipitation (slides 30 & 31).
- There is some tendency for lead time dependence (relatively higher skills for shorter lead forecasts) in monthly forecasts, and less so for seasonal forecasts (slide 21, 24, & 27).




# Observed Seasonal Anomalies

## Global and North America




# Observed Anomaly AMJ2020



# Observed Anomaly AMJ2020




# Observed Anomaly AMJ2020




# AMJ2020 CPC Seasonal Outlooks and NMME Forecasts

Temp nonEC  
HSS=-29




Temperature




Precipitation


Prec nonEC  
HSS=4



CPC



NMME





For the rationale behind CPC outlooks see [https://www.cpc.ncep.noaa.gov/products/archives/long\\_lead/PMD/2020/202003\\_PMD90D](https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2020/202003_PMD90D)

## Model Simulated/Forecast Ensemble Mean Anomalies





# Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
  - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
  - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach fr constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).





# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)







# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)







# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)







# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



# AMJ2020 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




# Model Simulated/Forecast Anomalies: Individual Runs

# Model Simulated/Forecast Anomalies: Individual Runs


- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? Geophys. Res. Lett., 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).

# AMJ2020 Anomaly Correlation for Individual AMIP Simulation with Observation -- z200(20N–90N)




# Observed & AMIP Ensemble Average Anomalies





## AMJ2020 z200(m) 18 runs/worst 2 runs/best 2 runs




# AMJ2020 Anomaly Correlation for Individual CFSv2 Forecast with Observation --- z200 (20N–90N)




Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 z200(m) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 z200(m) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast







# AMJ2020 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast







# AMJ2020 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 T2m(K) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies  
AMJ2020 T2m(K) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast



# Monthly Means Prec(mm/day) Observed & Forecasts

Monthly AMJ2020 Prec(mm/day) Observed & Forecasts




Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.

Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

# Monthly Means T2m(K) Observed & Forecasts

Monthly AMJ2020 T2m(K) Observed & Forecasts



Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.


Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

# Seasonal Forecasts from other multi-model systems and linear models


# C3S Seasonal Forecast

([https://climate.copernicus.eu/charts/c3s\\_seasonal/](https://climate.copernicus.eu/charts/c3s_seasonal/))

C3S multi-system seasonal forecast  
Mean 2m temperature anomaly  
Nominal forecast start: 01/03/20  
Variance-standardized mean





C3S multi-system seasonal forecast  
Mean precipitation anomaly  
Nominal forecast start: 01/03/20  
Variance-standardized mean




# North American Multi-Model Ensemble Seasonal Forecast




(<https://www.cpc.ncep.noaa.gov/products/NMME/>)



# 200mb Height from Linear Model



# Seasonal Forecasts from the Constructed Analog Model



# Seasonal Forecasts from WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME)

<https://www.wmolk.org/>

- LC-LRFMME seasonal forecast are based on forecasts provided by WMO recognized Global Producing Centers (GPCs) for Long-Range Forecasts to the LC-LRFMME. Contribution of all GPCs is acknowledged.
- Seasonal forecasts from GPCs are merged into a multi-model ensemble forecast.
- LC-LRFMME forecasts are based on GPC seasonal forecast systems run during the first week of the month for the next season. For example, forecasts runs in first week of January for the seasonal mean of February-March-April.
- Forecasts in slides 42-45 are from the Lead Center.
- For latest seasonal outlook guidance see  
<http://www.wmo.int/pages/prog/wcp/wcasp/LC-LRFMME/index.php>
- *For more information see visit Lead Center website; also see Graham, R., and Co-authors, 2011: New perspectives for GPCs, their role in the GFCS and a proposed contribution to a 'World Climate Watch'. Climate Research, 47, 47-55.*


# Simple Composite Map

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

[Unit : K]

2m Temperature : AMJ2020

(issued on Mar2020)



WMO Lead Centre  
for  
LRF MME


# Simple Composite Map

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

[Unit : mm]

Precipitation : AMJ2020

(issued on Mar2020)




WMO Lead Centre for  
LRF MME

# Probabilistic Multi-Model Ensemble Forecast

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

2m Temperature : AMJ2020

(issued on Mar2020)




WMO Lead Center for  
LRF MME

# Probabilistic Multi-Model Ensemble Forecast

CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

Precipitation : AMJ2020

(issued on Mar2020)



# Background & Methodology

# Attribution of Seasonal Climate Anomalies

- Goal
  - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
  - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

# Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
  - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
  - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

## Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

# Data

- Observations
  - SST: NCDC daily OI analysis (Reynolds et al., 2007)
  - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
  - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
  - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
  - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
  - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
  - 18 members
- All above seasonal mean anomalies are based on 1999-2010 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)