

Attribution of Seasonal Climate Anomalies

February-March-April 2021

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


Summary of Observed Conditions and Outlooks

- SST cold anomalies weakened in the central/eastern Pacific; tropical Indian Ocean and western and NE Pacific Ocean remained on the warm side. Also, one feature to note is the prevalence of cold SST anomalies (blue) in global oceans (slide 4).
- In general, the large-scale distribution of negative and positive precipitation anomalies in the equatorial Pacific and over the Maritime Continent ([a reflection of La Niña conditions](#)) was well predicted in CFSv2 and other MME forecast systems (slides 11, 33, 34, & 39).
- The average of FMA2021 height anomalies was in [positive phase of AO](#) pattern. The initialized CFSv2 and other multi-model ensemble MME forecast didn't capture the observed negative height anomalies over high latitudes leading to errors in forecasting cold temperature anomalies over North America, north Asia, and eastern Europe (slides 12, 13, 15, 16, 33, 34, & 38). Local trough over NA, however, was evident in initialized predictions (slide 15).
- The forecast NA temperature and precipitation anomalies in the CFSv2 and MME models were in the pattern consistent with [the La Niña composite](#), while they were different from the observations (slides 14, 16, 33, 34, 38, 39).
- The Apr2021 monthly forecasts from the shortest 3-day-leads initial conditions also missed most of observed temperature and precipitation anomalies (slides 30, 31). Extreme cold temperature anomaly over NA in February 2021 also reflected in the FMA 2021 seasonal mean (slide 31).

Observed Seasonal Anomalies


Global and North America

Observed Anomaly FMA2021




Observed Anomaly FMA2021



Prec(mm/day)






T2m(K)



z200(m)






Observed Anomaly FMA2021





CPC Seasonal Outlooks and NMME Forecasts


Temp nonEC
HSS=-14



Prec nonEC
HSS=10



NMME





For the rationale behind CPC outlooks see: https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2021/202101_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies





Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach fr constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).





FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)







FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)







FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)







FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



FMA2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs





Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? Geophys. Res. Lett., 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).


FMA2021 Anomaly Correlation for Individual AMIP Simulation with Observation -- z200(20N–90N)







Observed & AMIP Ensemble Average Anomalies
FMA2021 z200(m) 18 runs/worst 2 runs/best 2 runs







FMA2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N–90N)




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 z200(m) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 z200(m) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast







FMA2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast







FMA2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 T2m(K) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2021 T2m(K) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast



Monthly Means Prec(mm/day) Observed & Forecasts

Monthly FMA2021 Prec(mm/day) Observed & Forecasts




Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.

Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

Monthly Means T2m(K) Observed & Forecasts

Monthly FMA2021 T2m(K) Observed & Forecasts



Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.

Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

Seasonal Forecasts from other multi-model systems and linear models

C3S Seasonal Forecast


(https://climate.copernicus.eu/charts/c3s_seasonal/)

C3S multi-system seasonal forecast

Mean 2m temperature anomaly

Nominal forecast start: 01/01/21

Variance-standardized mean



ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA


FMA 2021

C3S multi-system seasonal forecast

Mean precipitation anomaly


Nominal forecast start: 01/01/21

Variance-standardized mean




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)






200mb Height from Linear Model

FMA2021 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N



Seasonal Forecasts from the Constructed Analog Model



Seasonal Forecasts from WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME)

<https://www.wmolk.org/>

- LC-LRFMME seasonal forecast are based on forecasts provided by WMO recognized Global Producing Centers (GPCs) for Long-Range Forecasts to the LC-LRFMME. Contribution of all GPCs is acknowledged.
- Seasonal forecasts from GPCs are merged into a multi-model ensemble forecast.
- LC-LRFMME forecasts are based on GPC seasonal forecast systems run during the first week of the month for the next season. For example, forecasts runs in first week of January for the seasonal mean of February-March-April.
- Forecasts in slides 42-45 are from the Lead Center.
- For latest seasonal outlook guidance see
<http://www.wmo.int/pages/prog/wcp/wcasp/LC-LRFMME/index.php>
- *For more information see visit Lead Center website; also see Graham, R., and Co-authors, 2011: New perspectives for GPCs, their role in the GFCS and a proposed contribution to a 'World Climate Watch'. Climate Research, 47, 47-55.*


Simple Composite Map

Beijing,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

[Unit : K]

2m Temperature : FMA2021

(issued on Jan2021)




Simple Composite Map

Beijing,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

[Unit : mm]

Precipitation : FMA2021

(issued on Jan2021)




WMO Lead Center for
LRF MMRE

Probabilistic Multi-Model Ensemble Forecast

Beijing,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

2m Temperature : FMA2021

(issued on Jan2021)




WMO Lead Centre
for
LRF MME

Probabilistic Multi-Model Ensemble Forecast

Beijing,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

Precipitation : FMA2021

(issued on Jan2021)



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: NCDC daily OI analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
 - 18 members
- All above seasonal mean anomalies are based on 1999-2010 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)