

Attribution of Seasonal Climate Anomalies April-May-June 2021

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


Summary of Observed Conditions and Outlooks

- SST cold anomalies weakened in the central/eastern Pacific; Warm SST anomalies weakened in the tropical Indian Ocean; NE Pacific Ocean remained on the warm side (slide 4).
- In general, the large-scale distribution of SST anomalies and precipitation anomalies over tropics were forecast reasonably well for CFSv2 and other MME forecast systems (slides 10, 11, 33, 34, & 39).
- The average of AMJ2021 height anomalies had a weak projection on [positive phase of AO](#) pattern. The AMIP simulation and initialized CFSv2 forecasts didn't predict well the variations of z200 anomalies over the extra-tropics, that led to errors in forecasting cold anomalies over the areas of east Asia, central Canada, and, the mid-SE US (slides 12, 13, 15, 16).
- The ensemble means of CFSv2 and MME models did not capture the observed anomalies in NA temperature and precipitation, while the best 4 members of reconstructed CFSv2 forecasts resembled well the observations (slides 26, 29).
- June 2021 monthly forecasts from the shortest 3-day-leads initial conditions captured the observed precipitation and temperature anomalies, especially, the positive precipitation anomalies and cold anomalies over the central to south-east US and warm temperatures over the west coast of the US (slides 30, 31).


Observed Seasonal Anomalies

Global and North America


Observed Anomaly AMJ2021



Observed Anomaly AMJ2021



Observed Anomaly AMJ2021



CPC Seasonal Outlooks and NMME Forecasts


Temperature

Precipitation


Temp nonEC
HSS=-8

Prec nonEC
HSS=14

CPC




NMME prob fcst TMP2m IC=202103 for lead 1 2021 AMJ



NMME

NMME prob fcst Prate IC=202103 for lead 1 2021 AMJ






For the rationale behind CPC outlooks see: https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2021/202103_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies


Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
 - **Seasonal forecast**: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - **Reconstructed forecast**: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).


AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)




AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



AMJ2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs

Model Simulated/Forecast Anomalies: Individual Runs


- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).

AMJ2021 Anomaly Correlation for Individual AMIP Simulation with Observation — z200(20N–90N)




Observed & AMIP Ensemble Average Anomalies


AMJ2021 z200(m) 18 runs/worst 2 runs/best 2 runs




AMJ2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N-90N)



Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 z200(m) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 z200(m) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast




AMJ2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S-30N)

— ensemble mean


■ 0-1-d-L ■ 2-3-d-L ■ 4-5-d-L ■ 6-7-d-L ■ 8-9-d-L



Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast




AMJ2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S-30N)

— ensemble mean


■ 0-1-d-L ■ 2-3-d-L ■ 4-5-d-L ■ 6-7-d-L ■ 8-9-d-L



Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 T2m(K) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies AMJ2021 T2m(K) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast



Monthly Means Prec(mm/day) Observed & Forecasts

Monthly AMJ2021 Prec(mm/day) Observed & Forecasts




Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.

Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

Monthly Means T2m(K) Observed & Forecasts

Monthly AMJ2021 T2m(K) Observed & Forecasts



Top row: Observed anomaly

Middle row: CFSv2 seasonal forecasts from the initial conditions from the month prior to the target season.

Bottom row: CFSv2 monthly forecasts from the last three days of the month prior to the target month.

Seasonal Forecasts from other multi-model systems and linear models

C3S Seasonal Forecast

(https://climate.copernicus.eu/charts/c3s_seasonal/)

C3S multi-system seasonal forecast

Mean 2m temperature anomaly


Nominal forecast start: 01/03/21

Variance-standardized mean

ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA

AMJ 2021

■ <-2.0°C ■ -2.0..-1.0 ■ -1.0..-0.5 ■ -0.5..-0.2 ■ -0.2..0.2 ■ 0.2..0.5 ■ 0.5..1.0 ■ 1.0..2.0 ■ > 2.0°C



C3S multi-system seasonal forecast

Mean precipitation anomaly


Nominal forecast start: 01/03/21

Variance-standardized mean

ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA

AMJ 2021


■ <-200mm ■ -200..-100 ■ -100..-50 ■ -50..-10 ■ -10..10 ■ 10..50 ■ 50..100 ■ 100..200 ■ > 200mm




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)

NMME Forecast of TMP2m Anom IC=202103 for Lead 1 2021AMJ





NMME Forecast of Prec. rate Anom IC=202103 for Lead 1 2021AMJ



200mb Height from Linear Model


AMJ2021 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N



Pattern COR: global=0.12, tropics(30S–30N)=0.25

Seasonal Forecasts from the Constructed Analog Model


CA HGT200 Prd for AMJ2021, ICs through Jun2021(m), Lead -3



Petao Peng CPC/NCEP/NWS/NOAA

Base Period 1991-2020


CA T2m Prd for AMJ2021, ICs through Jun2021(K), Lead -3



Petao Peng CPC/NCEP/NWS/NOAA

Base Period 1991-2020

CA Prec Prd for AMJ2021, ICs through Jun2021(mm/day), Lead -3



Petao Peng CPC/NCEP/NWS/NOAA

Base Period 1991-2020

Seasonal Forecasts from WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME)

<https://www.wmolc.org/>

- LC-LRFMME seasonal forecast are based on forecasts provided by WMO recognized Global Producing Centers (GPCs) for Long-Range Forecasts to the LC-LRFMME. Contribution of all GPCs is acknowledged.
- Seasonal forecasts from GPCs are merged into a multi-model ensemble forecast.
- LC-LRFMME forecasts are based on GPC seasonal forecast systems run during the first week of the month for the next season. For example, forecasts runs in first week of January for the seasonal mean of February-March-April.
- Forecasts in slides 42-45 are from the Lead Center.
- For latest seasonal outlook guidance see <http://www.wmo.int/pages/prog/wcp/wcasp/LC-LRFMME/index.php>
- *For more information see visit Lead Center website; also see Graham, R., and Co-authors, 2011: New perspectives for GPCs, their role in the GFCS and a proposed contribution to a 'World Climate Watch'. Climate Research, 47, 47-55.*



Simple Composite Map

Beijing, CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

[Unit : K]

2m Temperature : AMJ2021

(issued on Mar2021)



WMO Lead Centre for LRF MME



Simple Composite Map

Beijing, CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

[Unit : mm]

(issued on Mar2021)

Precipitation : AMJ2021




Probabilistic Multi-Model Ensemble Forecast

Beijing, CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

2m Temperature : AMJ2021

(issued on Mar2021)




Probabilistic Multi-Model Ensemble Forecast

Beijing, CPTEC, ECMWF, Exeter, Melbourne, Montreal, Moscow, Offenbach, Seoul, Tokyo, Toulouse, Washington

Precipitation : AMJ2021

(issued on Mar2021)



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: NCDC daily OI analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
 - 18 members
- All above seasonal mean anomalies are based on 1999-2010 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)