

# Attribution of Seasonal Climate Anomalies September-October-November 2021

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


# Summary of Observed Conditions and Outlooks

- SON2021 SST cold anomalies increased compared to ASO2021 in the equatorial central and eastern Pacific; the equatorial Atlantic and N Pacific Ocean remained on the warm side (slide 4). In general, the large-scale distribution of SST anomalies were predicted well (slide 10);
- The large scale distribution of dry, wet, and dry precipitation anomalies respectively in the equatorial western Indian Ocean, the eastern Indian Ocean - the Maritime Continent, and the equatorial western and central Pacific Ocean were predicted well in the initialized CFSv2 forecasts and other MME models (slides 11, 35, 36, 38 & 39).
- The initialized CFSv2 forecasts predicted most of observed 200mb height anomalies globally, except for the low (high) 200mb height anomalies in the region of Bering Strait and Alaska (central Canada); the forecasts also missed the large warm temperature anomalies over northern Canada; while the reconstructed z200mb and T2m seasonal mean had a better improved correlation skill (slides 12, 13, 15, &16).
- T2m anomalies over the US were well predicted by the CPC seasonal outlook (slide 7).
- The Oct. 2021 monthly mean forecasts with the shortest leads predicted well the large warm temperature anomalies in Canada (slide 32).


# Observed Seasonal Anomalies

## Global and North America




# Observed Anomaly SON2021



# Observed Anomaly SON2021




# Observed Anomaly SON2021




# CPC Seasonal Outlooks and NMME Forecasts

Temp nonEC  
HSS=84




Temperature

CPC




Precipitation

Prec nonEC  
HSS=36




NMME prob fct TMP2m IC=202108 for lead 1 2021 SON



NMME

NMME prob fct Prate IC=202108 for lead 1 2021 SON





For the rationale behind CPC outlooks see: [https://www.cpc.ncep.noaa.gov/products/archives/long\\_lead/PMD/2021/202108\\_PMD90D](https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2021/202108_PMD90D)

# Model Simulated/Forecast Ensemble Mean Anomalies





# Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
  - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
  - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach fr constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).





# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)







# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)







# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)







# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)







# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



# SON2021 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




# Model Simulated/Forecast Anomalies: Individual Runs

# Model Simulated/Forecast Anomalies: Individual Runs


- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? Geophys. Res. Lett., 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).

# SON2021 Anomaly Correlation for Individual AMIP Simulation with Observation -- z200(20N–90N)




# Observed & AMIP Ensemble Average Anomalies





## SON2021 z200(m) 18 runs/worst 2 runs/best 2 runs







# SON2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N–90N)




Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 z200(m) 40 runs/worst 4 runs/best 4 runs  
**Seasonal Forecast**







Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 z200(m) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast




# SON2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
**Seasonal Forecast**







Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
**Reconstructed Forecast**







# SON2021 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 T2m(K) 40 runs/worst 4 runs/best 4 runs  
**Seasonal Forecast**




Observed & CFSv2 Forecast Ensemble Average Anomalies  
SON2021 T2m(K) 40 runs/worst 4 runs/best 4 runs  
**Reconstructed Forecast**



# Monthly Means Prec(mm/day) Observed & Forecasts


## Monthly SON2021 Prec(mm/day) Observed & Forecasts

(at shortest leads)



# Monthly Means Prec(mm/day) Observed & Forecasts

## Monthly SON2021 Prec(mm/day) Observed & Forecasts (at long leads)



Top row: Observed anomaly

Middle row: CFSv2 monthly forecasts from the initial conditions near 10<sup>th</sup> day of the month prior to the target month.

Bottom row: CFSv2 monthly forecasts from the initial conditions near 20<sup>th</sup> day of the month prior to the target month.

# Monthly Means T2m(K) Observed & Forecasts

## Monthly SON2021 T2m(K) Observed & Forecasts


(at shortest leads)

Sep2021

Oct2021

Nov2021

SON2021



# Monthly Means T2m(K) Observed & Forecasts

## Monthly SON2021 T2m(K) Observed & Forecasts


(at long leads)

Sep2021

Oct2021

Nov2021

SON2021



Top row: Observed anomaly

Middle row: CFSv2 monthly forecasts from the initial conditions near 10<sup>th</sup> day of the month prior to the target month.

Bottom row: CFSv2 monthly forecasts from the initial conditions near 20<sup>th</sup> day of the month prior to the target month.



# Seasonal Forecasts from WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME)

<https://www.wmolk.org/>

- LC-LRFMME seasonal forecast are based on forecasts provided by WMO recognized Global Producing Centers (GPCs) for Long-Range Forecasts to the LC-LRFMME. Contribution of all GPCs is acknowledged.
- Seasonal forecasts from GPCs are merged into a multi-model ensemble forecast.
- LC-LRFMME forecasts are based on GPC seasonal forecast systems run during the first week of the month for the next season. For example, forecasts runs in first week of January for the seasonal mean of February-March-April.
- Forecasts in slides 42-45 are from the Lead Center.
- For latest seasonal outlook guidance see  
<http://www.wmo.int/pages/prog/wcp/wcasp/LC-LRFMME/index.php>
- *For more information see visit Lead Center website; also see Graham, R., and Co-authors, 2011: New perspectives for GPCs, their role in the GFCS and a proposed contribution to a 'World Climate Watch'. Climate Research, 47, 47-55.*

# LC-LRFMM Seasonal Forecasts (ensemble means)

(<https://www.wmoclc.org/>)



# LC-LRFMM Seasonal Forecasts (probabilities)


(<https://www.wmoclc.org/>)

## Probabilistic Multi-Model Ensemble Forecast

Beijing,Bologna,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

### 2m Temperature : SON2021

(issued on Aug2021)




## Probabilistic Multi-Model Ensemble Forecast

Beijing,Bologna,CPTEC,ECMWF,Exeter,Melbourne,Montreal,Moscow,Offenbach,Seoul,Tokyo,Toulouse,Washington

### Precipitation : SON2021

(issued on Aug2021)



# Seasonal Forecasts from other multi-model systems and linear models

# C3S Seasonal Forecast

([https://climate.copernicus.eu/charts/c3s\\_seasonal/](https://climate.copernicus.eu/charts/c3s_seasonal/))

C3S multi-system seasonal forecast


Mean 2m temperature anomaly

Nominal forecast start: 01/08/21

Variance-standardized mean

ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA/ECCC

SON 2021



C3S multi-system seasonal forecast


Mean precipitation anomaly

Nominal forecast start: 01/08/21

Variance-standardized mean



ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA/ECCC

SON 2021




# North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)






# 200mb Height from Linear Model

SON2021 200mb Eddy HGT(m)  
OBS vs. Linear Model Response to Tropical Heating  
Heating is converted from Prate in 15S–15N



# Seasonal Forecasts from the Constructed Analog Model



# Background & Methodology

# Attribution of Seasonal Climate Anomalies

- Goal
  - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
  - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

# Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
  - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
  - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

## Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

# Data

- Observations
  - SST: OI version 2 analysis (Reynolds et al., 2007)
  - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
  - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
  - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
  - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
  - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
  - 18 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)