

Attribution of Seasonal Climate Anomalies January-February-March 2022

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


Summary of Observed Conditions and Outlooks

- Tropical SST anomalies continued in La Nina conditions; the equatorial Atlantic and North Pacific SST anomalies remained on the warm side (slide 4). In general, the large-scale distribution of SST anomalies was predicted well (slide 10);
- Large-scale distribution of wet and dry precipitation anomalies in the equatorial eastern Indian Ocean, Maritime Continent and equatorial western, central Pacific Ocean ([a reflection of La Niña conditions](#)) were predicted well in the initialized CFSv2 and other MME models (slides 37-39).
- Initialized CFSv2 forecasts, in general, predicted the large-scale distribution of observed 200mb height anomalies, while the forecasts misplaced the observed locations of trough and low anomaly center over the northeast North America (slides 12, 15), leading to errors in forecasting cold anomalies over the region from central of Canada to central and southwest US (slides 7,13, 16).
- The spatial pattern of NA precipitation anomalies in the CFSv2 and MME models was consistent with [the La Niña composite](#) and captured the observed dry conditions over CA and southern US areas (Slides 7, 14).
- The monthly means from CFSv2 seasonal forecasts of z200, T2m, & Prec were in pattern consistent with the canonical La Nina signal (slides 30-32).
- March 2022 monthly mean forecasts from the shortest leads didn't predict the observed cold anomalies over the US central areas and most precipitation anomalies in general (slides 34, 35).


Observed Seasonal Anomalies

Global and North America


Observed Anomaly JFM2022



Observed Anomaly JFM2022



Observed Anomaly JFM2022



CPC Seasonal Outlooks and NMME Forecasts

Temperature

Precipitation

Temp nonEC
HSS=-26





Prec nonEC
HSS=39

CPC


NMME

Seasonal Temperature Outlook


Seasonal Precipitation Outlook



NMME prob fcst TMP2m IC=202112 for lead 1 2022 JFM



NMME prob fcst Prate IC=202112 for lead 1 2022 JFM






For the rationale behind CPC outlooks see: https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2021/202112_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies


Model Simulated/Forecast Ensemble Average Anomalies

- CFS **AMIP simulations** forced with observed sea surface temperatures (18 members ensemble)
- CFSv2 real time operational forecasts
 - **Seasonal forecast**: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - **Reconstructed forecast**: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC).


JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)




JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



JFM2022 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs


Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).


JFM2022 Anomaly Correlation for Individual AMIP Simulation with Observation — z200(20N–90N)




Observed & AMIP Ensemble Average Anomalies JFM2022 z200(m) 18 runs/worst 2 runs/best 2 runs




JFM2022 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N-90N)



Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 z200(m) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 z200(m) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast




JFM2022 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S-30N)

— ensemble mean


■ 0-1-d-L ■ 2-3-d-L ■ 4-5-d-L ■ 6-7-d-L ■ 8-9-d-L



Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast







Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast




JFM2022 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S-30N)

— ensemble mean


0-1-d-L 2-3-d-L 4-5-d-L 6-7-d-L 8-9-d-L



Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 T2m(K) 40 runs/worst 4 runs/best 4 runs Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies JFM2022 T2m(K) 40 runs/worst 4 runs/best 4 runs Reconstructed Forecast



z200(m) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) JFM2022 z200(m) eddy & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

T2m(k) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) JFM2022 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

Prec(mm/day) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) JFM2022 Prec(mm/day) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

z200(m) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst JFM2022 z200(m) eddy & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

T2m(k) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst JFM2022 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

Prec(mm/day) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst JFM2022 Prec(mm/day) & Obs



Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

Seasonal Forecasts from Multi-Model Ensemble Systems

- WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME). <https://www.wmolc.org/>
- Copernicus Climate Change Service (C3S) Multi-model seasonal forecasts. https://climate.copernicus.eu/charts/c3s_seasonal/
- North American Multi-Model Ensemble (NMME) seasonal forecasts. <https://www.cpc.ncep.noaa.gov/products/NMME/>

LC-LRFMM Seasonal Forecasts


(<https://www.wmolc.org/>)

Ensemble means

Simple Composite Map

Beijing,CMCC,CPTec,ECMWF,Exeter,Melbourne,Monreal,Moscow,Offenbach,Pretoria,Seoul,Tokyo,Toulouse,Washington


[Unit : K]



Simple Composite Map

Beijing,CMCC,CPTec,ECMWF,Exeter,Melbourne,Monreal,Moscow,Offenbach,Pretoria,Seoul,Tokyo,Toulouse,Washington

[Unit : mm]




Probabilities

Probabilistic Multi-Model Ensemble Forecast

Beijing,CMCC,CPTec,ECMWF,Exeter,Melbourne,Monreal,Moscow,Offenbach,Pretoria,Seoul,Tokyo,Toulouse,Washington


(issued on Dec2021)



Probabilistic Multi-Model Ensemble Forecast

Beijing,CMCC,CPTec,ECMWF,Exeter,Melbourne,Monreal,Moscow,Offenbach,Pretoria,Seoul,Tokyo,Toulouse,Washington


(issued on Dec2021)




C3S Seasonal Forecast

(https://climate.copernicus.eu/charts/c3s_seasonal/)

C3S multi-system seasonal forecast ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA/ECCC
Mean 2m temperature anomaly JFM 2022
Nominal forecast start: 01/12/21
Variance-standardized mean




C3S multi-system seasonal forecast ECMWF/Met Office/Météo-France/CMCC/DWD/NCEP/JMA/ECCC
Mean precipitation anomaly JFM 2022
Nominal forecast start: 01/12/21
Variance-standardized mean




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)

NMME Forecast of TMP2m Anom IC=202112 for Lead 1 2022JFM





NMME Forecast of Prec. rate Anom IC=202112 for Lead 1 2022JFM



200mb Height from Linear Model


JFM2022 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S-15N




Pattern COR: global=0.25, tropics(30S-30N)=0.46

Seasonal Forecasts from the Constructed Analog Model


CA HGT200 Prd for JFM2022, ICs through Mar2022(m), Lead -3



CA T2m Prd for JFM2022, ICs through Mar2022(K), Lead -3



CA Prec Prd for JFM2022, ICs through Mar2022(mm/day), Lead -3



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: OI version 2 analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation from CFSv2 (provided by Dr. Bhaskar Jha/CPC)
 - 18 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model (provided by Dr. Peitao Peng/CPC)
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model (provided by Dr. Peitao Peng/CPC)