

Attribution of Seasonal Climate Anomalies

February-March-April 2025

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)

Summary of Observed Conditions and Outlooks


- During FMA 2025, weak below-normal SST anomalies persisted in the central-eastern equatorial Pacific, consistent with weak La Niña conditions, while above-normal SSTs developed in the eastern tropical Pacific and along the coasts of Central and South America. Warm anomalies also persisted in the southern Indian Ocean, the northwestern and southern Pacific, and both tropical and North Atlantic basins, with the tropical Atlantic remaining anomalously warm for over a year (slide 4).
- CFSv2 forecasts captured the large-scale SST anomaly structure but exhibited a cold bias across the eastern Pacific and South American coastal regions (slide 10).
- AMIP, CFSv2, and MME forecasts consistently predicted enhanced precipitation over the Maritime Continent and the southwestern Pacific, and suppressed rainfall across the central-eastern equatorial Pacific—broadly aligning with observed tropical precipitation patterns (Slide 11, 37-39).
- Although the cold SST anomalies in the central Pacific were modest on an absolute basis, models simulated a [canonical La Niña response](#), likely due to [relatively cooler SST in the central Pacific](#) in a general warming background state of tropical and global oceans.
- CFSv2 forecasts reproduced the general warming trend in 200-mb height and land surface temperature but predicted a North Pacific blocking high resembling La Niña responses, deviating from observations and leading to erroneous predictions of strong cold anomalies over northwestern North America. (slides 12, 13, 15, 16).
- The predicted precipitation anomaly pattern over North America was consistent with the [La Niña response](#) and matched observations reasonably well (slide 14).
- Forecast skill for 200-mb height, T2m, and precipitation over North America improved at shorter lead times in April 2025 (slides 33-35).

Observed Seasonal Anomalies


Global and North America

Observed Anomaly FMA2025

SST(K)




Observed Anomaly FMA2025




Observed Anomaly FMA2025


Prec(mm/day)




T2m(K)



z200(m) (full anomalies)




z200(m) (eddy anomalies)




CPC Seasonal Outlooks and NMME Forecasts

CPC




Seasonal Temperature Outlook

Valid: Feb-Mar-Apr 2025
Issued: January 16, 2025




Seasonal Precipitation Outlook


Valid: Feb-Mar-Apr 2025
Issued: January 16, 2025



Prec nonEC
HSS=38




NMME prob fcst TMP2m IC=202501 for lead 1 2025 FMA



NMME

NMME prob fcst Prate IC=202501 for lead 1 2025 FMA




For the rationale behind CPC outlooks see https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2025/202501_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies


Model Simulated/Forecast Ensemble Average Anomalies

- AMIP simulations forced with observed sea surface temperatures (100 members ensemble)
- CFSv2 real time operational forecasts
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC). “w/o mean” is AC with area mean removed.


FMA2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




FMA2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




FMA2025 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)




FMA2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




FMA2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



FMA2025 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)



FMA2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs


Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).


FMA2025 Anomaly Correlation for Individual AMIP Simulation
with Observation -- z200(20N–90N)



Observed & AMIP Ensemble Mean Anomalies
FMA2025 z200(m) 100 runs/worst 4 runs/best 4 runs
(full anomalies: shaded; eddy anomalies: contours)





FMA2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N–90N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2025 z200(m) 40 runs/worst 4 runs/best 4 runs

Seasonal Forecast



(full anomalies: shaded; eddy anomalies: contours)




Observed & CFSv2 Forecast Ensemble Average Anomalies

Reconstructed Forecast


(full anomalies: shaded; eddy anomalies: contours)




FMA2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2025 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2025 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast




FMA2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2025 T2m(K) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
FMA2025 T2m(K) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast



$z200(m)$ Monthly Means from *Seasonal Forecast*

Monthly Means from Seasonal Fcst (40ensm) FMA2025 $z200(m)$ eddy & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

T2m(k) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) FMA2025 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th – 20th of the prior month.
- 4th row: 1st – 10th of the prior month.

Prec(mm/day) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) FMA2025 Prec(mm/day) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

$z200(m)$ Monthly Means from Monthly Forecast




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

T2m(k) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst FMA2025 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

Prec(/mm/day) Monthly Means from Monthly Forecast



Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.



Seasonal Forecasts from Multi-Model Ensemble Systems

- WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME).
<https://www.wmclc.org/>
- Copernicus Climate Change Service (C3S) Multi-model seasonal forecasts.
https://climate.copernicus.eu/charts/c3s_seasonal/
- North American Multi-Model Ensemble (NMME) seasonal forecasts.
<https://www.cpc.ncep.noaa.gov/products/NMME/>



LC-LRFMM Seasonal Forecasts

(<https://www.wmorc.org/>)

Ensemble means




Probabilities




C3S Seasonal Forecast

(https://climate.copernicus.eu/charts/c3s_seasonal/)




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)




200mb Height from Linear Model

FMA2025 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N



Seasonal Forecasts from the Constructed Analog Model



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: OI version 2 analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation based on GFS_FV3 (provided by Dr. Tao Zhang/CPC)
 - 100 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model.
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model.