

Attribution of Seasonal Climate Anomalies

March-April-May 2025

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)

Summary of Observed Conditions and Outlooks


- Based on FMA 2025 values, weak below-normal SST anomalies in the central-eastern Pacific continued to diminish. However, a pronounced east-to-west SST gradient--characteristic of La Niña events--remained. Warm anomalies in the eastern tropical Pacific and along the coasts of Central and South America also weakened. Equatorial Atlantic warming saw a slight decline as well. Meanwhile, positive SST anomalies persisted in the southern Indian Ocean, the northwestern and southern Pacific, and both the Northern and Southern Atlantic (slide 4).
- The CFSv2 forecast successfully captured the large-scale SST anomaly structure but exhibited a cold bias in its forecast for the eastern Pacific along the South American coast, the southern Indian Ocean, and the tropical Atlantic (slide 10).
- AMIP, CFSv2, and MME forecasts consistently showed enhanced rainfall over the Maritime Continent and the western/southwestern Pacific, with suppressed rainfall across the central-eastern equatorial Pacific – resembling La Niña responses. However, models overpredicted wet anomalies in the western tropical Pacific and northwest South America (Slide 11, 37-39).
- Despite weak central Pacific cold SST anomalies in absolute terms, models simulated a [canonical La Niña response](#), likely due to [relatively cooler SSTs in the central Pacific](#) against a background of warming oceans.
- CFSv2 captured the general warming trend in 200-mb height and land surface temperature but failed to forecast observed cold anomalies in North America due to misplacement of negative height anomalies from Southwest to Northeast (slides 12, 13, 15, 16).
- Predicted North American precipitation largely matched typical [La Niña signals](#) but diverged from observations (slide 14).
- May 2025 forecast skill for 200-mb height, T2m, and precipitation over North America improved with shorter leads (slides 33-35).

Observed Seasonal Anomalies


Global and North America

Observed Anomaly MAM2025


SST(K)



Observed Anomaly MAM2025




Observed Anomaly MAM2025




CPC Seasonal Outlooks and NMME Forecasts

CPC




Seasonal Temperature Outlook

Valid: Mar-Apr-May 2025
Issued: February 20, 2025




Seasonal Precipitation Outlook


Valid: Mar-Apr-May 2025
Issued: February 20, 2025



Prec nonEC
HSS=0




NMME prob fcst TMP2m IC=202502 for lead 1 2025 MAM



NMME

NMME prob fcst Rate IC=202502 for lead 1 2025 MAM




For the rationale behind CPC outlooks see https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2025/202502_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies


Model Simulated/Forecast Ensemble Average Anomalies

- AMIP simulations forced with observed sea surface temperatures (100 members ensemble)
- CFSv2 real time operational forecasts
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC). “w/o mean” is AC with area mean removed.


MAM2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




MAM2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




MAM2025 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)




MAM2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




MAM2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



MAM2025 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)



MAM2025 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs


Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).


MAM2025 Anomaly Correlation for Individual AMIP Simulation
with Observation -- $z200(20N-90N)$



Observed & AMIP Ensemble Mean Anomalies
MAM2025 z200(m) 100 runs/worst 4 runs/best 4 runs
(full anomalies: shaded; eddy anomalies: contours)



MAM2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N–90N)





Observed & CFSv2 Forecast Ensemble Average Anomalies

MAM2025 z200(m) 40 runs/worst 4 runs/best 4 runs

Seasonal Forecast

(full anomalies: shaded; eddy anomalies: contours)





Observed & CFSv2 Forecast Ensemble Average Anomalies


MAM2025 z200(m) 40 runs/worst 4 runs/best 4 runs

Reconstructed Forecast


(full anomalies: shaded; eddy anomalies: contours)




MAM2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies
MAM2025 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
MAM2025 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast




MAM2025 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
MAM2025 T2m(K) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
MAM2025 T2m(K) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast



$z200(m)$ Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) MAM2025 $z200(m)$ eddy & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

T2m(k) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) MAM2025 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

Prec(mm/day) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) MAM2025 Prec(mm/day) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

$z200(m)$ Monthly Means from Monthly Forecast




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

T2m(k) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst MAM2025 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

Prec(/mm/day) Monthly Means from Monthly Forecast



Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.



Seasonal Forecasts from Multi-Model Ensemble Systems

- WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME).
<https://www.wmclc.org/>
- Copernicus Climate Change Service (C3S) Multi-model seasonal forecasts.
https://climate.copernicus.eu/charts/c3s_seasonal/
- North American Multi-Model Ensemble (NMME) seasonal forecasts.
<https://www.cpc.ncep.noaa.gov/products/NMME/>



LC-LRFMM Seasonal Forecasts

(<https://www.wmorc.org/>)

Ensemble means





Probabilities




C3S Seasonal Forecast

(https://climate.copernicus.eu/charts/c3s_seasonal/)




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)




200mb Height from Linear Model

MAM2025 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N



Seasonal Forecasts from the Constructed Analog Model



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: OI version 2 analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation based on GFS_FV3 (provided by Dr. Tao Zhang/CPC)
 - 100 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model.
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model.