



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
February 11, 2008**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**




Overview

- **A moderate-strong MJO continues with the enhanced phase centered over the Maritime Continent.**
- **The MJO is expected to continue shifting eastward over the next 1-2 weeks to the western Pacific and the South Pacific Convergence Zone (SPCZ).**
- **Enhanced rainfall from Indonesia to the western Pacific including the SPCZ can be expected over the period. Tropical cyclogenesis is favored for the eastern Indian Ocean and waters to the north of Australia (Week 1) and to the northeast of Australia during Week 2.**
- **As MJO convection shifts eastward, an extension of the Pacific Jet Stream is more likely favoring increased chances for elevated rainfall across the northern US West Coast (mid-late Week 2). Given continued MJO propagation, circulation changes favor below-average temperatures across the eastern US during weeks 2-3.**



850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the magnitude of anomalous wind vectors



Strong northerly flow developed across the Arabian Sea during the past week.


Strong westerly anomalies are evident across the Indian Ocean during the last five days - in part associated with tropical cyclone activity.

Easterly anomalies continue across the western Pacific during the last five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Time



Longitude

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow

Weak intraseasonal activity was evident from August to October.


Westerly anomalies shifted eastward from the Indian Ocean to the Date Line during the first cycle of the recent MJO activity in November.

During December, the second MJO cycle is evident as westerly anomalies again developed across the Indian Ocean and shifted eastward while easterly anomalies strengthened in the western and central Pacific.

The next MJO cycle is evident as westerly anomalies have propagated eastward from the Indian Ocean to Indonesia with renewed easterly anomalies just west of the Date Line.



OLR Anomalies: Last 30 days



Drier-than-normal conditions, positive OLR anomalies (red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)


Wet (dry) conditions were observed across the western Pacific (Indian Ocean/western Maritime continent) during mid January.

As the MJO propagated eastwards, enhanced rainfall once again developed across the Indian Ocean and later Indonesia by early February. Dry conditions shifted across the Maritime continent to the western Pacific over the period.

Enhanced convection is evident near Hawaii during early February.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

Intraseasonal variability was evident during September and October but was less coherent and had a longer period than the current MJO activity.

Moderate-to-strong MJO activity has been evident since mid-November. Enhanced convection shifted from the Indian Ocean to the southwest Pacific during December and January while suppressed convection shifted from Africa to the Pacific.


During the past week, enhanced convection has shifted eastward to the Maritime continent.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation




The global velocity potential pattern indicates a coherent wave 1 structure typical of the MJO in the current phase. Large-scale suppressed (enhanced) conditions are evident across much of the western (eastern) hemisphere.



200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation
Negative anomalies (green shading) indicate favorable conditions for precipitation

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



The MJO was weak or incoherent during much of August and September.

The MJO strengthened during October but coherent propagation was short-lived.


Moderate-to-strong MJO activity developed in mid-November and has continued into February.

The MJO did weaken somewhat in early January as velocity potential anomalies became less coherent but recently the MJO has become more organized once again.



200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the magnitude of anomalous wind vectors




Westerly anomalies continue across the Pacific Ocean and have shifted slightly eastward during the last five days.

Well-defined anti-cyclonic (A) and cyclonic (C) circulations are evident throughout the period and are consistent with the current phase of the MJO.



200-hPa Zonal Wind Anomalies (m s^{-1})



Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow


Cycle 1 of the ongoing MJO activity is most clearly evident in the upper-levels by eastward propagation of easterly anomalies globally from early November to mid-December.

MJO cycle 2 signal was somewhat weaker especially as it shifted across the central Pacific Ocean due to a strengthening La Nina base state.



Weekly Heat Content Evolution in the Equatorial Pacific

Time



Longitude

Beginning in February, negative heat content anomalies developed across the eastern equatorial Pacific and continued until June 2007.

Kelvin wave activity (downwelling phases indicated by dashed lines) has been observed since May and has affected the sub-surface temperature departures at varying levels across the Pacific Ocean. The strongest wave occurred during May and June.

During September and October, negative heat content anomalies increased markedly across the eastern Pacific Ocean.

Most recently, increasingly positive anomalies have developed across the western Pacific and are shifted eastward associated with the latest downwelling Kelvin wave.



MJO Index -- Information



- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present

Plots put current MJO activity in historical context



Ensemble GFS MJO Forecasts


Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GEFS ensemble mean predicts continued eastward propagation during the next 1-2 weeks.

Confidence is high during the first week as most ensemble members predict similar evolution.






Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)


Spatial map of OLR anomalies for the next 15 days



The GEFS indicates a moderate-strong MJO signal early with a decrease in amplitude during week 2. Enhanced rainfall is expected across much of the Maritime continent and the western Pacific. Dry conditions are anticipated across Africa and the Indian Ocean.

Eastward propagation is expected during the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa wind vectors for the next 20 days
(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical MJO forecast indicates moderate MJO activity during the upcoming 1-2 week period. Enhanced rainfall is expected across the Maritime continent and the western Pacific Ocean.


Dry conditions are forecast to enter the Indian Ocean over the period.





MJO Composites – Global Tropics

Precipitation Anomalies



850-hPa Wind Anomalies

