



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
December 28, 2009**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remained weak during the past week.**
- **The MJO index signal is related to higher frequency coherent tropical variability rather than a large-scale, long-lived coherent MJO signal.**
- **The MJO is expected to remain weak during the upcoming 1-2 week period.**
- **The MJO is not expected to contribute substantially to anomalous tropical rainfall during the period.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades:
Easterly anomalies

Red shades:
Westerly anomalies



Westerly anomalies have strengthened across parts of the equatorial Atlantic and central Africa.


Easterly wind anomalies persisted across much of the Maritime Continent and Indian Ocean.

Westerly wind anomalies persisted across the central and eastern Pacific during the last five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



From July into September, easterly (westerly) anomalies prevailed across the Indian Ocean (Indonesia) (blue and orange boxes).

Several westerly wind bursts (red circles) occurred during this period. The westerly wind bursts became more frequent and stronger during September and October.


Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November and were associated with the MJO.

During mid-late December, westerly anomalies strengthened across much of the eastern Pacific Ocean.



OLR Anomalies: Last 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)




During late November and early December, suppressed convection (red ovals) was evident over the Americas, the Maritime Continent and Australia, while enhanced convection associated with the MJO was evident across the Western Pacific.

During early-mid December, enhanced convection developed across the west-central Pacific while suppressed convection persisted across the Maritime continent.

During mid-to-late December, the enhanced convection across the central Pacific waned and drier-than-average conditions across the Maritime Continent relaxed. Enhanced convection was also evident across much of the Americas and central Africa.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

During most of September and October, generally enhanced (suppressed) convection was evident across the western Pacific (eastern Indian Ocean) (blue and orange boxes).

Beginning in late October, enhanced convection developed across Africa, shifted eastward across the Maritime Continent, and spread into the Pacific. Later, suppressed convection developed across the Indian Ocean and also spread eastward during late November and early December.

Most recently, the strongest anomalies are with enhanced convection near the Date Line, which has shifted slightly east. Some enhancement is also evident across the Indian Ocean.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

Anomaly intensity varied during September and early October but the overall pattern remained generally persistent with upper-level divergence (convergence) across the western Pacific (parts of Western Hemisphere) (red box).

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (blue box).


During December, the coherent MJO pattern weakened. Eastward propagation is again evident but is related to higher frequency tropical variability and not coherent MJO activity.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation


Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern does not indicate a structure consistent with a coherent MJO.



200-hPa Vector Wind Anomalies (m s^{-1})



Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

During the last five to ten days, strong easterly anomalies developed across the eastern and central Pacific (blue box). The wind field also indicates some mid-latitude influence into the Tropics.


Westerly anomalies (red shades) stretch across much of the globe in the northern Hemisphere subtropics.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westerly anomalies across the Indian Ocean had persisted for much of the period since July 2009 (vertical dashed black line).


In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the latest MJO activity.



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



During December 2008 – January 2009, negative heat content anomalies returned and then strengthened in the central and eastern equatorial Pacific as La Niña conditions redeveloped.

The negative anomalies weakened during January-March 2009, with anomalies becoming positive since late March.

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average (blue box).

Multiple Kelvin waves shifted eastward between August and December (last three dashed black lines).



MJO Index -- Information


- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods or when the ENSO signal is large.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates an increase in amplitude with very fast eastward propagation. This is consistent with higher frequency tropical variability and not a coherent MJO.



MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.
Plots put current MJO activity in historical context.**



Ensemble GFS (GEFS) MJO Forecast


Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS MJO index forecasts indicate an eastward propagating signal across the Indian Ocean into the Maritime Continent over the period.

Ensemble spread is low during Week-1 but increases during Week-2.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast shows suppressed convection (red shades) across the western Pacific early in the period with enhanced convection (blue shades) across the Indian Ocean. Enhanced convection develops across the Maritime Continent during Week-2.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


The statistical forecast indicates suppressed convection initially over the Maritime Continent through Week-1 while enhanced convection would be present across the Indian Ocean throughout most of the period.





MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)



850-hPa Wind Anomalies (Nov-Mar)

