



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
February 1, 2010**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO has weakened during the past week as enhanced convection has become more persistent over the central tropical Pacific Ocean, consistent with El Niño.**
- **Based on the latest observations and some MJO forecast tools, the MJO is expected to continue to weaken.**
- **The weakening MJO activity, however, is expected to contribute to suppressed rainfall across Indonesia and enhanced rainfall across the central Pacific. There also remains an elevated threat for tropical cyclogenesis in the south Pacific Ocean.**
- **The current weakening MJO activity would tend to reinforce El Niño related impacts over the US as the convection continues to remain enhanced in the central Pacific.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades:  
Easterly anomalies

Red shades:  
Westerly anomalies



Westerly anomalies diminished over the eastern Indian Ocean/Maritime Continent, while easterly anomalies increased across the western Indian Ocean.


Westerly anomalies increased across the central Pacific and expanded eastward.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time  
↓



Longitude


Several westerly wind bursts (red circles) occurred during the July to October period. The westerly wind bursts became more frequent and stronger during September and October.

Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November associated with the MJO.

The westerly (easterly) anomalies (red box) evident in the Indian (western Pacific) Ocean during late December and early January have shifted eastward during January.



# OLR Anomalies: Last 30 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**


During early January, enhanced convection shifted to the Maritime Continent while wet conditions were observed across Central America.

During mid January, enhanced convection shifted eastward into the western Pacific while suppressed convection developed in the Indian Ocean.

In late January, enhanced convection became centered over the Date Line while suppressed convection continued to shift eastward over the Maritime Continent.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of the Bureau of Meteorology (BOM) - Australia)**

For much of the October 2009 to January 2010 period, the MJO has been active.

The October to early December 2009 period saw two periods of suppressed convection shift eastward from the Indian Ocean into the western Pacific (dotted lines) and one episode of enhanced convection (dashed line).

After a brief break during mid-late December, enhanced convection developed in the Indian Ocean and shifted eastward to the western and central Pacific during mid to late January. An area of suppressed convection across the Indian Ocean and Maritime Continent followed.

Time



Longitude




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Longitude

Anomaly intensity varied during September and early October but the overall pattern remained generally persistent with upper-level divergence (convergence) across the western Pacific (parts of Western Hemisphere) (red box).

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (blue box).

During early-to-mid December, the coherent MJO pattern weakened. Eastward propagation evident during the second half of December was related to higher frequency tropical variability and not large-scale coherent MJO activity (orange box).

Eastward propagation associated with the MJO was again evident during early-mid January.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The anomalous velocity potential pattern indicates upper-level divergence over the central and eastern Pacific with upper-level convergence evident across parts of South America, the Indian Ocean and the Maritime Continent.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



During the last five days, westerly anomalies (red shades) increased across Africa and the Indian Ocean.

An anomalous anticyclonic circulation is evident north of the equator across the subtropical Pacific (blue box).




# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time  
↓



In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the MJO.

Easterly anomalies have dominated much of the central and eastern Pacific during the second half of December and January.


Most recently, westerly anomalies have developed over much of the western Hemisphere into the Indian Ocean.



# Weekly Heat Content Evolution in the Equatorial Pacific

EQ. Upper-Ocean Heat Anoms. (deg C)

Time  
↓



Longitude

The negative anomalies associated with La Nina weakened during January-March 2009, with anomalies becoming positive since late March.

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average (blue box).

Multiple Kelvin waves shifted eastward between August and January 2010 (last three dashed black lines).

West of the Date Line, some cooling is present in the upwelling phases of the most recent Kelvin waves.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the US CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, Submitted.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the past week, the enhanced phase of the MJO remained generally stationary across the west-central Pacific. El Nino conditions are significantly contributing to the observed signal.



# MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.**




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GFS MJO index forecasts indicate some eastward propagation into phase 8 over the period.








# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

**Spatial map of OLR anomalies for the next 15 days**



The GEFS ensemble mean forecast indicates enhanced convection (blue shades) across the west-central Pacific. Suppressed convection (red shades) is forecast to shift slightly eastward over the period.

**Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days**






# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical forecast indicates a more progressive signal that the dynamical model forecast with enhanced convection weakening across the Pacific Ocean and suppressed convection shifting from the Indian Ocean to the Maritime continent.





# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)

## 850-hPa Wind Anomalies (Nov-Mar)

