



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
February 22, 2010**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO is weak as enhanced convection remains persistent over the central tropical Pacific Ocean, consistent with El Nino.**
- **Based on the latest observations and forecasts of weak MJO activity by MJO index model forecasts, the MJO is expected to remain weak during the next 1-2 weeks.**
- **Impacts across the Tropics and U.S. are expected to be more consistent with El Nino.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades:  
Easterly anomalies

Red shades:  
Westerly anomalies



Easterly anomalies continued across the Maritime continent but have weakened over south central Africa.


Westerly anomalies continued from the Date Line across the eastern Pacific.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time  
↓



Longitude

Several westerly wind bursts (red circles) occurred during the July to October period. The westerly wind bursts became more frequent and stronger during September and October.


Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November associated with the MJO.

The westerly (easterly) anomalies (red box) evident in the Indian (western Pacific) Ocean during late December and early January shifted eastward during January but propagation diminished by February.



# OLR Anomalies: Last 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During mid-to-late January, enhanced convection continued near the Date Line while suppressed convection continued across the eastern Indian Ocean and expanded to include the Maritime continent.


In late January and early February, enhanced convection shifted slightly eastward while suppressed convection intensified over the Maritime continent.

Enhanced (suppressed) convection continued across the central Pacific (Maritime continent) regions during mid-February. Also, wetter-than-average conditions developed across the eastern Pacific, Mexico and the Gulf of Mexico.





# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

The MJO has been active for much of the October 2009 to January 2010 period.

The October to early December 2009 period saw two periods of suppressed convection shift eastward from the Indian Ocean into the western Pacific (dotted lines) and one episode of enhanced convection (dashed line).

After a brief break during mid-late December, enhanced convection developed in the Indian Ocean and shifted eastward to the western and central Pacific during mid to late January. An area of suppressed convection across the Indian Ocean and Maritime Continent followed.

During February these anomalous areas have become more stationary.




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



Longitude

Anomaly intensity varied during September and early October but the overall pattern remained generally persistent with upper-level divergence (convergence) across the western Pacific (parts of Western Hemisphere) (red box).

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (blue box).

During early-to-mid December, the coherent MJO pattern weakened.

Eastward propagation associated with the MJO was again evident during early-mid January. During February, the MJO became much less coherent and anomalies became more stationary in nature – especially upper-level convergence near 120E.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



Upper-level divergence is indicated over the eastern Indian Ocean and the central/eastern Pacific while upper-level convergence is focused across Africa, the eastern Maritime Continent and the western Pacific.




# 200-hPa Vector Wind Anomalies ( $m s^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



During the last five to ten days, westerly anomalies (red shades) continued across much of Africa, the Indian Ocean and the Maritime continent with only minor eastward movement.


The depiction of anomalous anticyclonic circulations both north and south of the equator across the sub-tropical Pacific ("A", blue boxes) weakened during the last five days.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the MJO.

Easterly anomalies dominated much of the central and eastern Pacific during the second half of December and January.


During February westerly anomalies developed over the Indian Ocean and Maritime continent but have shown only minor eastward propagation to date.





# Weekly Heat Content Evolution in the Equatorial Pacific

Time  
↓



Longitude

The negative anomalies associated with La Nina weakened during January-March 2009, with anomalies becoming positive since late March.

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average (blue box).

Multiple Kelvin waves shifted eastward between August and January 2010 (last three dashed black lines).

West of the Date Line, some below-average anomalies are present in the upwelling phases of the most recent Kelvin waves.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the US CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, Submitted.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During late January and early February, the enhanced phase of the MJO remained generally stationary across the west-central Pacific. Some eastward movement of the index is indicated for a brief time during early-mid February before weakening dramatically.





# MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.**



# Ensemble GFS (GEFS) MJO Forecast


**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

**RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days**

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GFS MJO index forecasts indicate a generally weak signal remaining in phases 6 and 7 with no eastward propagation.

El Nino conditions continue to contribute to the amplitude and non-steady behavior of the MJO index forecast.






# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast indicates little change in anomalies over the period, consistent with the MJO index plot on the previous slide.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical forecast indicates a weak MJO signal.








# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)



## 850-hPa Wind Anomalies (Nov-Mar)

