



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
March 15, 2010**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remains weak as anomalous convection across the Maritime continent and Pacific remains consistent with El Nino conditions.**
- **Some model MJO index forecasts do indicate an increase in the signal with propagation across the Indian Ocean during the period. However, these particular models have not verified well in recent weeks. Nevertheless, potential MJO strengthening will need to be monitored.**
- **Based on the latest observations and some MJO index forecasts, the MJO is expected to remain weak during the next 1-2 weeks.**
- **Impacts across the Tropics are more likely to be consistent with El Nino.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades:
Easterly anomalies

Red shades:
Westerly anomalies

Westerly anomalies over southern Asia and the Indian Ocean have ended and have been replaced by weak easterly anomalies in some areas.




Westerly anomalies continue in the western Pacific south of the equator and are partly linked to tropical cyclone activity.

Strong westerly anomalies are again evident across the eastern Pacific during the last five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westerly wind bursts (red circles) occurred during September and October.


Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November associated with the MJO.

Weaker and shorter-lived MJO activity was evident during January.

Winds during March have been close to average across much of the deep Tropics.



OLR Anomalies: Last 30 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)


Enhanced (suppressed) convection (blue/red ovals) continued across the central Pacific (Maritime continent) regions during mid February. Wet conditions were also evident from the eastern Pacific to the Caribbean.

During late February, enhanced convection continued across the eastern Pacific, Central America and parts of the Caribbean. Wetter-than-average conditions developed across much of Africa.

In early-to-mid March, wet conditions in the central Pacific expanded eastward south of the equator while suppressed convection continued across the Maritime continent.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

During October to early December 2009, two periods of suppressed convection shifted eastward from the Indian Ocean into the western Pacific (dotted lines) and one episode of enhanced convection (dashed line).

After a brief break during mid-late December, enhanced convection developed in the Indian Ocean and shifted eastward to the western and central Pacific during mid to late January. An area of suppressed convection across the Indian Ocean and Maritime Continent followed.

During February and March these anomalous areas have become stationary.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Anomaly intensity varied during September and early October but the overall pattern remained generally persistent with upper-level divergence (convergence) across the western Pacific (parts of Western Hemisphere) (red box).

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (blue box).

Eastward propagation associated with the MJO was again evident during early-mid January.


During February, the MJO weakened and anomalies became more stationary during February and incoherent on the intraseasonal time scale in March.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The anomalous velocity potential pattern is not coherent.

Upper-level convergence is strongest across the western Maritime continent while upper-level divergence is indicated near the Date Line.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



During the last five to ten days easterly anomalies have returned to parts of the central and eastern Pacific (red boxes) and northern South America while a strong subtropical jet continues across the western Hemisphere north of the equator.




200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time
↓



In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the MJO.

Easterly anomalies dominated much of the central and eastern Pacific during the second half of December and January.


During late January and February westerly anomalies were evident over the Indian Ocean and Maritime continent.

Longitude



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



Longitude

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average.

Multiple Kelvin waves shifted eastward between August and March 2010 (last four dashed black lines).

West of the Date Line, some below-average anomalies are now present.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, Submitted.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During late January, February and early March the signal has generally remained across phases 7, 8 and 1 with varying amplitude indicating weak or no MJO activity.

El Nino conditions contributed to some of the signal during this period.



MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.
Plots put current MJO activity in historical context.**



Ensemble GFS (GEFS) MJO Forecast


Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS forecasts indicate incoherent MJO activity during the next the 1-2 weeks.

El Nino conditions continue to contribute to some of the amplitude and non-steady behavior of the MJO index forecast.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast indicates little change in position of anomalies with a weakening signal over the period, consistent with the MJO index plot on the previous slide.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


The statistical forecast indicates a only weak MJO activity.





MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)



850-hPa Wind Anomalies (Nov-Mar)

