



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
March 29, 2010**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- The MJO continued to show signs of strengthening and weak MJO activity is indicated.
- Not all observational fields indicate MJO activity and so it is too early to determine whether this subseasonal variation will mature into a long-lived, coherent MJO event.
- Most dynamical model MJO index forecasts indicate a gradual weakening signal as it shifts across the Maritime continent into the western Pacific during the next two weeks.
- Rainfall across the Maritime continent associated with this activity may aid drought stricken areas and enhance rainfall across the western Pacific during Week-2 while drier-than average conditions are expected across parts of Africa over the period.

Additional potential impacts across the global tropics are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>





# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades:  
Easterly anomalies

Red shades:  
Westerly anomalies



Westerly anomalies expanded and strengthened in the Indian Ocean during the last five days.

Easterly anomalies continued near the Date Line and expanded slightly east during the last five days.

Westerly anomalies continued across the eastern Pacific and developed in the equatorial Atlantic during the last five days.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow


Westerly wind bursts (red circles) occurred during September and October.

Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November associated with the MJO.

Weaker and shorter-lived MJO activity was evident during January.

Winds during the first half of March have been close to average across much of the deep Tropics although most recently easterly anomalies have increased just west of the Date Line.

Time  
↓




Longitude



# OLR Anomalies: Last 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**




Enhanced (suppressed) convection (blue/red ovals) continued across the central Pacific (Maritime continent) regions during late February and early March. Wet conditions were also evident from the eastern Pacific to the Caribbean and across parts of Africa.

During mid-March, enhanced convection continued across the central and eastern Pacific with a less expansive area of suppressed convection over Indonesia.

In mid-to-late March, anomalous convection was more variable than in recent weeks with enhanced (suppressed) convection near the Date Line (Indonesia) reduced. Wet conditions developed over parts of the Indian Ocean while dry conditions were evident over much of South America.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of the Bureau of Meteorology (BOM) - Australia)**

During October to early December 2009, two periods of suppressed convection shifted eastward from the Indian Ocean into the western Pacific (dotted lines) and one episode of enhanced convection (dashed line).

After a brief break during mid-late December, enhanced convection developed in the Indian Ocean and shifted eastward to the western and central Pacific during mid to late January. An area of suppressed convection across the Indian Ocean and Maritime Continent followed.


During February and March these anomalous areas became stationary and most recently have weakened.



# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation  
Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



Longitude

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (red box).

Eastward propagation associated with the MJO was again evident during early-mid January.

In February, the MJO weakened and anomalies became more stationary and incoherent on the intraseasonal time scale during the first half of March.

Most recently, weak upper-level divergence has developed over most of the eastern hemisphere where some areas in this region had experienced upper-level convergence.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The anomalous velocity potential pattern is more coherent than in recent weeks. The anomalies are generally small, however. Upper-level convergence is indicated across the western hemisphere while upper-level convergence is indicated over the eastern hemisphere.




# 200-hPa Vector Wind Anomalies ( $m s^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




During the last five days westerly anomalies have shifted eastward from the Indian Ocean to the western Pacific (red boxes).



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the MJO.


Easterly anomalies dominated much of the central and eastern Pacific during the second half of December and January.

During March easterly anomalies have been evident over the eastern Pacific Ocean – a change from late January and early February.



# Weekly Heat Content Evolution in the Equatorial Pacific

Time  
↓



Longitude

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average.

Multiple Kelvin waves shifted eastward between August and March 2010 (last four dashed black lines).

West of the Date Line, some below-average anomalies are now present.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, Submitted.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates an eastward propagating signal during the past ten days with the center located across the western Maritime continent.



# MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.**




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GFS forecasts indicate a weak but eastward propagating signal during the next 1-2 weeks.








# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast indicates weak enhanced convection (blue shades) shifting from the eastern Indian Ocean to the western Pacific over the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical forecast indicates weak-to-moderate MJO activity with enhanced convection shifting east from the eastern Indian Ocean to the western Pacific over the next two weeks.





# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)

## 850-hPa Wind Anomalies (Nov-Mar)

