



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
May 17, 2010**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO weakened during the past week with the enhanced convective phase located in the Maritime continent. Eastward propagation slowed during the past week.**
- **The majority of dynamical model MJO index forecasts indicate further weakening of the MJO signal during the period. These forecasts have not verified well during the past week.**
- **Based on the latest observations and statistical MJO forecasts, weak MJO activity is forecast to continue during the next week. Uncertainty is high due to differences in forecast tools.**
- **The MJO is expected to contribute to enhanced rainfall across parts of the eastern Indian Ocean, Maritime Continent and India and drier-than-average conditions across central Africa.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>



850-hPa Vector Wind Anomalies (m s^{-1})


Note that shading denotes the zonal wind anomaly

Blue shades:
Easterly anomalies

Red shades:
Westerly anomalies

During the previous five days, a large area of easterly anomalies was evident from the Indian Ocean to the central Pacific.

A narrow area of westerly anomalies is now evident across the equatorial Indian Ocean and western Maritime continent.




Easterly anomalies have shifted eastward during the last five days to include much of the eastern Pacific.

Westerly anomalies weakened during the last five days across the Atlantic and Africa.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during November and early December associated with the MJO.

Weaker and shorter-lived MJO activity was evident during January.

Easterly anomalies have persisted in the west-central Pacific since mid-March (black box). Westerly anomalies (red oval) increased across the eastern Pacific during late April associated with the MJO and entered the Indian Ocean in mid May.



OLR Anomalies: Last 30 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)


In mid-to-late April, suppressed convection continued across central South America while enhanced convection was evident across the southern Maritime continent.

During late April and early May, suppressed convection developed over much of the west-central Pacific and continued over parts of Brazil. Enhanced convection developed over the eastern Pacific, Central America, central Africa and the western Indian Ocean.

In early to mid May, enhanced convection shifted east to the Indian Ocean and Maritime continent. Suppressed convection developed across the eastern Pacific, Central America, South America and Africa.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

MJO activity was evident during early December 2009 and again during January 2010.

The MJO was not active during February and March as anomalous convection was more persistent across the Maritime continent (suppressed) and west-central Pacific (enhanced).

Anomalies were small during the month of April.

Enhanced convection developed across the Indian Ocean in early May and has shifted eastward associated with MJO activity.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

Eastward propagation associated with the MJO was evident during November and again during early-mid January (red boxes).

During February and the first half of March, the MJO weakened and anomalies became more stationary and incoherent on the intraseasonal time scale (black boxes).

In mid-March, weak upper-level divergence (convergence) developed over Africa and the Indian Ocean (Maritime continent) and these anomalies propagated eastward.


In late April and early May, anomalies increased and eastward propagation is evident.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The current anomalous velocity potential pattern has degraded to a weak “Wave 2” pattern with large scale upper-level divergence located across the Indian Ocean/Maritime continent and northern South America. Upper-level convergence is evident over the western Pacific and Africa.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies across the tropical central Pacific have shifted eastward to the eastern Pacific and South America during the last five days (red boxes).




200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during November associated with the MJO.

Easterly anomalies dominated much of the central and eastern Pacific during the second half of December and most of January (black oval).


For several weeks, westerly anomalies have been in place in the central Pacific while easterlies have dominated the eastern Pacific (red box). Since early May, however, eastward propagation of westerly anomalies is evident across the Pacific.



Weekly Heat Content Evolution in the Equatorial Pacific

EQ. Upper-Ocean Heat Anoms. (deg C)

Time
↓



Longitude

From May 2009 through March 2010, heat content anomalies remained above-average for much of the period.

From November 2009 – February 2010 three ocean Kelvin waves contributed to the change in heat content across the eastern Pacific (last three dashed black lines).

During April 2010 heat content anomalies have decreased across the Pacific in association with the upwelling phase of a Kelvin wave. Currently, negative heat content anomalies extend across the central and east-central Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, Accepted.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the past week, the MJO index indicates a decrease in eastward propagation. The signal remains centered across the Maritime continent region.



MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.
Plots put current MJO activity in historical context.**




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS forecasts indicate a weakening MJO signal during the period. Similar forecasts of a decrease in amplitude during the past week did not verify well.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast indicates enhanced (suppressed) convection over the Indian Ocean and Maritime continent (Central America and Africa) early in the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





Statistical MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical forecast is consistent with the GFS early in the period but indicates continued propagation.


This forecast shows enhanced convection from the Indian Ocean to the western Pacific during Week-1. Suppressed convection is forecast to enter the Indian Ocean by Week-2.





MJO Composites – Global Tropics

Precipitation Anomalies (May-Sep)



850-hPa Wind Anomalies (May-Sep)

