



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
October 4, 2010**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO signal was generally weak during the past week but recently has showed signs of strengthening.**
- **Dynamical models forecast a strengthening MJO over the next 2 weeks.**
- **Based on the latest observations and dynamical forecasts, the MJO signal is expected to further strengthen over the next 2 weeks. It is uncertain whether this potential activity will emerge as a more coherent long lived MJO event.**
- **The MJO is expected to contribute to wet conditions in the western Pacific and Indonesia and dry conditions in Central America.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies persisted across the western Pacific during the last five to ten days.


Westerly anomalies continued across the eastern Pacific but weakened in the Atlantic.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Easterly anomalies have persisted in the west-central Pacific since mid-March (black box) consistent with the development of La Nina conditions.


Enhanced westerly anomalies (red dotted ovals) occurred across the eastern Pacific on separate occasions during late April, late May and early-to-mid July and these were in part associated with MJO activity.

During mid-August, westerly anomalies strengthened at 90E, enhancing low-level convergence over the western Maritime Continent.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**




In early September, conditions were wetter-than-average across parts of southern Asia, Indonesia, and western Africa. Suppressed convection persisted near the Date Line.

Enhanced convection continued over west Africa during mid September and suppressed convection continued across the west-central Pacific.

During the end of September, enhanced convection expanded over the Indian Ocean and the western Maritime Continent and developed over Central America and the Caribbean Sea. Suppressed rainfall persisted across the west-central Pacific.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

Enhanced convection, in part associated with the MJO, developed across the Indian Ocean in early May and shifted eastward. Suppressed convection developed across much of Africa in its wake.

Since mid-July, generally enhanced (suppressed) convection has prevailed across the western Maritime continent (Date Line) (red box).

Considerable intraseasonal variability is evident during the period as enhanced convection has shifted both eastward and westward in this area during the period.




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



In late April and into May, anomalies increased and eastward propagation was evident, coincident with the MJO.

From early June to early July, anomalies became more stationary in nature (red box) with upper-level convergence primarily located across the central Pacific and divergence stretching from the Atlantic to the Indian Ocean.

Eastward propagation was evident during mid-July associated with the MJO.

Eastward propagation in August and September has mainly been associated with higher frequency coherent tropical variability rather than the MJO.


Longitude



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The anomalous velocity potential pattern shows upper-level divergence from Africa to eastern Asia and upper-level convergence across the Pacific. This pattern has become more organized over the past few days and exhibits a wave-1 structure, consistent with an MJO signal.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies strengthened over the Maritime continent (blue box). Easterly anomalies continued across the Atlantic Ocean (red box).



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

Westerly (easterly) anomalies prevailed across the central (eastern) Pacific (red box) for much of the period during March and April (black dotted box).


In early May, there was some eastward propagation of westerly anomalies across the Pacific in association with the MJO at that time (dashed black line).

Westerly anomalies have persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since early July.

Eastward propagation of westerly anomalies in August and September are not associated with the MJO.



# Weekly Heat Content Evolution in the Equatorial Pacific



From Aug 2009 through March 2010, heat content anomalies remained above-average for much of the period.

From November 2009 – February 2010 three ocean Kelvin waves contributed to the change in heat content across the eastern Pacific (last three dashed black lines).

During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Currently, negative heat content anomalies extend across the central and eastern Pacific with positive anomalies in the western Pacific.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, In Press.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the past week, the MJO index has indicated a strengthening signal.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GFS forecasts indicate a strengthening MJO signal during the next few days and eastward propagation by the end of Week 1 and through Week 2. Model spread is low increasing confidence in this forecast.







# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

**Spatial map of OLR anomalies for the next 15 days**

**Time-longitude section of (7.5 S-7.5 N) OLR anomalies for the last 180 days and for the next 15 days**



The GEFS ensemble mean forecast indicates a moderate signal of enhanced convection over the Indian Ocean and Maritime Continent during Week 1. During week 2, a strong signal of enhanced convection is forecast over the Maritime Continent and western Pacific and suppressed convection is forecast over Central America.




# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

Weak MJO activity is forecast during the period.





# MJO Composites – Global Tropics

## Precipitation Anomalies (May-Sep)

## 850-hPa Wind Anomalies (May-Sep)

