



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
May 23, 2011**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO signal weakened during the past week, but the index did indicate some eastward propagation, albeit at a slower pace than during most of this current event.**
- **The dynamical model MJO index forecasts indicate continued eastward propagation, but a significant weakening of the signal is forecast to continue during the period.**
- **Based on the latest observations and majority of model forecasts, the MJO is not expected to contribute significantly to rainfall anomalies across the Tropics.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies persisted across eastern equatorial Africa and the Indian Ocean during the last five day, with a slight expansion northward.


Westerly anomalies decreased in coverage across the western Pacific while easterly anomalies expanded near the Date Line.

Westerly anomalies persisted across the eastern Pacific Ocean and most of the tropical Atlantic Basin with a slight increase in magnitude.



850-hPa Zonal Wind Anomalies ($m s^{-1}$)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

Easterly anomalies have persisted in the west-central Pacific since October (black box) consistent with La Nina conditions.

In late January, easterly winds weakened and westerly anomalies developed near the Date Line due to MJO activity.


During March, easterlies strengthened near the Date Line, while westerly wind anomalies increased dramatically in strength at the end of the month.

A recent burst of westerly winds associated with the MJO moved across the Pacific. Most recently, easterly anomalies have returned to the western Pacific.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During mid-to-late April, enhanced convection continued from the Indian Ocean to the Western Pacific while suppressed convection continued across Africa and the southern CONUS.

The enhanced convection pattern over parts of the Eastern Hemisphere strengthened during early May. Suppressed convection persisted across Africa and the western Indian Ocean.


During mid-May, enhanced convection developed across much of the tropical Pacific Basin and parts of Central America and northern South America. The area of suppressed convection extended eastward from Africa to cover the entire Indian Ocean.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
5-Dec-2010 to 22-May-2011 + 14 days

Time
↓



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

MJO activity was experienced during late November into December and once again during January. During both periods, enhanced convection developed near 80E and shifted to the Maritime continent followed by an area of suppressed convection.

Enhanced convection was evident across northern South America during much of February and March.

During late March and April, two periods of distinct eastward movement of enhanced convection are noted. Most recently, an area of suppressed convection developed near 80E while the enhanced phase of the weakening MJO is evident near the Prime Meridian.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



During late November and early December, some eastward propagation associated with the MJO is evident in velocity potential anomalies.

During mid-to-late January, the MJO strengthened and upper-level divergence shifted eastward from 120E and upper-level convergence shifted from Africa to near the Date Line.

Eastward propagation of anomalies was observed during March associated with weak MJO activity.


Most recently during May, evidence of MJO activity with a fast period can be seen. Upper-level divergence (convergence) is observed across much of the Western Hemisphere (Indian Ocean).



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows anomalous upper-level divergence from portions of northern South America, across Africa to the western Indian Ocean. Anomalous upper-level convergence is evident from the eastern Indian Ocean to the central Pacific Ocean.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies persisted across the equatorial central Pacific and Indian Ocean during the last five to ten days.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since November.

Some strengthening and eastward propagation of westerly anomalies was evident in late March. In early April there was some eastward propagation of an easterly/westerly anomaly couplet over the eastern Hemisphere.

Significant eastward propagation was evident in May (dashed line) associated with the MJO.



Weekly Heat Content Evolution in the Equatorial Pacific



Beginning in April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Since the beginning of January 2011, positive heat content anomalies have shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution



- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months



The MJO index indicates the continued progression of the MJO signal with some weakening during the last week.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present. Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecasts a continued weakening of the signal during most of Week-1, with a weak strengthening during later Week-1. The signal remains weak overall during the entire two week period.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast indicates a weak signal supporting continued dry conditions from the Indian Ocean to the western Pacific early in the period. Later in Week-1, some enhanced convection is indicated across the western Pacific

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


The forecast calls for little to no MJO activity during the period.






MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)



850-hPa Wind Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>