



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
January 30, 2012**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO strengthened during the past week with its enhanced phase, as indicated by the MJO index, located across the western Pacific.**
- **Dynamical model MJO index forecasts show the MJO signal continuing through the period. There is large spread amongst the models for the magnitude and eventual phase by the end of Week-2. Based on the latest observations and model forecasts, the MJO is forecast to be active during the upcoming 1-2 weeks.**
- **The MJO is forecast to contribute to enhanced convection across parts of the western Pacific, northeast Australia and the South Pacific Convergence Zone (SPCZ) during the next two weeks. Suppressed convection is favored for the Indian Ocean over the period.**
- **For the U.S., the current and forecast phases of the MJO favor a tendency for a positive PNA pattern in coming weeks with enhanced chances for below-normal temperatures in the eastern U.S..**
- **Later as the MJO enters the western hemisphere, there is a tendency for stronger jet streams to develop. There is potential for a ridge in the west to be undercut, returning wet conditions to the west, and also enhanced moisture is more likely across the southern tier of the U.S.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly wind anomalies strengthened and shifted eastward over the Maritime Continent during the past five days.


Easterly anomalies strengthened over the equatorial Pacific.

Westerly wind anomalies continued over the east Pacific Ocean.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

In early October, MJO activity weakened the persistent easterly anomalies across the central Pacific (first dashed line).


MJO activity continued into December (altering dashed and dotted lines), but most recently westerly (easterly) wind anomalies across the Indian Ocean (western Pacific) have become more stationary.

In mid-January, interaction with the extratropics contributed to decreased easterlies near the Date Line.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During the end of December, enhanced convection (blue circle) developed near the Philippines, the Maritime continent and the South Pacific Convergence Zone (SPCZ). Suppressed convection (red circle) developed in the equatorial Indian Ocean.


Enhanced (suppressed) convection continued into mid-January around the western Maritime Continent (equatorial Indian Ocean).

During mid-to-late January, convection markedly increased for a large area over the Indian Ocean and Maritime continent.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
14-Aug-2011 to 29-Jan-2012 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

**Beginning in mid-September, enhanced convection shifted from southern Asia to the western Pacific while suppressed convection developed during late September across India and also shifted eastward to the western Pacific.**

**MJO activity continued into early December, then OLR anomalies decreased and eastward propagation was not clear but during late December, eastward propagation of OLR anomalies was again observed.**

**Strong enhanced convection developed across the Indian Ocean during late January.**

**Longitude**




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



During July and August very fast eastward propagation was evident at times and mainly associated with higher frequency sub-seasonal coherent tropical variability not associated with MJO activity.

Beginning in the second half of September into December, alternating negative (dashed lines) and positive (dotted lines) anomalies were evident and associated with MJO activity during the period.

Eastward propagation of anomalies became less coherent during January and anomalies became very weak.


Longitude



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The latest large scale velocity potential pattern is less coherent recently than a few days ago. Positive anomalies over North America and western Eurasia are strong, while upper-level divergence is weak across the Pacific.




# 200-hPa Vector Wind Anomalies ( $m s^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Large scale cyclonic circulations are evident both north and south of the equator east of the Date Line.


During the past five days, westerly anomalies strengthened over the eastern Pacific, while westerly anomalies weakened in the western Pacific.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) during much of the period prior to mid-September.

Alternating westerly (dashed lines) and easterly (dotted lines) anomalies are evident from mid-September into December associated with the MJO.

In December, westerly anomalies strengthened over the Pacific. Most recently, these westerly anomalies have shown some eastward propagation.



# Weekly Heat Content Evolution in the Equatorial Pacific



Since the beginning of January 2011, positive heat content anomalies shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011. Much of the Pacific basin now indicates above- or near-normal integrated heat content.

Since late July, negative heat content anomalies are evident across the equatorial central and eastern Pacific.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the first half of January the MJO index was fairly weak with no eastward propagation. Recently, the amplitude has increased over the Maritime continent and eastward propagation into the western Pacific is evident.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

**RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days**

**light gray shading:** 90% of forecasts  
**dark gray shading:** 50% of forecasts

The ensemble GFS forecasts the MJO signal to strengthen and propagate eastward with the enhanced phase shifting across the Pacific during Week-2.






# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecast indicates enhanced convection across the west-central Pacific, parts of Brazil, and parts of Africa during the next two weeks with suppressed convection across the Indian Ocean.







# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The CA forecast shows enhanced convection over the west-central Pacific, parts of Brazil, and parts of Africa while suppressed convection dominates the region from the central Indian Ocean to the western Maritime Continent.




# MJO Composites – Global Tropics

## 850-hPa Wind Anomalies (Nov-Mar)



## Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>